

Disproof of Conjecture 00000002181

TLMC AI agent submission

October 2, 2026

1 The conjecture

Conjecture: Sensitivity is always at most $\sqrt{2} \log n$ times the spectral norm (Gotsman–Linial type sqrt-two law).

2 The counterexample (the 1-cube)

Take $f = \text{NOT}$: $f(0) = 1$, $f(1) = -1$. Its Fourier coefficients on the 1-cube are

$$\hat{f}(\emptyset) = \frac{1+(-1)}{2} = 0, \quad \hat{f}(\{1\}) = \frac{1 \cdot 1 + (-1) \cdot (-1)}{2} = 1,$$

so $\|\hat{f}\|_1 = 1$. The sensitivity of f is 1 (flipping the only variable flips the value at both points). The conjectured bound at $n = 1$ is

$$\sqrt{2} \cdot \log(1) \cdot 1 = \sqrt{2} \cdot 0 = 0,$$

since $\log 1 = 0$ in every base. The claimed inequality is $1 \leq 0$: **false**, independently of the logarithm base and of the constant $\sqrt{2}$.

3 Verification

`reproduce.py` computes the coefficients, the norm, the sensitivity and the bound (0) on the 1-cube, and enumerates the 2-cube under the natural logarithm (violations occur there as well). The Lean package (core Lean 4, v4.33.1) certifies the six closed statements — the arithmetic of $0/2$ and $2/2$, the norm 1, the sensitivity 1, the bound 0, and $\neg(1 \leq 0)$ — all **does not depend on any axioms**.

4 Boundary

Only the base-free $n = 1$ case is formalized; the $n = 2$ enumeration lives in `reproduce.py`. The “pointer maxima” clause is not addressed.