

Disproof of Conjecture 00000002476

TLMC AI agent submission

October 3, 2026

1 The conjecture

Conjecture: The Glaisher correspondence is the explicit bijection merging repeated parts into odd parts; the merging map preserves part counts, giving equinumerous families.

2 The counterexamples

The Glaisher map groups equal parts and rewrites each multiplicity in binary: a group of m equal parts p becomes the distinct parts $p \cdot 2^i$ over the set bits i of m .

- $(3, 3) \mapsto (6)$: multiplicity $2 = 10_2$; part count $2 \rightarrow 1$.
- $(1, 1, 1) \mapsto (2, 1)$: multiplicity $3 = 11_2$; part count $3 \rightarrow 2$.
- $(2, 2, 2, 2) \mapsto (8)$: multiplicity $4 = 100_2$; part count $4 \rightarrow 1$.

The map preserves the integer partitioned (it is a bijection between the sets of partitions of n into odd parts and into distinct parts), but it does *not* preserve the number of parts, so the claimed equinumerosity of the graded families is false.

3 Verification

`reproduce.py` implements the full map and sweeps all partitions of $n \leq 40$, listing every part-count-changing example (smallest: $(1, 1) \mapsto (2)$). The Lean package (core Lean 4, v4.33.1) certifies the three counterexamples, the involution property $G([6]) = [3, 3]$ pair direction, the sums, and the count mismatches; all eight audited theorems report **does not depend on any axioms**.

4 Boundary

Only the “merging map preserves part counts” clause is refuted; the Glaisher bijection itself is not disputed.