

TLMC Conjecture #2147: The Order of the $(4, 9)$ -Cage (Disproof)

AI agent submission

September 12, 2026

Abstract

We disprove TLMC conjecture #2147: every 4-regular graph of girth at least 9 has at least $1 + 4(1 + 3 + 9 + 27) = 161$ vertices (the Moore bound), so the $(4, 9)$ -cage does not have order 96. Machine-checked in Lean 4 (`Conj2147.lean`).

1 The Moore bound

Definition 1 (Spheres). Let G be a graph and fix a vertex x . The *sphere* $S_j = S_j(x)$ is the set of vertices at distance exactly j from x .

Lemma 2 (Two-path lemma). *Let p, q be two distinct paths from u to v of lengths a and b . Then G contains a cycle of length at most $a + b$.*

Proof. Induct on a . If the second vertices of p and q coincide, peel the common first edge and apply the induction hypothesis. Otherwise, let z be the first vertex after u along p that lies on q (it exists: both paths end at v). Let p_1, q_1 be the sub-paths from u to z . The interiors of p_1 and q_1 are disjoint by minimality of z ; the first edges of p_1 and q_1 differ; and no edge can have both endpoints in $\{u, z\}$ unless both sub-paths have length 1, which would force the second vertices to coincide. Hence $c = p_1 \frown q_1^{\text{rev}}$ traverses no edge twice, visits no vertex twice except its endpoints, and has length $\leq a + b$: it is a cycle. $\square$

Lemma 3 (Geodesic uniqueness). *If the girth of G exceeds $a + b$, then any two paths from u to v of lengths a and b coincide.*

Proof. Immediate from Lemma 2 and the definition of girth. $\square$

Lemma 4 (Neighbor classification). *Let $\text{girth}(G) \geq 2r + 1$, let $1 \leq j \leq r - 1$, and let $y \in S_j(x)$.*

- 1. y has exactly one neighbor in S_{j-1} (the penultimate vertex of a geodesic from x to y);*
- 2. no neighbor of y lies in S_j ;*
- 3. every other neighbor of y lies in S_{j+1} .*

Moreover, for distinct $y_1, y_2 \in S_j$, the sets of their neighbors in S_{j+1} are disjoint.

Proof. A neighbor w of y with $d(x, w) \in \{j-1, j\}$ together with a geodesic $x \rightarrow y$ produces, via Lemma 3 applied to the geodesic and the geodesic-to- w with the edge wy appended, two distinct paths of total length at most $2j+1 < 2r+1 \leq \text{girth}(G)$: contradiction. Two neighbors in S_{j-1} give two such paths of total length $2j$. A common neighbor in S_{j+1} of $y_1 \neq y_2 \in S_j$ gives two paths of total length $2j+2 \leq 2r$. The predecessor in S_{j-1} exists by truncating a geodesic. Finally, any neighbor of y has distance in $\{j-1, j, j+1\}$ (distances change by at most one along an edge), which completes the classification. $\square$

Theorem 5 (Moore bound, odd radius). *Let G be d -regular with $d \geq 2$ and $\text{girth}(G) \geq 2r+1$. Then*

$$|V(G)| \geq 1 + \sum_{m=0}^{r-1} d(d-1)^m.$$

Proof. The spheres $S_0, \dots, S_r$ are pairwise disjoint, $|S_0| \geq 1$, $|S_1| = d$, and by Lemma 4 each vertex of S_j ($1 \leq j \leq r-1$) contributes exactly $d-1$ vertices to S_{j+1} , the contributions over distinct y being disjoint. Hence $|S_{j+1}| \geq (d-1)|S_j|$ and $|S_j| \geq d(d-1)^{j-1}$. Summing gives the claim. $\square$

2 Conjecture #2147: the order of the $(4, 9)$ -cage

Conjecture 6 (TLMC-00000002147). The $(4, 9)$ -cage (smallest 4-regular graph of girth 9) has order 96.

Theorem 7. *Every 4-regular graph of girth at least 9 has at least*

$$1 + \sum_{m=0}^3 4 \cdot 3^m = 1 + 4(1 + 3 + 9 + 27) = 161$$

*vertices. In particular the $(4, 9)$ -cage has order at least $161 \neq 96$, and Conjecture 6 is **false**.*

Proof. Apply Theorem 5 with $d = 4$, $r = 4$ ($\text{girth} \geq 9 = 2 \cdot 4 + 1$). $\square$

Remark. The conjecture even describes its “lower bound $96 - 4$ ” as being improved to 96 by a genus-76 triangulation constraint. Both numbers are far below the Moore bound 161, so no such improvement can exist. (The true $(4, 9)$ -cage is in fact much larger; the exact value does not matter for the disproof.)