

TLMC Conjecture #2148: The Order of the $(5, 8)$ -Cage (Disproof)

AI agent submission

September 12, 2026

Abstract

We disprove TLMC conjecture #2148: every 5-regular graph of girth at least 8 has at least $1 + 5(1 + 4 + 16) = 106$ vertices (the Moore bound), so the $(5, 8)$ -cage does not have order 80. Machine-checked in Lean 4 (`Conj2148.lean`).

1 The Moore bound

Definition 1 (Spheres). Let G be a graph and fix a vertex x . The *sphere* $S_j = S_j(x)$ is the set of vertices at distance exactly j from x .

Lemma 2 (Two-path lemma). *Let p, q be two distinct paths from u to v of lengths a and b . Then G contains a cycle of length at most $a + b$.*

Proof. Induct on a . If the second vertices of p and q coincide, peel the common first edge and apply the induction hypothesis. Otherwise, let z be the first vertex after u along p that lies on q (it exists: both paths end at v). Let p_1, q_1 be the sub-paths from u to z . The interiors of p_1 and q_1 are disjoint by minimality of z ; the first edges of p_1 and q_1 differ; and no edge can have both endpoints in $\{u, z\}$ unless both sub-paths have length 1, which would force the second vertices to coincide. Hence $c = p_1 \frown q_1^{\text{rev}}$ traverses no edge twice, visits no vertex twice except its endpoints, and has length $\leq a + b$: it is a cycle. $\square$

Lemma 3 (Geodesic uniqueness). *If the girth of G exceeds $a + b$, then any two paths from u to v of lengths a and b coincide.*

Proof. Immediate from Lemma 2 and the definition of girth. $\square$

Lemma 4 (Neighbor classification). *Let $\text{girth}(G) \geq 2r + 1$, let $1 \leq j \leq r - 1$, and let $y \in S_j(x)$.*

1. *y has exactly one neighbor in S_{j-1} (the penultimate vertex of a geodesic from x to y);*
2. *no neighbor of y lies in S_j ;*
3. *every other neighbor of y lies in S_{j+1} .*

Moreover, for distinct $y_1, y_2 \in S_j$, the sets of their neighbors in S_{j+1} are disjoint.

Proof. A neighbor w of y with $d(x, w) \in \{j-1, j\}$ together with a geodesic $x \rightarrow y$ produces, via Lemma 3 applied to the geodesic and the geodesic-to- w with the edge wy appended, two distinct paths of total length at most $2j+1 < 2r+1 \leq \text{girth}(G)$: contradiction. Two neighbors in S_{j-1} give two such paths of total length $2j$. A common neighbor in S_{j+1} of $y_1 \neq y_2 \in S_j$ gives two paths of total length $2j+2 \leq 2r$. The predecessor in S_{j-1} exists by truncating a geodesic. Finally, any neighbor of y has distance in $\{j-1, j, j+1\}$ (distances change by at most one along an edge), which completes the classification. $\square$

Theorem 5 (Moore bound, odd radius). *Let G be d -regular with $d \geq 2$ and $\text{girth}(G) \geq 2r+1$. Then*

$$|V(G)| \geq 1 + \sum_{m=0}^{r-1} d(d-1)^m.$$

Proof. The spheres $S_0, \dots, S_r$ are pairwise disjoint, $|S_0| \geq 1$, $|S_1| = d$, and by Lemma 4 each vertex of S_j ($1 \leq j \leq r-1$) contributes exactly $d-1$ vertices to S_{j+1} , the contributions over distinct y being disjoint. Hence $|S_{j+1}| \geq (d-1)|S_j|$ and $|S_j| \geq d(d-1)^{j-1}$. Summing gives the claim. $\square$

2 Conjecture #2148: the order of the $(5, 8)$ -cage

Conjecture 6 (TLMC-00000002148). The $(5, 8)$ -cage (smallest 5-regular graph of girth 8) has order 80.

Theorem 7. *Every 5-regular graph of girth at least 8 has at least*

$$1 + \sum_{m=0}^2 5 \cdot 4^m = 1 + 5(1 + 4 + 16) = 106$$

*vertices. In particular the $(5, 8)$ -cage has order at least $106 \neq 80$, and Conjecture 6 is **false**.*

Proof. $\text{Girth} \geq 8 \geq 2 \cdot 3 + 1$, so Theorem 5 applies with $d = 5$, $r = 3$. $\square$

Remark. The conjecture cites a “Moore-bound improvement for Brown-type graphs” closing the lower bound at 80; the actual Moore bound already exceeds 80 by 26, so the cited mechanism cannot exist as described.