

Disproof of Conjecture 00000001230: the Jacobian of the 2×2 grid is $\mathbb{Z}/4$, which is not a sum of $\mathbb{Z}/\gcd(S)$

jilint777

2026-10-04

Abstract

Conjecture 00000001230 asserts that the Jacobian (sandpile group) of the grid graph $[m_1] \times \cdots \times [m_n]$ is isomorphic to a group $\bigoplus_S (\mathbb{Z}/\gcd(S))^{e(S)}$, where S runs over the subsets of $\{m_1, \dots, m_n\}$. The exponents $e(S)$ are not specified. We show that no choice of exponents works. The 2×2 grid is the 4-cycle C_4 , and its Jacobian $\mathbb{Z}/4$ has an element of order 4. For $(m_1, m_2) = (2, 2)$, every modulus $\gcd(S)$ is 0 or 2. In any product of copies of $\mathbb{Z}$ and $\mathbb{Z}/2$, an element x with $4x = 0$ also has $2x = 0$. So $\mathbb{Z}/4$ does not even embed into any such group. The same graph defeats the edge-count convention $[m] = \text{path with } m \text{ edges}$, where the moduli are 0 and 1. The cube $[2]^3$ (Jacobian $\mathbb{Z}/2 \oplus \mathbb{Z}/8 \oplus \mathbb{Z}/24$) gives a counterexample in dimension 3. The disproof is formalized in Lean 4 (core library only), with the grid graph, its Laplacian, the sandpile group and the conjectured groups all defined from scratch.

1 The conjecture and its precise reading

Definition: $\text{Jac}(K_{m,n})$ is the sandpile group (Jacobian) of the complete bipartite graph.

Conjecture: The Jacobian of the n -dimensional grid graph $[m_1] \times \cdots \times [m_n]$ is isomorphic to the explicit decomposition $\bigoplus_S (\mathbb{Z}/\gcd(S))^{e(S)}$ over subsets S of the m_i ; the two-dimensional case is given explicitly by the $K_{m,n}$ formula and torus gluing.

The Chinese version says the same. The statement never defines $e(S)$, so we read it in the weakest possible form:

Claim. For every $n \geq 1$ and every $m_1, \dots, m_n \geq 1$ there exist exponents $e(S) \geq 0$, one for each subset S , such that $\text{Jac}([m_1] \times \cdots \times [m_n]) \cong \bigoplus_S (\mathbb{Z}/\gcd(S))^{e(S)}$.

Any explicit formula for $e(S)$, including whatever the second clause (“given explicitly by the $K_{m,n}$ formula and torus gluing”) is meant to produce in dimension 2, is a special case. Refuting the Claim therefore refutes the conjecture, whatever the exponents are meant to be. The second clause is too vague to state precisely. We do not need it, since the conjecture is a conjunction and the first clause already fails.

Conventions covered. The counterexample defeats all of the following readings at the same time (Section 5):

- $[m]$ is the path with m *vertices* (standard), or the path with m *edges*;

- $\gcd(\emptyset) = 0$ (so $\mathbb{Z}/\gcd(\emptyset) = \mathbb{Z}$), $\gcd(\emptyset) = 1$, or the empty set is skipped;
- “subsets of the m_i ” are subsets of the index set $\{1, \dots, n\}$ (so repeated values give repeated summands) or of the value set $\{m_1, \dots, m_n\}$;
- the exponents are finite, or arbitrary cardinals (infinite direct sums);
- “Jacobian” is the sandpile group $\mathbb{Z}^{V \setminus \{s\}} / \tilde{\Delta} \mathbb{Z}^{V \setminus \{s\}}$, the Baker–Norine Jacobian $\text{Pic}^0 = \text{Div}^0 / \text{Prin}$, or even the full cokernel $\mathbb{Z}^V / \Delta \mathbb{Z}^V$;
- dimension $n = 2$, or the claim restricted to $n \geq 3$.

2 Definitions

Definition 1. For $m_1, \dots, m_n \geq 1$, the grid graph $[m_1] \times \dots \times [m_n]$ has vertex set $\{(x_1, \dots, x_n) : 0 \leq x_i < m_i\}$. Two vertices are adjacent iff their ℓ^1 distance is 1, that is, they differ by 1 in exactly one coordinate. Equivalently, it is the Cartesian product $P_{m_1} \square \dots \square P_{m_n}$ of paths.

Definition 2. Let $G = (V, E)$ be a connected graph with Laplacian $\Delta = D - A$, and fix a sink $s \in V$. The *reduced Laplacian* $\tilde{\Delta}$ is Δ with the row and column of s deleted. The *sandpile group* (Jacobian, critical group) is

$$K(G) = \mathbb{Z}^{V \setminus \{s\}} / \tilde{\Delta} \mathbb{Z}^{V \setminus \{s\}}.$$

Its order is $\det \tilde{\Delta}$, the number of spanning trees (matrix-tree theorem). Up to isomorphism it does not depend on s . It is isomorphic to the Baker–Norine Jacobian $\text{Pic}^0(G) = \text{Div}^0(G) / \Delta \mathbb{Z}^V$, where Div^0 is the set of integer vectors with coordinate sum 0. Also $\text{Pic}(G) = \mathbb{Z}^V / \Delta \mathbb{Z}^V \cong \mathbb{Z} \oplus K(G)$.

Throughout, $\mathbb{Z}/0 = \mathbb{Z}$ and $\mathbb{Z}/1 = 0$.

3 The obstruction

Lemma 3. Let $(d_i)_{i \in I}$ be any family, indexed by any set I , with $d_i \in \{0, 1, 2\}$. Let $P = \prod_{i \in I} \mathbb{Z}/d_i$. If $x \in P$ and $4x = 0$, then $2x = 0$. The same holds in every subgroup of P , in particular in the direct sum $\bigoplus_{i \in I} \mathbb{Z}/d_i$.

Proof. Work coordinatewise. If $d_i = 0$, then $4x_i = 0$ in $\mathbb{Z}$ forces $x_i = 0$. If $d_i = 2$ or $d_i = 1$, then $2x_i = 0$ always. $\square$

Lemma 4. Let G be an abelian group with an element g such that $4g = 0$ and $2g \neq 0$. Then for every family as in Lemma 3 there is no injective homomorphism $G \rightarrow \prod_i \mathbb{Z}/d_i$. In particular $G \not\cong \bigoplus_i \mathbb{Z}/d_i$.

Proof. Let φ be an injective homomorphism. Then $4\varphi(g) = \varphi(4g) = 0$, so $2\varphi(g) = 0$ by Lemma 3. So $\varphi(2g) = 0 = \varphi(0)$, and injectivity gives $2g = 0$, a contradiction. $\square$

In Lean, a homomorphism is just a map φ with $\varphi(a + b) = \varphi(a) + \varphi(b)$, and $\varphi(0) = 0$ is derived from this. Lemma 3 also gives cancellation, $y + y = y \Rightarrow y = 0$, in $\prod_i \mathbb{Z}/d_i$.

4 The 2×2 grid

The grid $[2] \times [2]$ has vertices $(0, 0), (0, 1), (1, 0), (1, 1)$ and edges $(0, 0)(0, 1), (0, 0)(1, 0), (0, 1)(1, 1), (1, 0)(1, 1)$. It is the 4-cycle C_4 . Its Laplacian, in this vertex order, is

$$\Delta = \begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 2 & 0 & -1 \\ -1 & 0 & 2 & -1 \\ 0 & -1 & -1 & 2 \end{pmatrix}.$$

With sink $s = (0, 0)$, the reduced Laplacian is

$$\tilde{\Delta} = \begin{pmatrix} 2 & 0 & -1 \\ 0 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}, \quad \det \tilde{\Delta} = 4.$$

Proposition 5. *Let $g = [e_0] \in K(C_4)$ be the class of one chip on $(0, 1)$. Then $4g = 0$ and $2g \neq 0$. In fact $K(C_4) \cong \mathbb{Z}/4$, via $x \mapsto 3x_0 + x_1 + 2x_2 \pmod{4}$.*

Proof. $\tilde{\Delta}(3, 1, 2)^\top = (6 - 2, 2 - 2, -3 - 1 + 4)^\top = (4, 0, 0)^\top = 4e_0$, so $4g = 0$. Suppose $\tilde{\Delta}z = 2e_0$ with $z \in \mathbb{Z}^3$, that is, $2z_0 - z_2 = 2$, $2z_1 - z_2 = 0$ and $-z_0 - z_1 + 2z_2 = 0$. The second equation makes z_2 even, say $z_2 = 2a$. Then $z_1 = a$ and $z_0 = 1 + a$, and the third equation becomes $2a = 1$, which is impossible. So $2g \neq 0$.

For the isomorphism, the vector $(3, 1, 2)$ is the first row of $\text{adj } \tilde{\Delta} = 4\tilde{\Delta}^{-1}$. It annihilates every column of $\tilde{\Delta}$ modulo 4 (columns $(2, 0, -1), (0, 2, -1), (-1, -1, 2)$ give 4, 0, 0). So $\psi(x) = 3x_0 + x_1 + 2x_2 \pmod{4}$ is well defined on $K(C_4)$. It is onto, since $\psi(3a, 0, 0) = 9a \equiv a$. It is injective: if $w = x - y$ has $\psi(w) \equiv 0$, write $3w_0 + w_1 + 2w_2 = 4q$. Then $w_0 - w_1$ is even, say $2r$, and $z = (q, q - r, r + w_1 + w_2)$ is an integer solution of $\tilde{\Delta}z = w$. $\square$

Theorem 6. *For $(m_1, m_2) = (2, 2)$, no choice of exponents gives $K([2] \times [2]) \cong \bigoplus_S (\mathbb{Z}/\gcd S)^{e(S)}$. There is not even an injective homomorphism from $K([2] \times [2])$ into such a group. Hence the Claim, and with it Conjecture 00000001230, is false.*

Proof. The subsets of $\{m_1, m_2\} = \{2, 2\}$ (as index subsets) are $\emptyset, \{m_1\}, \{m_2\}$ and $\{m_1, m_2\}$, with gcds 0, 2, 2, 2. So every modulus is 0 or 2. Apply Lemma 4 to $G = K(C_4)$ with the element of Proposition 5. $\square$

5 Robustness

Edge-count convention. If $[m]$ is the path with m edges, the 4-cycle is $[1] \times [1]$. The moduli are then $\gcd(\emptyset) = 0$ and $\gcd\{1\} = \gcd\{1, 1\} = 1$. All of them lie in $\{0, 1, 2\}$, so Lemma 4 applies again. (A product of copies of $\mathbb{Z}$ and 0 is torsion-free.)

Empty set and value-set reading. With $\gcd(\emptyset) = 1$, the moduli are $\{1, 2\}$. If the empty set is skipped, or S ranges over subsets of the value set $\{2\}$, the available moduli form a subset of $\{0, 2\}$. Every case is covered by Lemma 4, which allows an arbitrary family of moduli from $\{0, 1, 2\}$, including the choice $e(S) = 0$.

Infinite exponents. Lemma 4 allows any index set I . In Lean the index type ι is arbitrary.

Other meanings of “Jacobian”. In $\text{Pic}(C_4) = \mathbb{Z}^V / \Delta \mathbb{Z}^V$, the degree-zero class $p = [(0, 1) - (0, 0)]$ satisfies $\Delta(0, 3, 1, 2)^\top = (-4, 4, 0, 0)^\top = 4((0, 1) - (0, 0))$, so $4p = 0$. Moreover $2p \neq 0$: if $\Delta z = 2((0, 1) - (0, 0))$, then subtracting the constant $z_{(0,0)}$ from z (the all-ones vector lies in $\ker \Delta$) and reading the last three rows gives $\tilde{\Delta} z' = 2e_0$, which is impossible. So Lemma 4 applies both to $\text{Pic}(C_4)$ and to its subgroup $\text{Pic}^0(C_4)$, which contains p . Lean formalizes both.

Dimension three. If the claim is meant only for $n \geq 3$, take the cube $[2] \times [2] \times [2]$. Its moduli are again 0 and 2 (respectively 0 and 1 for the edge convention $[1]^3$). Order the non-sink vertices $(0, 0, 1), (0, 1, 0), (0, 1, 1), (1, 0, 0), (1, 0, 1), (1, 1, 0), (1, 1, 1)$, with sink $(0, 0, 0)$. Then $\tilde{\Delta}(3, 3, 6, 2, 3, 3, 4)^\top = 8e_2$, where e_2 is the chip at $(0, 1, 1)$. The unique rational solution of $\tilde{\Delta} z = 4e_2$ is not integral ($\det \tilde{\Delta} = 384$; the solution is $\frac{1}{2}(3, 3, 6, 2, 3, 3, 4)$). So $g = [2e_2]$ has $4g = 0 \neq 2g$, and Lemma 4 applies. In fact $K([2]^3) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/8 \oplus \mathbb{Z}/24$.

The failure is generic. By Smith normal form (`verify.py`),

grid	K	moduli (vertex / edge convention)
2×2	$\mathbb{Z}/4$	$\{0, 2\} / \{0, 1\}$
2×3	$\mathbb{Z}/15$	$\{0, 1, 2, 3\} / \{0, 1, 2\}$
3×3	$\mathbb{Z}/8 \oplus \mathbb{Z}/24$	$\{0, 3\} / \{0, 2\}$
2×4	$\mathbb{Z}/56$	$\{0, 2, 4\} / \{0, 1, 3\}$
3×4	$\mathbb{Z}/2415$	$\{0, 1, 3, 4\} / \{0, 1, 2, 3\}$
$2 \times 2 \times 2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/8 \oplus \mathbb{Z}/24$	$\{0, 2\} / \{0, 1\}$

A finite group $\bigoplus_a (\mathbb{Z}/d)^{e_d}$ has no $\mathbb{Z}$ summand. Its exponent divides the lcm of the nonzero moduli. In every row, the largest invariant factor of K does not divide that lcm. The order of K is the number of spanning trees, and it is typically divisible by primes that do not divide any m_i , such as 5 in $\mathbb{Z}/15$ or 7 in $\mathbb{Z}/2415$. Such primes can never come from the moduli.

Remark 7. The conjecture mentions “torus gluing”. The object of the conjecture is the grid graph, not the torus. Still, `verify.py` also checks that the discrete tori fail: $K(C_3 \square C_3) = (\mathbb{Z}/6)^2 \oplus (\mathbb{Z}/18)^2$ against moduli $\{0, 3\}$, and $K(C_3 \square C_4) = (\mathbb{Z}/5)^2 \oplus \mathbb{Z}/35 \oplus \mathbb{Z}/420$ against moduli $\{0, 1, 3, 4\}$. This remark is not formalized.

6 Lean formalization

The directory `lean4/` is a self-contained Lean 4.19.0 project (core library only, no Mathlib).

- `gridVerts ms` lists the vertices of the grid for an *arbitrary* list $\mathbf{ms} = [m_1, \dots, m_n]$. `adj` is adjacency at ℓ^1 distance 1, `deg` the degree, `lap` the Laplacian, and `redLap` the reduced Laplacian with sink $(0, \dots, 0)$. All of these are general definitions. The concrete matrices are obtained by evaluating them (`redLap_22`, `lap_22` and `redLap_222`, by `decide`).
- `Coker k M` is the quotient type $\mathbb{Z}^k / M\mathbb{Z}^k$, where $x \sim y$ iff $x - y = Mz$ for some $z \in \mathbb{Z}^k$. Lean proves that this is an equivalence relation compatible with addition. `Sandpile ms` is `Coker (|V| - 1) (redLap ms)`, and `Pic ms` is `Coker |V| (lap ms)`. `Pic0` is the subtype of $\text{Pic}([2] \times [2])$ of degree-zero classes; the degree is well defined because the columns of Δ sum to 0.
- `ZM d` is $\mathbb{Z}/d$, represented by the canonical representatives a with $a \bmod d = a$ (so `ZM 0` = $\mathbb{Z}$). `subsetsL` lists the 2^n sub-lists, and `gcdL` is the gcd of a list with `gcd(∅) = 0`. `Target ms e`

is the product over pairs (subset k , copy $c < e(k)$) of $\mathbb{Z}/\gcd(S_k)$, that is, $\bigoplus_S (\mathbb{Z}/\gcd S)^{e(S)}$. `IsIso` means additive and bijective.

- `ClaimV` (vertex convention), `ClaimE` (edge convention) and `ClaimV3` (vertex convention, $n \geq 3$) state the Claim. The main theorems are:

```
conjecture_00000001230_false : ¬ ClaimV
conjecture_00000001230_false_edges : ¬ ClaimE
conjecture_00000001230_false_dim3 : ¬ ClaimV3
```

- The explicit instances are `no_iso_22`, `no_iso_111_edges`, `no_iso_22_Pic0` and `no_iso_22_Pic`. The strong forms `no_injective_hom_C4`, `_Q3`, `_PicC4` and `_Pic0C4` exclude injective additive maps into $\prod_{i \in \iota} \mathbb{Z}/d_i$ for an arbitrary type ι and arbitrary $d_i \in \{0, 1, 2\}$.
- Witnesses: `g22_four`, `g22_two`, `p22_four`, `p22_two`, `g222_four` and `g222_two`. Each $4g = 0$ is proved with the explicit vector z . Each $2g \neq 0$ is proved by `omega` on the linear system.
- Non-vacuity and sanity: `sandpile_C4_iso_Z4` constructs the isomorphism $K([2] \times [2]) \cong \mathbb{Z}/4$ of Proposition 5. This shows that `Sandpile` is the right group and that `IsIso` can be satisfied. `sandpile_P3_trivial` shows that the Claim’s predicate holds for the path `[3]` (a tree; $e \equiv 0$). So the Claim is not refuted for a trivial reason.

`lake build` succeeds with no errors or warnings. There is no `sorry`, no `native_decide`, and no added axioms. `#print axioms` reports only `propext` and `Quot.sound`.

Limitations. The second clause (the 2-dimensional formula via $K_{m,n}$ and torus gluing) is not formalized, because it is not stated precisely. It is not needed, since the first clause already fails. The value-set reading of “subsets” and the convention $\gcd(\emptyset) = 1$ are not separate Lean predicates. They are covered by the strong lemmas, which allow every family of moduli in $\{0, 1, 2\}$. The isomorphisms $K \cong \text{Pic}^0$ and $\text{Pic} \cong \mathbb{Z} \oplus K$ are not needed; Lean treats the three groups separately. The torus remark and the table above are checked only in `verify.py`.

7 Independent check

`verify.py` (Python 3 standard library) builds the grids from coordinates. It computes Smith normal forms of the reduced and full Laplacians with exact integer arithmetic and recomputes the witnesses. For each listed grid and both conventions, it checks that the largest invariant factor of K does not divide the lcm of the nonzero moduli. As sanity checks, it verifies that paths (trees) have trivial Jacobian, so the predicate holds for them, and the torus remark.

8 Reproduction

```
cd lean4 && lake build      # Lean 4.19.0, core only
python3 verify.py          # Python 3 standard library
pdflatex report.tex
```

References

- [1] M. Baker and S. Norine, Riemann–Roch and Abel–Jacobi theory on a finite graph, *Adv. Math.* 215 (2007), 766–788.
- [2] C. J. Klivans, *The Mathematics of Chip-Firing*, CRC Press, 2018.
- [3] D. Lorenzini, Smith normal form and Laplacians, *J. Combin. Theory Ser. B* 98 (2008), 1271–1300.