

Disproof of Conjecture 00000007672: $\gcd(F_6, F_3) = 1$ is not a multiple of $F_{\gcd(6,3)} = 1 + q$

jilint777

2026-10-04

Abstract

Conjecture 00000007672 asserts that for the q -Fibonacci polynomials $F_n(q) = \sum_j \begin{bmatrix} n-1-j \\ j \end{bmatrix}_q q^{j^2}$, the greatest common divisor $\gcd(F_n, F_m)$ in $\mathbb{Z}[q]$ equals $F_{\gcd(n,m)}$ times a correction factor. We show that $\gcd(F_6, F_3) = 1$ in $\mathbb{Z}[q]$, while $F_{\gcd(6,3)} = F_3 = 1 + q$. Since 1 is not $(1 + q)$ times any polynomial, the claim fails. More generally the claim fails for $(3, 3k)$ for every $k \geq 2$. The main counterexample is fully formalized in Lean 4 (core library only), including the proof that 1 is a gcd of F_6 and F_3 in $\mathbb{Z}[q]$.

1 The conjecture

The statement (English version) reads:

Definition: The q -Fibonacci polynomials are $F_n(q) = \sum [n-1-j \text{ choose } j]_q q^{j^2}$, with Gaussian binomial coefficients. *Conjecture:* $\gcd(F_n(q), F_m(q))$ in $\mathbb{Z}[q]$ equals $F_{\gcd(n,m)}(q)$ times a correction factor from a finite exceptional set E depending only on primes $p \equiv \pm 2 \pmod{5}$, and the count of $E \cap [1, x]$ is $O(\log \log x)$.

The Chinese version says the same thing. We refute the first clause:

$$\gcd(F_n, F_m) = F_{\gcd(n,m)} \cdot c_{n,m} \quad \text{for some polynomial } c_{n,m}. \quad (1)$$

The “correction factor” multiplies a polynomial and is supposed to yield the gcd, which lies in $\mathbb{Z}[q]$. So it is a polynomial, in $\mathbb{Z}[q]$ or at least in $\mathbb{Q}[q]$. Our argument works for $c_{n,m} \in \mathbb{Q}[q]$ too (Remark 4). If $c_{n,m}$ were allowed to be an arbitrary rational function, (1) would hold for any polynomial in place of $\gcd(F_n, F_m)$ and would say nothing.

2 Definitions and small values

The Gaussian binomial coefficient is $\begin{bmatrix} n \\ k \end{bmatrix}_q = \prod_{i=1}^k \frac{1-q^{n-k+i}}{1-q^i}$ for $0 \leq k \leq n$, and 0 otherwise. It satisfies the q -Pascal rule $\begin{bmatrix} n+1 \\ k+1 \end{bmatrix}_q = \begin{bmatrix} n \\ k \end{bmatrix}_q + q^{k+1} \begin{bmatrix} n \\ k+1 \end{bmatrix}_q$. Only the terms with $0 \leq j \leq n-1-j$ contribute to F_n . We get

$$\begin{aligned} F_3 &= \begin{bmatrix} 2 \\ 0 \end{bmatrix}_q + \begin{bmatrix} 1 \\ 1 \end{bmatrix}_q q = 1 + q, \\ F_6 &= \begin{bmatrix} 5 \\ 0 \end{bmatrix}_q + \begin{bmatrix} 4 \\ 1 \end{bmatrix}_q q + \begin{bmatrix} 3 \\ 2 \end{bmatrix}_q q^4 = 1 + (q + q^2 + q^3 + q^4) + (q^4 + q^5 + q^6) \\ &= 1 + q + q^2 + q^3 + 2q^4 + q^5 + q^6. \end{aligned}$$

As a sanity check, $F_n(1) = \sum_j \binom{n-1-j}{j}$ is the Fibonacci number $1, 1, 2, 3, 5, 8, \dots$

Lemma 1. $F_3(-1) = 0$ and $F_6(-1) = 2$. Moreover $F_6 = F_3 \cdot (-1 + 2q - q^2 + 2q^3 + q^5) + 2$.

Proof. Direct computation. $F_6(-1) = 1 - 1 + 1 - 1 + 2 - 1 + 1 = 2$. □

3 The gcd of F_6 and F_3 in $\mathbb{Z}[q]$

Proposition 2. 1 is a greatest common divisor of F_6 and F_3 in $\mathbb{Z}[q]$. That is, every $D \in \mathbb{Z}[q]$ dividing both F_6 and F_3 divides 1 , i.e. $D = \pm 1$.

Proof. Write $DA = F_3 = 1 + q$ and $DB = F_6$ with $A, B \in \mathbb{Z}[q]$. Degrees add in $\mathbb{Z}[q]$, because the top coefficient of a product is the product of the top coefficients and $\mathbb{Z}$ has no zero divisors. Hence $\deg D + \deg A = 1$, so D or A is constant.

If $A = a$ is constant, then $D(0)a = 1$ gives $a \neq 0$, and $D(-1)a = F_3(-1) = 0$ gives $D(-1) = 0$. But then $F_6(-1) = D(-1)B(-1) = 0$, contradicting $F_6(-1) = 2$ (Lemma 1).

If $D = d$ is constant, then $d \cdot A(0) = 1$ forces $d = \pm 1$, so $D \cdot d = 1$ and $D \mid 1$. □

Theorem 3. For $(n, m) = (6, 3)$ the clause (1) fails. Conjecture 00000007672 is false.

Proof. $\gcd(6, 3) = 3$ and $F_3 = 1 + q$. By Proposition 2 the gcd of F_6 and F_3 in $\mathbb{Z}[q]$ is ± 1 , which is unique up to units. If $\pm 1 = (1 + q)c$ for a polynomial c , then evaluating at $q = -1$ gives $\pm 1 = 0$, a contradiction. More generally, no divisor G of F_6 is a multiple of F_3 . If $G = F_3c$ and $GH = F_6$, then $2 = F_6(-1) = F_3(-1)c(-1)H(-1) = 0$. □

Remark 4. The same evaluation argument works if the correction factor c is allowed to lie in $\mathbb{Q}[q]$, or even to be any rational function without a pole at $q = -1$. Over $\mathbb{Q}[q]$ the gcd is also 1, because $F_6 \equiv 2 \pmod{1 + q}$.

Remark 5 (Infinitely many failures). The polynomials F_n satisfy Schur's recurrence $F_n = F_{n-1} + q^{n-2}F_{n-2}$ (checked for $n \leq 30$ in `verify.py`). Put $f_n = F_n(-1)$. Then $f_{2k} = f_{2k-1} + f_{2k-2}$ and $f_{2k+1} = f_{2k} - f_{2k-1}$, so $f_{2k+1} = f_{2k-2}$ and $f_{2k} = f_{2k-2} + f_{2k-4}$. The even-indexed values are therefore Fibonacci numbers, and $f_n > 0$ for every $n \neq 3$. Hence $1 + q = F_3$ divides no F_{3k} with $k \geq 2$, and the argument of Theorem 3 refutes (1) for every pair $(3, 3k)$, $k \geq 2$. The Lean project checks $f_{3k} \neq 0$ for $k = 2, \dots, 6$. So the failure cannot be confined to a finite exceptional set of pairs, under any reading of the set E .

4 Lean formalization

The directory `lean4/` is a self-contained Lean 4.19.0 project that uses only the core library (no Mathlib).

- Polynomials in $\mathbb{Z}[q]$ are coefficient lists. `PEq` means equal coefficients, `Dvd a b` is $\exists c, a \cdot c = b$ in $\mathbb{Z}[q]$, and `IsGcd G a b` is the standard definition of a greatest common divisor (a common divisor that every common divisor divides).
- `gauss` defines Gaussian binomials by the q -Pascal rule. `gauss_product_formula` checks $\begin{bmatrix} n \\ k \end{bmatrix}_q (q; q)_k (q; q)_{n-k} = (q; q)_n$ for all $k \leq n \leq 7$. `qfib n` is $\sum_{j < n} \begin{bmatrix} n-1-j \\ j \end{bmatrix}_q q^{j^2}$ exactly as in the conjecture. `qfib_at_one` checks $F_n(1) = \text{Fib}_n$ for $n \leq 15$.

- `common_divisor_dvd_one` and `isGcd_one` formalize Proposition 2. They use the degree argument (`coeff_mull_top`, `const_or_const`) and evaluation (`evalL_mull`, `evalL_congr`).
- `GcdClause` is the clause (1): for all $n, m \geq 1$ and every gcd G of F_n, F_m in $\mathbb{Z}[q]$, $G = F_{\gcd(n,m)} \cdot c$ for some $c \in \mathbb{Z}[q]$. `conjecture_00000007672.false` proves $\neg \text{GcdClause}$. `no_divisor_of_F6_is_multiple_of_F3` proves the stronger statement from the proof of Theorem 3.

`lake build` succeeds. There is no `sorry`, no `native_decide`, and no added axioms. `#print axioms` shows at most `propext`, `Classical.choice` and `Quot.sound`. The numerical facts are proved by `decide`; `qfib_mult3_at_neg_one` uses `decide +kernel`, which is checked by the Lean kernel.

5 Reproduction

```
cd lean4 && lake build      # Lean 4.19.0, core only
python3 verify.py          # standard library only
pdflatex report.tex
```

`verify.py` computes Gaussian binomials from the product formula by exact division, which is independent of the q -Pascal rule used in Lean. It recomputes F_n , finds $\gcd(F_6, F_3) = 1$ by the Euclidean algorithm in $\mathbb{Q}[q]$ (and in $\mathbb{Z}[q]$ by Gauss’s lemma, since F_3 is primitive), and lists all 76 pairs $n < m \leq 30$ for which $F_{\gcd(n,m)} \nmid F_n$ or $F_{\gcd(n,m)} \nmid F_m$.

References

- [1] L. Carlitz, Fibonacci notes 3: q -Fibonacci numbers, *Fibonacci Quart.* 12 (1974), 317–322.
- [2] G. E. Andrews, Fibonacci numbers and the Rogers–Ramanujan identities, *Fibonacci Quart.* 42 (2004), 3–19.