

Infinitely many highly transitive complete designs

Counterexample to TLMC 00000008411

ziangni-sys

4 October 2026

Abstract

For every $v \geq 5$, the complements of single points in a v -point set form a 3 -($v, v-1, v-3$) design. Every point permutation is an automorphism, and the automorphism group is 3-transitive. Different values of v give nonisomorphic designs. This supplies infinitely many isomorphism classes with the fixed strength $t = 3$.

1 Scope of the assertion

The source defines a t -(v, k, λ) design by the condition that each t -element subset is contained in exactly λ blocks. It then asserts that for fixed $t \geq 3$ only finitely many isomorphism classes have a t -transitive automorphism group.

The following family consists of *complete designs*, often called trivial designs in classification terminology. The stated definition does not exclude complete designs. Our parameters nevertheless satisfy $3 < k < v$. We make no claim about a different classification problem that adds a noncomplete-design hypothesis. Only t is fixed in the source; the parameters v, k, λ are allowed to vary.

2 An explicit infinite family

For each $v \geq 5$, let X_v be a set of v points and put

$$\mathcal{B}_v = \{X_v \setminus \{x\} : x \in X_v\}.$$

Distinct omitted points give distinct blocks, so this is a simple block system with exactly v blocks, each of cardinality $v-1$. It is precisely the family of all $(v-1)$ -element subsets of X_v .

Proposition 1. *($X_v, \mathcal{B}_v$) is a 3 -($v, v-1, v-3$) design, and its automorphism group is 3-transitive.*

Proof. Fix any three-element subset $T \subseteq X_v$. The block $X_v \setminus \{x\}$ contains T exactly when $x \notin T$. There are $v-3$ such points, and distinct points give distinct blocks. This proves the design property with $\lambda = v-3 > 0$.

For every permutation σ of X_v ,

$$\sigma(X_v \setminus \{x\}) = X_v \setminus \{\sigma(x)\}.$$

Thus every permutation preserves the block family and is an automorphism. Given any two ordered triples of distinct points, send the first triple to the second in the prescribed order. This bijection between their three-element ranges extends to a permutation of X_v , because the two complements have equal cardinality $v-3$. The resulting automorphism sends the triples to one another, proving 3-transitivity. $\square$

Proposition 2. *These designs have infinitely many distinct isomorphism classes.*

Proof. An isomorphism of block designs includes a bijection of their point sets. Consequently the designs for v and w can be isomorphic only if $v = w$. Taking $v = n + 5$ for every natural number n therefore gives an infinite sequence of pairwise nonisomorphic designs. The value $t = 3$ is fixed throughout, so this contradicts the claimed finiteness. $\square$

The finiteness clause already fails, so the subsequent classification list cannot establish the conjecture as stated.

3 Formal correspondence

The Lean project defines the actual point set as `Fin (n+5)`. The block omitting i is `Finset.univ.erase i`, and the block family is its finite image over all points. The theorem `block_injective` rules out duplicate blocks, while `block_card` and `number_of_blocks` verify the sizes.

The definition `IsThreeDesign` requires every block to have $n + 4$ points and quantifies over every actual three-element subset, counting the blocks containing it. The theorem `incidence_count` identifies this filtered family with the omitted points outside that subset and computes its cardinality $n + 2$. Thus `actual_design` proves the full defining incidence condition, not only a numerical parameter identity.

`IsAutomorphism` is the actual block-preservation biconditional for a permutation of the point set. The theorem `permutation_preserves` proves it for every permutation. Ordered triples of distinct points are represented by embeddings `Fin 3` into the point set. The theorem `three_transitive` constructs a bijection of their ranges and uses the finite-set extension theorem to produce an actual permutation with the required action on all three entries.

`Isomorphic` requires an actual point bijection preserving block membership in both directions. The theorem `isomorphic_iff` proves that two family members are isomorphic exactly when their natural-number parameters are equal. The formal quotient by this isomorphism relation is proved infinite by an injective map from $\mathbb{N}$. Hence the final theorem `conjecture_00000008411` includes actual designs, actual 3-transitivity, the inequalities $3 < k < v$, and an infinite type of isomorphism classes.

4 Verification

Lean 4.19.0 is used with Mathlib commit `c44e0c8ee63ca166450922a373c7409c5d26b00b`. Run `lake build`, then `lake env lean Main.lean` in the `lean` directory. The axiom audits of the design theorem, transitivity theorem and final theorem contain only the standard logical axioms `propext`, `Classical.choice` and `Quot.sound`. There are no admitted proofs, extra axioms, native evaluation shortcuts, or auxiliary computational dependencies.