
a
Descriptor groups supporting each readout
(relative permutation drop, %)
Intra-nt backbone
Glycosidic
Helical/ torsion
Local neighborhood
Inter-nt edge
Global/ center
Unclassified
20.4
0.1
3.9
2.0
0.4
2.4
2.0
-0.1
2.1
1.4
0.3
10.0
30.6
23.9
12.2
2.2
2.4
31.5
7.9
0.1
0.4
45
40
35
30
25
20
15
10
5
0
Nucleotide
Base-pair
SASA
b
Solvent-accessibility signal is not carried only by explicit proxies
SASA Pearson r
0.96
0.95
0.94
0.93
0.92
0.91
0.90
0.948
0.948
0.942
0.942
all 935
no explicit SASA
no SASA + global
no SASA +global+density
c
Atom-channel controls
Weighted recovery (%)
64
62
60
58
56
54
61.9
58.5 (-3.4)
57.3 (-4.7)
57.4 (-4.5)
61.6 (-0.3)
61.0 (-0.9)
7-atom
N mask
N zero
N dummy
5-atom
3-atom
d
Family calibration (families with n ≥ 20)
Purine
AdoCbl-II
tRNA
5S rRNA
5.8S rRNA
GOLLD
RNaseP bact b
ROOL
15.0
13.5
12.3
8.5
7.5
6.9
5.6
5.4
0.0
2.5
5.0
7.5
10.0
12.5
15.0
17.5
Mean per-chain ECE (%)
e
Method and resolution strata
(adjusted resolution effect n.s.)
NMR/other
cryo-EM >4.0 A
cryo-EM 3.5-4.0 A
cryo-EM 3.0-3.5 A
cryo-EM <=3.0 A
X-ray >3.0 A
X-ray 2.5-3.0 A
X-ray 2.0-2.5 A
X-ray <=2.0 A
44
70
64
62
65
55
71
73
54
40
50
60
70
80
Weighted recovery (%)
f
Chain length (nt)
201-500
101-200
50-100
<50
51 (n=231)
80 (n=701)
68 (n=993)
54 (n=2210)
30
40
50
60
70
80
90
Weighted recovery (%)