Medicinal Chemistry 1030042 Midterm Exam

Class:

Dr. An-Rong Lee Mar. 15, 2024

Name:

1. Give reasons why amyl nitrite is suitable for emergency use in the case of angina pectoris while pentaerythritol tetranitrate for preventive purpose. (10%)

Amyl nitrite

Pentaerythritol tetranitrate


- 2. Briefly describe Digoxin Immune Fab as an antidote for digoxin overdose (10%)
- 3. (a) Amiodarone is one of the most effective antiarrhythmics with a very broad spectrum; however, it is a last choice in clinical uses. Why? (10%) (b) From the structure features of dronedarone, explain how it canimprove those drawbacks of amiodarone? (10%)

Amiodarone

Dronedarone

- 4. Inclisiran is newly approved for the treatment of atherosclerotic cardiovascular disease and heterozygous familial hypercholesterolemia. Briefly describe its mechanism of action (for a more readily comprehensible answer, you may also draw an illustrative figure). (10%)
- 5. Clopidogrel is used as a substitute for aspirin. Now from the structure features of prasugrel, give explanations why it is a better drug than clopidogrel in terms of efficacy of inhibiting platelet aggregation. (10%)

Clopidogrel



Prasugrel

112 學年第二學期大學部藥物化學(II)期中考試

113-3-15

姓名: 言节往往日子

學號: 410030013

一、 選擇題 (20%)

1. 下列利尿劑中,何者最適用於腎功能不佳之水腫患者?

2. 下列利尿劑中,何者常用於青光眼治療?

3. 下列何者為 spironolactone 的主要活性代謝產物?

4. 下列降血壓藥,何者最適合與 thiazide 利尿劑併用,協助平衡血鉀濃度?

5. 下列 ACE 抑制劑中,何者不屬於前驅藥?

A. Benazepril B. Enalapril C. Lisinopril D. Ramipril

6. 左式結構為一類降血壓藥物,其螯合於作用標的活性中心金

屬離子的基團為? A. 酸基 a B. 酸基 b C. 酮基 c D. 胺基 d

7. Angiotensin II receptor blockers (ARBs)的簡式(如下圖),何者並非

ARBs 中R的結構?

8. 下列何者是最適用於治療良性前列腺肥大?

9. a-Methyldopa 可經由何種反應而產生活性代謝物?

A. 先經 Decarboxylation 再經 Methylation B. 先經 Hydroxylation 再經 Decarboxylation C. 先經 Decarboxylation 再經 Hydroxylation D. 先經 Hydroxylation 再經 Methylation

10. Sodium nitroprusside (Na₂Fe(CN)₅NO·2H₂O 結構中的 Fe 與 NO 基團的價數分別為? A. + II 與 + I B. + III 與 -I C. + II 與 0 D. + III 與 0

二、 問答題 (20%)

1. 請以反應方程式敘述 omeprazole (結構如下) 制酸之作用機制 (8%)。

2. 請敘述 4 種主要利尿劑及其作用機制,並各舉一代表藥物 (12%)。