

Conditional Probability

Last Updated : 23 May, 2025


Conditional probability defines the probability of an event occurring based on a given condition or prior knowledge of another event. Conditional probability is **the likelihood of an event occurring**, given that another event has already occurred. In probability this is denoted as A given B, expressed as $P(A | B)$, indicating the probability of event A when the event B has already occurred.

Example :

We flip two coins. What is the probability that both are heads, given that at least one of them is heads?

We are given that there is at least one head. So there are three possibilities HH, HT, TH. Only one outcome has both coins as heads: HH.

So our answer is $= 1/3$



We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Got It !

Let's consider two events A and B, then the **formula for the conditional probability** of B when A has already occurred is given by:

$$P(B|A) = \frac{P(B \cap A)}{P(A)}$$

Where,

P (A ∩ B) represents the probability of both events A and B occurring simultaneously.

P(A) represents the probability of event A occurring.

Steps to Find Probability of One Event Given Another Has Already Occurred

To **calculate** the conditional probability, we can use the following step-by-step method:

Step 1: *Identify the Events. Let's call them Event A and Event B.*

Step 2: *Determine the Probability of Event A i.e., **P(A)***

Step 3: *Determine the Probability of Event B i.e., **P(B)***

Step 4: *Determine the Probability of Event A and B i.e., **P(A ∩ B)**.*

Step 5: *Apply the Conditional Probability Formula and calculate the required probability.*

Conditional Probability Examples

There are various examples of conditional probability as in real life all the events are related to each other and happening any event affects the probability of another event. For **example**, if it rains, the probability of road accidents increases as roads have less friction. Let's consider some problem-based examples here:

1) Tossing a Coin

Let's consider two events in tossing two coins,

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Sample space for tossing two coins is:

$$S = \{HH, HT, TH, TT\}$$

The conditional probability of getting a head on the second coin (B) given that we got a head on the first coin (A) is $= P(B|A)$

Since the coins are independent (one coin's outcome does not affect the other),

$P(B|A) = P(B) = 0.5$ (50%), which is the probability of getting a head on a single coin toss.

2) Drawing Cards

In a deck of 52 cards where two cards are being drawn, then let's consider the events be.

- **A:** Drawing a red card on the first draw, and
- **B:** Drawing a red card on the second draw.

The conditional probability of drawing a red card on the second draw (B) given that we drew a red card on the first draw (A) is $= P(B|A)$

After drawing a red card on the first draw, there are 25 red cards and 51

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Properties of Conditional Probability

Some of the common properties of conditional property are:

1: Let's consider an event A in any sample space S of an experiment.

$$P(S | A) = P(A | A) = 1$$

2: For any two events A and B of a sample space S, and an event X such that $P(X) \neq 0$,

$$P((A \cup B) | X) = P(A | X) + P(B | X) - P((A \cap B) | X)$$

3: The order of sets or events is important in conditional probability, i.e.,

$$P(A | B) \neq P(B | A)$$

4: The complement formula for probability only holds conditional probability if it is given in the context of the first argument in conditional probability i.e.,

$$P(A' | B) = 1 - P(A | B)$$

$$P(A | B') \neq 1 - P(A | B)$$

5: For any two or three independent events, the intersection of events can be calculated using the following formula:

- For the intersection of two events A and B

$$P(A \cap B) = P(A) P(B)$$

- For the intersection of three events A, B, and C,

$$P(A \cap B \cap C) = P(A) P(B) P(C)$$

Conditional Probability and Independent Events

With the help of conditional probability, we can tell apart dependent and independent events. When the probability of one event happening doesn't influence the probability of any other event, then events are called independent, otherwise dependent events.

Conditional Probability of Independent Events

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

as $P(A)$ as there is no effect of event B on the probability of event A. For independent events, A and B, the conditional probability of A and B concerning each other is given as follows:

- $P(B | A) = P(B)$
- $P(A | B) = P(A)$

Check, [Probability Formulas](#)

Conditional Probability vs Joint Probability vs Marginal Probability

The difference between Conditional Probability, **Joint Probability**, and **Marginal Probability** is given in the following table:

Parameter	Conditional Probability	Joint Probability	Marginal Probability
Definition	The probability of an event occurring is given. That another event has already occurred.	The probability of two or more events occurring simultaneously.	The probability of an event occurring without considering any other events.
Calculation	$P(A B)$	$P(A \cap B)$	$P(A)$
Variables involved	Two or more events	Two or more events	Single event.

Conditional Probability and Bayes' Theorem

Bayes' Theorem is a key concept in probability theory that updates beliefs

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

$$P(A|B) = (P(B|A) \times P(A)) / P(B)$$

Where:

- $P(A|B)$ is the updated probability of hypothesis A given evidence B (posterior probability).
- $P(B|A)$ is the likelihood of evidence B given hypothesis A.
- $P(A)$ is the initial belief in hypothesis A (prior probability).
- $P(B)$ is the total probability of observing evidence B.

Bayes' Theorem allows for iterative updates as new data is gathered, making it useful in fields like machine learning, medical diagnosis, and spam filtering. It provides a systematic approach to reasoning under uncertainty.

Here's a breakdown of how Bayes' Theorem works:

- **Prior Probability $P(A)$:** Initial belief in hypothesis A before evidence.
- **Likelihood $P(B|A)$:** Probability of observing evidence B if A is true.
- **Evidence $P(B)$:** Normalization factor, total probability of B across all hypotheses.
- **Posterior Probability $P(A|B)$:** Updated probability of A after considering evidence B.

Multiplication Rule of Probability

Multiplication Rule of Probability, when applied in the context of conditional probability, helps us calculate the probability of the intersection of two events when the probability of one event depends on the occurrence of the other event. **This rule is crucial in understanding the joint probability of events under specific conditions.**

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

$$P(A \cap B) = P(A) \times P(B | A)$$

Here's what each term represents:

- **$P(A \cap B)$** : This denotes the probability that both events A and B occur simultaneously.
- **$P(A)$** : This represents the probability of event A happening.
- **$P(B|A)$** : This is the conditional probability of event B occurring given that event A has already occurred.

How to Apply the Multiplication Rule?

To apply the Multiplication Rule in the context of conditional probability, we can use the following steps:

- First, we calculate the probability of event A occurring.
- Then, we compute the probability of event B occurring given that event A has occurred.
- Multiplying these probabilities together gives us the joint probability of both events happening under the specified conditions.
- This rule is particularly useful when dealing with events that are not independent, meaning that the occurrence of one event affects the probability of the other event.

Applications of Conditional Probability

Various applications of conditional probability are,

Finance and Risk Management

- **Example:** Assessing the probability of default for a borrower given certain financial indicators.
- **Application:** Banks and financial institutions use conditional probability to evaluate the risk associated with loans and investments.

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- **Example:** Determining the probability of a patient having a specific disease given the results of diagnostic tests.
- **Application:** Conditional probability is crucial in medical diagnoses and decision-making, helping healthcare professionals make informed decisions based on test results.

Marketing and Customer Relationship Management (CRM)

- **Example:** Predicting the probability of a customer making a purchase based on their past buying behavior.
- **Application:** Businesses use conditional probability to tailor marketing strategies, optimize customer experiences, and personalize product recommendations.

Machine Learning and Artificial Intelligence

- **Example:** Predicting the likelihood of a user clicking on a particular ad based on their online behavior.
- **Application:** Conditional probability is fundamental in machine learning algorithms for tasks such as classification, recommendation systems, and natural language processing.

Weather Forecasting

- **Example:** Estimating the probability of rain tomorrow given today's weather conditions.
- **Application:** Meteorologists use conditional probability to make weather predictions based on historical data and current atmospheric conditions.

Solved Examples of Conditional Probability

Question 1: A bag contains 5 red balls and 7 blue balls. Two balls are drawn without replacement. What is the probability that the second

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Let the events be,

Event A: The first ball drawn is red.

Event B: The second ball drawn is red.

$P(A) = 5/12$ and $P(B) = 4/11$ (as first ball drawn is already red, thus only 4 red balls remain in the bag)

Therefore, probability of the second ball drawn being red given that the first ball drawn was red is $4/11$.

Question 2: In a survey among a group of students, 70% play football, 60% play basketball, and 40% play both sports. If a student is chosen at random and it is known that the student plays basketball, what is the probability that the student also plays football?

Solution:

Let's assume there are 100 students in the survey.

Number of students who play football = $n(A) = 70$

Number of students who play basketball = $n(B) = 60$

Number of students who play both sports = $n(A \cap B) = 40$

To find the probability that a student plays football given that they play basketball, we use the **conditional probability formula**:

$$P(A|B) = n(A \cap B) / n(B)$$

Substituting the values, we get:

$$P(A|B) = 40 / 60 = 2/3$$

Therefore, probability that a randomly chosen student who plays basketball also plays football is $2/3$.

Question 3: In a certain city, it rains 30% of the day. A weather forecaster correctly predicts rain 80% of the time when it rains, and correctly predicts no rain 90% of the time when it doesn't rain. If the forecast predicts rain, what is the probability that it will rain?

Solution:

Let R be the event that it rains, and F be the event that rain is

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

0.31

$$P(R|F) = (0.8 \times 0.3) / 0.31 = 0.7742$$

Question 4: A fair die is rolled twice. Given that the sum of the two rolls is even, what is the probability that the first roll was an even number?

Solution:

Total favorable outcomes for an even sum:

Both rolls even: 9 outcomes (e.g., (2, 2), (2, 4), ...)

Both rolls odd: 9 outcomes (e.g., (1, 1), (1, 3), ...)

Total = 18 outcomes.

Favorable outcomes where the first roll is even:

First roll even (2, 4, 6), second roll even (2, 4, 6): 9 outcomes.

Conditional probability: $P(\text{First roll even} \mid \text{Sum even}) = 9 / 18 = 1 / 2$

Question 5: In a bag, there are 4 red balls and 6 blue balls. Two balls are drawn without replacement. What is the probability that the second ball is red, given that the first ball drawn was blue?

Solution:

After drawing a blue ball, there are 4 red balls and 5 blue balls left.

$P(\text{Second is red} \mid \text{First was blue}) = 4 / (4 + 5) = 4/9 = 0.444$

Articles Related to Conditional Probability:

[Probability Theory](#)

[Addition Rule for Probability](#)

[Complement of a Set](#)

[Types of Events in Probability](#)

[Coin Toss Probability Formula](#)

[Mutually Exclusive Events](#)

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- [Probability Class 12 Notes](#)
- [Probability Class 12 NCERT Solutions](#)

[Comment](#)[More info](#)[Advertise with us](#)[Next Article](#)[Bayes' Theorem](#)

Similar Reads

1. Conditional Probability vs Bayes Theorem
2. Conditional Probability Practice Question
3. Bayes's Theorem for Conditional Probability
4. Conditional PDF
5. Conditional Convergence
6. Conditional Probability and Independence - Probability | Class 12 Maths
7. Conditional Statements & Implications - Mathematical Reasoning | Class 11 Maths
8. Non-Mutually Exclusive Events
9. Dependent Events
10. Rules of Inference



Corporate & Communications Address:

A-143, 7th Floor, Sovereign Corporate
Tower, Sector- 136, Noida, Uttar Pradesh
(201305)

Registered Address:

K 061, Tower K, Gulshan Vivante
Apartment, Sector 137, Noida, Gautam

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).



Advertise with us

Company

About Us
Legal
Privacy Policy
Careers
In Media
Contact Us
Corporate Solution
Campus Training Program

Tutorials

Python
Java
C++
PHP
GoLang
SQL
R Language
Android

Data Science & ML

Data Science With Python
Machine Learning
ML Maths
Data Visualisation
Pandas
NumPy
NLP
Deep Learning

Python Tutorial

Python Examples
Django Tutorial
Python Projects
Python Tkinter

Explore

Job-A-Thon
Offline Classroom Program
DSA in JAVA/C++
Master System Design
Master CP
Videos

DSA

Data Structures
Algorithms
DSA for Beginners
Basic DSA Problems
DSA Roadmap
DSA Interview Questions
Competitive Programming

Web Technologies

HTML
CSS
JavaScript
TypeScript
ReactJS
NextJS
NodeJs
Bootstrap
Tailwind CSS

Computer Science

GATE CS Notes
Operating Systems
Computer Network
Database Management System

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

DevOps

- Git
- AWS
- Docker
- Kubernetes
- Azure
- GCP
- DevOps Roadmap

System Design

- High Level Design
- Low Level Design
- UML Diagrams
- Interview Guide
- Design Patterns
- OOAD
- System Design Bootcamp
- Interview Questions

School Subjects

- Mathematics
- Physics
- Chemistry
- Biology
- Social Science
- English Grammar

Databases

- SQL
- MYSQL
- PostgreSQL
- PL/SQL
- MongoDB

Preparation Corner

- Company-Wise Recruitment Process
- Aptitude Preparation
- Puzzles
- Company-Wise Preparation

More Tutorials

- Software Development
- Software Testing
- Product Management
- Project Management
- Linux
- Excel
- All Cheat Sheets

Courses

- IBM Certification Courses
- DSA and Placements
- Web Development
- Data Science
- Programming Languages
- DevOps & Cloud

Programming Languages

- C Programming with Data Structures
- C++ Programming Course
- Java Programming Course
- Python Full Course

Clouds/Devops

- DevOps Engineering
- AWS Solutions Architect Certification
- Salesforce Certified Administrator Course

GATE 2026

- GATE CS Rank Booster
- GATE DA Rank Booster
- GATE CS & IT Course - 2026
- GATE DA Course 2026
- GATE Rank Predictor

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).