

Probability Formulas

Last Updated : 22 May, 2025

Probability Formulas are essential mathematical tools used in calculating the probability. Below is the main formula for probability.

Probability Formula

E : Event

$$P(E) = \frac{\text{Number of favourable outcomes}}{\text{Total number of possible outcomes}}$$

$0 \leq P(E) \leq 1$

Probability Formula

Probability of an Event = (Count of favorable outcomes) / (Total number of possible outcomes for the event)

$$P(A) = n(E) / n(S)$$

$$0 \leq P(A) \leq 1$$

Here, $P(A)$ signifies the probability of an event A, where $n(E)$ is the count of favorable outcomes, and $n(S)$ is the total number of possible outcomes for the event.

When considering the **complementary event**, represented as $P(A')$, which denotes the non-occurrence of event A. Then the formula will be:

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Got It !

$P(A')$ is the opposite of event A , indicating that either event $P(A)$ occurs or its complement $P(A')$ occurs.

Therefore, now we can say; $P(A) + P(A') = 1$

Learn,

- [Basic Concepts of Probability](#)
- [Events in Probability](#)
- [Types of Events in Probability](#)

Terms Related to Probability Formula

Some of the most common terms related to probability formulas are:

- **Experiment:** An Experiment is an action or procedure conducted to generate a particular outcome.
- **Sample Space:** The [Sample Space](#) includes the complete potential outcomes that come from an experiment. For example, when flipping a coin, the sample space includes {head, tail}.
- **Favorable Outcome:** A favorable outcome is the result that aligns with the intended or expected conclusion. In the case of rolling two dice, examples of favorable outcomes resulting in a sum of 4 are (1,3), (2,2), and (3,1).
- **Trial:** A trial denotes the execution of a random experiment.
- **Random Experiment:** A [Random Experiment](#) is characterized by a well-defined set of possible outcomes. An example of a random experiment is tossing a coin, where the result could be either heads or tails. That means the result would be uncertain.
- **Event:** An [Event](#) denotes the total outcomes that come from a random

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- **Equally Likely Events:** Equally Likely Events have identical probabilities of occurrence. The outcome of one event does not impact the outcome of another.
- **Exhaustive Events:** An [Exhaustive Event](#) occurs when the set of all possible outcomes covers the complete sample space.
- **Mutually Exclusive Events:** [Mutually Exclusive Events](#) are those that cannot occur simultaneously. For example, when we toss the coin, the result will be either a head or a tail, but we cannot get both at the same time.

Probability of an Event

In [Probability theory](#), an event represents a set of possible outcomes derived from an experiment. It often forms a subset of the overall sample space. If we represent the probability of an event E as $P(E)$, the following principles apply:

The probability $P(E)$ lies between 0 and 1.

- For an impossible event (E), $P(E) = 0$.
- For a certain event (E), $P(E) = 1$.

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

The sum of the probabilities of all possible outcomes in a random experiment is equal to 1.


Example : In a rolling die experiment

Possible Outcomes : $\{ 1, 2, 3, 4, 5, 6 \}$
then , $P(1) + P(2) + P(3) + P(4) + P(5) + P(6) = 1$

The different Probability Formulas are discussed below:

For a particular event E, probability formula will be $P(E) = n(E) / n(S)$

Here, $n(E)$ represents the number of outcomes favorable to event E, and $n(S)$ denotes the total count of outcomes within the sample space.



Probability Formulas

Complement Rule :

$$P(A) = 1 - P(A')$$

Basic Probability Formula :

$$P(E) = \frac{\text{Number of favourable outcomes}}{\text{total number of outcomes}}$$

< ▶ > 1 / 4

Classical Probability Formula

$P(A) = \text{Number of Favorable Outcomes} / \text{Total Number of Possible Outcomes}$

When we deal with an event that is the union of two separate events, for example, A and B, the probability of the union will be:

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Joint Probability Formula

It represents the common elements that constitute the distinct subsets of both events A and B. The formula can be expressed as:

$$P(A \cap B) = P(A|B)P(B) = P(B|A)P(A)$$

- $P(A \cap B)$ is the **joint probability**, meaning the probability that **both** events A and B occur.
- $P(A|B)$ is the **conditional probability** of A given that B has already occurred.
- $P(B|A)$ is the **conditional probability** of B given that A has already occurred.
- $P(A)$ and $P(B)$ are the probabilities of events A and B occurring individually.

Addition Rule for Mutually Exclusive Events

If events A and B are mutually exclusive, that means they cannot happen at the same time; the probability of either event occurring is equal to the sum of their respective probabilities. Then:

$$P(A \cap B) = 0$$

Thus, the **Addition Rule** for mutually exclusive events becomes:

$$P(A \cup B) = P(A) + P(B)$$

Complementary Rule Formula

If A is an event, then the probability of not A is expressed by the [complementary rule](#):

$$P(\text{not } A) = 1 - P(A) \text{ or } P(A') = 1 - P(A)$$

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- $P(A)$ is the probability that event A occurs.
- $P(A')$ is the probability that event A **does not** occur.
- Since an event either happens or it doesn't, their probabilities must add up to 1.

Some probability formulas based on complementary rules are as follows:

- $P(A \cap A') = 0$
- $P(A' \cap B) = P(B) - P(A \cap B)$
- $P(A \cap B') = P(A) - P(A \cap B)$
- $P(A \cup B) = P(A \cap B') + P(A' \cap B) + P(A \cap B)$
- $P(A \cap B) + P(A' \cap B') = 1$ (Not always true)

Conditional Rule Formula

In the case, where the occurrence of event A is already known, the probability of event B is going to occur, referred to as [conditional probability](#). It can be calculated using the formula:

$$P(B|A) = P(A \cap B) / P(A)$$

$P(B|A)$: Probability of event B when event A has already occurred.

Relative Frequency Formula

The [relative frequency](#) formula is based on frequencies observed in real-world data. This formula is given as

$$P(A) = \text{Number of Times Event A Occurs} / \text{Total Number of Trials or Observations}$$

Probability Formula with the Multiplication Rule

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- $P(A \cap B) = P(A) \cdot P(B)$ (in case of independent events)
- $P(A \cap B) = P(A) \cdot P(B|A)$ (in case of dependent events)

Disjoint Event

Two events **A** and **B** are **disjoint** (or **mutually exclusive**) if **they cannot happen at the same time**. This means their intersection is empty:

$$P(A \cap B) = 0$$

Bayes' Theorem

Bayes' Theorem calculates the probability of event A given the occurrence of event B. The [Bayes Theorem](#) Formula is given as

$$P(A|B) = P(B|A) \times P(A) / P(B)$$

- $P(A|B)$ = Probability of **A happening given that B has occurred** (posterior probability).
- $P(B|A)$ = Probability of **B happening given that A has occurred**.
- $P(A)$ = Probability of **A happening** (prior probability).
- $P(B)$ = Probability of **B happening** (total probability of evidence).

Dependent Probability Formula

When two events **depend** on each other, the probability of one event affects the probability of the other. The formula for **dependent probability** is:

$$P(B \text{ and } A) = P(A) \times P(B | A)$$

- $P(A \cap B)$ = Probability of **both A and B occurring**.

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Independent Probability Formula

Two events **A** and **B** are **independent** if the occurrence of one does **not affect** the probability of the other.

For **independent events**, the probability of both occurring is:

$$P(A \text{ and } B) = P(A) \times P(B)$$

Binominal Probability Formula

The Binomial Probability Formula is given as

$$P(x) = {}^nC_k \cdot p^x (1 - p)^{n-x}$$

$$P(x) = \left[\frac{n!}{x!(n-x)!} \right] \cdot p^x (1 - p)^{n-x}$$

Where,

n = Total number of events

x = Total number of successful events.

p = Success Probability in a single trial.

${}^nC_r = [n!/r!(n-r)!]$

$1 - p$ = Probability of failure.

Learn [Binomial Distribution](#)

Normal Probability Formula

The Normal probability formula is given by:

$$P(x) = (1/\sqrt{2\pi})e^{(-x^2/2)}$$

Learn [Normal Distribution](#)

Experimental Probability formula

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Probability $P(x)$ = Number of times an event occurs / Total number of trials.

Theoretical Probability Formula

The [Theoretical Probability](#) Formula is,

$P(x)$ = Number of Favorable outcomes/ Number of Possible outcomes.

Standard Deviation Probability Formula

The [Standard Deviation Probability](#) Formula is given as

$$P(x) = (1/\sigma\sqrt{2\pi})e^{-(x-\mu)^2/2\sigma^2}$$

Bernoulli Probability Formula

A random variable X will have a [Bernoulli Distribution](#) with probability p; the formula is,

$P(X = x) = p^x (1 - p)^{1-x}$, for $x = 0, 1$ and $P(X = x) = 0$ for other values of x

Here, 0 is failure and 1 is the success.

Probability Formula for Class 10

In Class 10, we have to study basic probability, such as the probability of tossing a coin, tossing 2 coins, tossing 3 coins, throwing a die, throwing two dice, probability of drawing a card from well well-shuffled deck. All these questions can be solved with only one formula. The Probability Formula Class 10 is given as

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

$n(E)$ is number of trials in which Event Occurred

$n(S)$ is number of Sample Space

Probability Formula for Class 12

The various formula used in Probability Class 12 is tabulated below:

Various Probability Formulas	
Experimental or Empirical Probability Formula	$P(E) = \text{Number of times an event occurs} / \text{Total number of trials.}$
Classical or Theoretical Probability Formula	$P(E) = \text{Number of Favorable Outcomes} / \text{Total Number of Possible Outcomes}$
Addition Probability Formula	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$
Joint Probability Formula	$P(A \cap B) = P(A) \cdot P(B)$
Addition Rule for Mutually Exclusive Events	$P(A \text{ or } B) = P(A) + P(B)$
Complementary Rule Formula	$P(\text{not } A) = 1 - P(A) \text{ or } P(A') = 1 - P(A).$ $P(A) + P(A') = 1$
Conditional Rule Formula	$P(B A) = P(A \cap B) / P(A)$
	$P(A) = \text{Number of Times Event A}$

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Various Probability Formulas	
Disjoint Event	$P(A \cap B) = 0$
Bayes' Theorem	$P(A B) = P(B A) \times P(A) / P(B)$
Dependent Probability Formula	$P(B \text{ and } A) = P(A) \times P(B A)$
Independent Probability Formula	$P(A \text{ and } B) = P(A) \times P(B)$
Binominal Probability Formula	$P(x) = {}^nC_x \cdot p^x (1 - p)^{n-x}$ or $P(r) = [n! / r!(n-r)!] \cdot p^r (1 - p)^{n-r}$
Normal Probability Formula	$P(x) = (1/\sqrt{2\pi}) e^{(-x^2/2)}$
Standard Deviation Probability Formula	$P(x) = (1/\sigma\sqrt{2\pi}) e^{-(x-\mu)^2/2\sigma^2}$
Bernoulli Probability Formula	$P(X = x) = p^x (1 - p)^{1-x}$, for $x = 0, 1$ and $P(X = x) = 0$ for other values of x .

Also, Check:

Probability Topics	
Coin Toss Probability	Card Probability
Tricks to Solve Probability Questions	Interesting Facts on Probability

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Probability Topics	
Probability Quiz	Real-Life Applications of Probability

Solved Examples on Probability Formulas

Example 1: Select a card at random from a standard deck. What is the probability of drawing a card with a feminine face?

Solution:

In a standard deck containing 52 cards: Total possible outcomes = 52

Event A = drawing a card with a feminine face

The number of favorable events (considering only queens as feminine faces) = 4

Therefore, the probability $P(A)$ is calculated using the formula:

$P(A) = \text{Number of Favorable Outcomes} \div \text{Total Number of Outcomes}$

$$P(A) = 4/52$$

$$P(A) = 1/13.$$

Example 2: If the Probability of event E, denoted as $P(E) = 0.35$, what is the probability of the complement event 'not E'?

Solution:

Given that $P(E) = 0.35$, we can use the complementary probability formula:

$$P(E) + P(\text{not } E) = 1$$

Substituting the known value:

$$P(\text{not } E) = 1 - P(E)$$

$$P(\text{not } E) = 1 - 0.35$$

$$\text{Hence, } P(\text{not } E) = 0.65$$

Example 3: Dangerous fires are very rare, around 1% but the smoke is

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Probability of dangerous Fire when there is smoke by using Bayes theorem:

$$P(\text{Fire}) = 0.01$$

$$P(\text{Smoke}) = 0.20$$

$$P(\text{Fire}|\text{Smoke}) = 0.80$$

$$P(\text{Fire}|\text{Smoke}) = \{P(\text{Fire})P(\text{Smoke}|\text{Fire})\}/P(\text{Smoke})$$

We can substitute these values:

$$P(\text{Fire} | \text{Smoke}) = (0.01 \times 0.80) / 0.20$$

$$P(\text{Fire} | \text{Smoke}) = 0.008 / 0.20$$

$$P(\text{Fire} | \text{Smoke}) = 0.04 = 4\%.$$

Example 4: Within a bag, there are 2 green bulbs, 4 orange bulbs, and 6 white bulbs. When a bulb is randomly chosen from the bag, what is the probability of picking either a green bulb or a white bulb?

Solution:

We are given a bag containing:

- **2 Green bulbs**
- **4 Orange bulbs**
- **6 White bulbs**
- **Total bulbs = 2 + 4 + 6 = 12**

*We need to find the probability of picking **either a green or a white bulb**.*

E = picking either a green bulb or a white bulb

P(E) = (Number of green bulbs + Number of white bulbs) / Total number of bulbs

$$P(E) = (2+6)/12$$

$$P(E) = 8/12$$

$$P(E) = 2/3.$$

Practice Questions on the Probability Formulas

Question 1. From a collection of marbles in a bag—8 red, 9 blue, and 6

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Question 2. In a drawer containing 6 black pens, 4 blue pens, and 7 red pens, a pen is drawn at random. What is the probability that the pen is either black or blue?

Question 3. Drawing one card from a thoroughly shuffled deck of 52 cards, determine the probability that the card will:

- Be a king.
- Not to be a king.

Question 4. According to a survey, 70% of individuals enjoy chocolate, and among those chocolate enthusiasts, 60% also have a liking for vanilla. What is the probability that an individual likes vanilla, given their fondness for chocolate?

Question 5. Determine the probability of rolling an odd number when a six-sided die is rolled.

[Comment](#)[More info](#)[Advertise with us](#)[Next Article](#)[Probability Formulas](#)

Similar Reads

1. [Probability in Maths](#)
2. [Card Probability](#)
3. [Experimental Probability](#)
4. [Probability | Set-2](#)
5. [Classical Probability in R](#)
6. [Probability of At Least One](#)
7. [Events in Probability](#)
8. [Chance and Probability](#)
9. [How to Calculate Probability](#)

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

**Corporate & Communications Address:**

A-143, 7th Floor, Sovereign Corporate Tower, Sector- 136, Noida, Uttar Pradesh (201305)

Registered Address:

K 061, Tower K, Gulshan Vivante Apartment, Sector 137, Noida, Gautam Buddh Nagar, Uttar Pradesh, 201305



Advertise with us

Company

About Us
Legal
Privacy Policy
Careers
In Media
Contact Us
Corporate Solution
Campus Training Program

Tutorials

Python
Java
C++
PHP
GoLang
SQL
R Language
Android

Explore

Job-A-Thon
Offline Classroom Program
DSA in JAVA/C++
Master System Design
Master CP
Videos

DSA

Data Structures
Algorithms
DSA for Beginners
Basic DSA Problems
DSA Roadmap
DSA Interview Questions
Competitive Programming

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

ML Maths
Data Visualisation
Pandas
NumPy
NLP
Deep Learning

JavaScript
TypeScript
ReactJS
NextJS
NodeJs
Bootstrap
Tailwind CSS

Python Tutorial

Python Examples
Django Tutorial
Python Projects
Python Tkinter
Web Scraping
OpenCV Tutorial
Python Interview Question

DevOps

Git
AWS
Docker
Kubernetes
Azure
GCP
DevOps Roadmap

School Subjects

Mathematics
Physics
Chemistry
Biology
Social Science
English Grammar

Preparation Corner

Company-Wise Recruitment Process
Aptitude Preparation
Puzzles
Company-Wise Preparation

Computer Science

GATE CS Notes
Operating Systems
Computer Network
Database Management System
Software Engineering
Digital Logic Design
Engineering Maths

System Design

High Level Design
Low Level Design
UML Diagrams
Interview Guide
Design Patterns
OOAD
System Design Bootcamp
Interview Questions

Databases

SQL
MYSQL
PostgreSQL
PL/SQL
MongoDB

More Tutorials

Software Development
Software Testing
Product Management
Project Management
Linux
Excel

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

DSA and Placements

Web Development

Data Science

Programming Languages

DevOps & Cloud

Clouds/Devops

DevOps Engineering

AWS Solutions Architect Certification

Salesforce Certified Administrator Course

C++ Programming Course

Java Programming Course

Python Full Course

GATE 2026

GATE CS Rank Booster

GATE DA Rank Booster

GATE CS & IT Course - 2026

GATE DA Course 2026

GATE Rank Predictor

@GeeksforGeeks, Sanchhaya Education Private Limited, All rights reserved

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).