


Bayes' Theorem

Last Updated : 26 Apr, 2025

Bayes' Theorem is a mathematical formula that helps determine the **conditional probability** of an event based on prior knowledge and new evidence.

It adjusts probabilities when new information comes in and helps make better decisions in uncertain situations.



Bayes' Theorem helps us update probabilities based on prior knowledge and new evidence. In this case, knowing that the pet is quiet (new information), we can use Bayes' Theorem to calculate the updated probability of the pet being a cat or a dog, based on how likely each animal is to be quiet.

Bayes Theorem and Conditional Probability

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Got It !

occurred.

The general statement of Bayes' theorem is "The conditional probability of an event A, given the occurrence of another event B, is equal to the product of the event of B, given A, and the probability of A divided by the probability of event B." i.e.

For example, if we want to find the probability that a white marble drawn at random came from the first bag, given that a white marble has already been drawn, and there are three bags each containing some white and black marbles, then we can use Bayes' Theorem.

Check: [Bayes's Theorem for Conditional Probability](#)

Bayes Theorem Formula

For any two events A and B, Bayes's formula for the Bayes theorem is given by:

$$P(A|B) = \frac{P(B|A) P(A)}{P(B)}$$

Formula for the Bayes theorem

Where,

Related searches

< Bayes Theorem Vs. Conditional Probability

Bayes Theorem Math Ia >

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- **P(A)** and **P(B)** are the probabilities of events A and B, also, P(B) is never equal to zero.
- **P(A|B)** is the probability of event A when event B happens,
- **P(B|A)** is the probability of event B when A happens.

Bayes Theorem Statement

Bayes's Theorem for n sets of events is defined as,

Let E_1, E_2, \dots, E_n be a set of events associated with the sample space S, in which all the events E_1, E_2, \dots, E_n have a non-zero probability of occurrence. All the events E_1, E_2, \dots, E form a partition of S. Let A be an event from space S for which we have to find the probability, then according to Bayes theorem,

$$P(E_i | A) = \frac{P(E_i) \cdot P(A|E_i)}{\sum_{k=1}^n P(E_k) \cdot P(A|E_k)}$$

for $k = 1, 2, 3, \dots, n$

Bayes Theorem Derivation

The proof of Bayes's, Theorem is given as, according to the conditional probability formula,

$$P(E_i | A) = \frac{P(E_i \cap A)}{P(A)} \dots \text{(i)}$$

Then, by using the multiplication rule of probability, we get

$$P(E_i \cap A) = P(E_i) \cdot P(A | E_i) \dots \text{(ii)}$$

Now, by the total probability theorem,

$$P(A) = \sum_{k=1}^n P(E_k) \cdot P(A | E_k) \dots \text{(iii)}$$

Substituting the value of $P(E_i \cap A)$ and $P(A)$ from eq (ii) and eq(iii) in eq(i) we get,

$$P(E_i | A) = \frac{P(E_i) \cdot P(A|E_i)}{\sum_{k=1}^n P(E_k) \cdot P(A|E_k)}$$

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Bayes' theorem is also known as the formula for the probability of “causes”. As we know, the E_i 's are a partition of the sample space S , and at any given time, only one of the events E_i occurs. Thus, we conclude that the Bayes theorem formula gives the probability of a particular E_i , given that event A has occurred.

Terms Related to Bayes Theorem

After learning about **Bayes** theorem in detail, let us understand some important terms related to the concepts we covered in the formula and derivation.

Hypotheses

- [Hypotheses](#) refer to possible events or outcomes in the sample space, they are denoted as E_1, E_2, \dots, E_n .
- Each hypothesis represents a distinct scenario that could explain an observed event.

Priori Probability

- [Priori Probability](#) $P(E_i)$ is the initial probability of an event occurring before any new data is taken into account.
- It reflects existing knowledge or assumptions about the event.
- **Example:** The probability of a person having a disease before taking a test.

Posterior Probability

- [Posterior probability](#) $P(E_i|A)$ is the updated probability of an event after considering new information.
- It is derived using the Bayes Theorem.
- **Example:** The probability of having a disease given a positive test result.

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

- The probability of an event A based on the occurrence of another event B is termed [conditional Probability](#).
- It is denoted as $P(A|B)$ and represents the probability of A when event B has already happened.

Joint Probability

- When the probability of two or more events occurring together and at the same time is measured, it is marked as [Joint Probability](#).
- For two events A and B, it is denoted by joint probability is denoted as $P(A \cap B)$.

Random Variables

- Real-valued variables whose possible values are determined by random experiments are called [random variables](#).
- The probability of finding such variables is the experimental probability.

Bayes Theorem Applications

Bayesian inference is very important and has found application in various activities, including medicine, science, philosophy, engineering, sports, law, etc., and Bayesian inference is directly derived from Bayes theorem.

Some of the Key Applications are:

- **Medical Testing** → Finding the real probability of having a disease after a positive test.
- **Spam Filters** → Checking if an email is spam based on keywords.
- **Weather Prediction** → Updating the chance of rain based on new data.
- **AI & Machine Learning** → Used in [Naïve Bayes classifiers](#) to predict outcomes.

Check, [Bayes' Life Applications of Bayes theorem](#)

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Difference Between Conditional Probability and Bayes Theorem

The difference between Conditional Probability and Bayes's. The theorem can be understood with the help of the table given below.

Bayes Theorem	Conditional Probability
Bayes's Theorem is derived using the definition of conditional probability. It is used to find the reverse probability.	Conditional Probability is the probability of event A when event B has already occurred.
Formula: $P(A B) = [P(B A)P(A)] / P(B)$	Formula: $P(A B) = P(A \cap B) / P(B)$
Purpose: To update the probability of an event based on new evidence.	Purpose: To find the probability of one event based on the occurrence of another.
Focus: Uses prior knowledge and evidence to compute a revised probability.	Focus: Direct relationship between two events.

Theorem of Total Probability

Let E_1, E_2, \dots, E_n be **mutually exclusive and exhaustive events** of a sample space S , and let E be any event that occurs with some E_i . Then, prove that :

$$P(E) = \sum_{i=1}^n P(E/E_i) \cdot P(E_i)$$

Proof:

Let S be the sample space

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

$S = E_1 \cup E_2 \cup E_3 \cup \dots \cup E_n$ and $E_i \cap E_j = \emptyset$ for $i \neq j$.

Now, consider the event $E: E = E \cap S$

Substituting S with the union of E_i 's:

$$\Rightarrow E = E \cap (E_1 \cup E_2 \cup E_3 \cup \dots \cup E_n)$$

Using distributive law:

$$\Rightarrow E = (E \cap E_1) \cup (E \cap E_2) \cup \dots \cup (E \cap E_n)$$

Since the events E_i are mutually exclusive, the intersections $E \cap E_i$ are also **mutually exclusive**. Therefore:

$$P(E) = P\{(E \cap E_1) \cup (E \cap E_2) \cup \dots \cup (E \cap E_n)\}$$

$$\Rightarrow P(E) = P(E \cap E_1) + P(E \cap E_2) + \dots + P(E \cap E_n)$$

{Therefore, $(E \cap E_1), (E \cap E_2), \dots, (E \cap E_n)$ are pairwise disjoint}

$$\Rightarrow P(E) = P(E/E_1) \cdot P(E_1) + P(E/E_2) \cdot P(E_2) + \dots + P(E/E_n) \cdot P(E_n) \text{ [by [multiplication theorem](#)]}$$

$$\Rightarrow P(E) = \sum_{i=1}^n P(E/E_i) \cdot P(E_i)$$

Articles Related to Bayes' Theorem-

- [Conditional Probability](#)
- [Probability Distribution](#)

Bayes Theorem for Programmers-

- [Bayes Theorem in Artificial Intelligence](#)
- [Bayes Theorem in Machine Learning](#)
- [Bayes' Theorem in Data Mining](#)

Solved Examples of Bayes's Theorem

Example 1: A person has undertaken a job. The probabilities of completion of the job on time with and without rain are 0.44 and 0.9, and 5, respectively. If the probability that it will rain is 0.45, then determine the probability that the job will be completed on time.

Solution:

Let E_1 be the event that the mining job will be completed on time

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

*By multiplication law of probability,
 $P(E_1) = 0.44$, and $P(E_2) = 0.95$*

Since, events A and B form partitions of the sample space S, by total probability theorem, we have

$$\begin{aligned} P(E) &= P(A) P(E_1) + P(B) P(E_2) \\ \Rightarrow P(E) &= 0.45 \times 0.44 + 0.55 \times 0.95 \\ \Rightarrow P(E) &= 0.198 + 0.5225 = 0.7205 \end{aligned}$$

So, the probability that the job will be completed on time is 0.7205

Example 2: There are three urns containing 3 white and 2 black balls, 2 white and 3 black balls, and 1 black and 4 white balls, respectively. There is an equal probability of each urn being chosen. One ball is equal probability chosen at random. What is the probability that a white ball will be drawn?

Solution:

*Let E_1 , E_2 , and E_3 be the events of choosing the first, second, and third urn respectively. Then,
 $P(E_1) = P(E_2) = P(E_3) = 1/3$*

*Let E be the event that a white ball is drawn. Then,
 $P(E/E_1) = 3/5$, $P(E/E_2) = 2/5$, $P(E/E_3) = 4/5$*

*By theorem of total probability, we have
 $P(E) = P(E/E_1) \cdot P(E_1) + P(E/E_2) \cdot P(E_2) + P(E/E_3) \cdot P(E_3)$
 $\Rightarrow P(E) = (3/5 \times 1/3) + (2/5 \times 1/3) + (4/5 \times 1/3)$
 $\Rightarrow P(E) = 9/15 = 3/5$*

Example 3: A card from a pack of 52 cards is lost. From the remaining cards of the pack, two cards are drawn and are found to be both hearts. Find the probability of the lost card being a heart.

Solution:

Let E_1 , E_2 , E_3 , and E_4 be the events of losing a card of hearts, clubs, spades, and diamonds respectively.

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

$P(E|E_1)$ = probability of drawing 2 hearts, given that a card of hearts is missing

$$\Rightarrow P(E|E_1) = {}^{12}C_2 / {}^{51}C_2 = (12 \times 11) / 2! \times 2! / (51 \times 50) = 22/425$$

$P(E|E_2)$ = probability of drawing 2 clubs, given that a card of clubs is missing

$$\Rightarrow P(E|E_2) = {}^{13}C_2 / {}^{51}C_2 = (13 \times 12) / 2! \times 2! / (51 \times 50) = 26/425$$

$P(E|E_3)$ = probability of drawing 2 spades, given that a card of hearts is missing

$$\Rightarrow P(E|E_3) = {}^{13}C_2 / {}^{51}C_2 = 26/425$$

$P(E|E_4)$ = probability of drawing 2 diamonds, given that a card of diamonds is missing

$$\Rightarrow P(E|E_4) = {}^{13}C_2 / {}^{51}C_2 = 26/425$$

Therefore,

$P(E_1|E)$ = probability of the lost card is being a heart, given the 2 hearts are drawn from the remaining 51 cards

$$\Rightarrow P(E_1|E) = P(E_1) \cdot P(E|E_1) / P(E_1) \cdot P(E|E_1) + P(E_2) \cdot P(E|E_2) + P(E_3) \cdot P(E|E_3) + P(E_4) \cdot P(E|E_4)$$

$$\Rightarrow P(E_1|E) = (1/4 \times 22/425) / \{(1/4 \times 22/425) + (1/4 \times 26/425) + (1/4 \times 26/425) + (1/4 \times 26/425)\}$$

$$\Rightarrow P(E_1|E) = 22/100 = 0.22$$

Hence, The required probability is 0.22.

Example 4: Suppose 15 men out of 300 men and 25 women out of 1000 are good orators. An orator is chosen at random. Find the probability that a male person is selected.

Solution:

Given,

- Total Men = 300
- Total Women = 1000
- Good Orators among Men = 15
- Good Orators among Women = 25

$$P(\text{Male Orator}) = \text{Numbers of male orators} / \text{total no of orators} = 15/40 = 3/8$$

Example 5: A man is known to speak the lies 1 out of 4 times. He throws a die and reports that it is a six. Find the probability that it **Bayes'** actually a six.

Solution:

In a throw of a die, let

E_1 = event of getting a six,

E_2 = event of not getting a six and

E = event that the man reports that it is a six.

Then, $P(E_1) = 1/6$, and $P(E_2) = (1 - 1/6) = 5/6$

$P(E|E_1)$ = probability that the man reports that six occurs when six has actually occurred

$\Rightarrow P(E|E_1)$ = probability that the man speaks the truth

$\Rightarrow P(E|E_1) = 3/4$

$P(E|E_2)$ = probability that the man reports that six occurs when six has not actually occurred

$\Rightarrow P(E|E_2)$ = probability that the man does not speak the truth

$\Rightarrow P(E|E_2) = (1 - 3/4) = 1/4$

Probability of getting a six ,given that the man reports it to be six

$P(E_1|E) = P(E|E_1) \times P(E_1) / P(E|E_1) \times P(E_1) + P(E|E_2) \times P(E_2)$ [by Bayes theorem]

$\Rightarrow P(E_1|E) = (3/4 \times 1/6) / \{(3/4 \times 1/6) + (1/4 \times 5/6)\}$

$\Rightarrow P(E_1|E) = (1/8 \times 3) = 3/8$

Hence the probability required is $3/8$.

Comment

More info

Advertise with us

Next Article

Probability Distribution - Function,
Formula, Table

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

1. Bayes' Theorem in Data Mining
2. Probability Theory
3. Bayes Theorem in Machine learning
4. Theoretical Probability
5. Nyquist Sampling Theorem
6. Bernoulli Naive Bayes
7. Mid Point Theorem
8. Multiplication Theorem
9. Real Life Applications of Bayes' Theorem
10. Multinomial Naive Bayes

**Corporate & Communications Address:**

A-143, 7th Floor, Sovereign Corporate Tower, Sector- 136, Noida, Uttar Pradesh (201305)

Registered Address:

K 061, Tower K, Gulshan Vivante Apartment, Sector 137, Noida, Gautam Buddh Nagar, Uttar Pradesh, 201305



Advertise with us

Company

About Us

Legal

Explore

Job-A-Thon

Offline Classroom Program

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Corporate Solution
Campus Training Program

Tutorials

Python
Java
C++
PHP
GoLang
SQL
R Language
Android

Data Science & ML

Data Science With Python
Machine Learning
ML Maths
Data Visualisation
Pandas
NumPy
NLP
Deep Learning

Python Tutorial

Python Examples
Django Tutorial
Python Projects
Python Tkinter
Web Scraping
OpenCV Tutorial
Python Interview Question

DevOps

Git
AWS
Docker
Kubernetes
Azure
GCP
DevOps Roadmap

DSA

Data Structures
Algorithms
DSA for Beginners
Basic DSA Problems
DSA Roadmap
DSA Interview Questions
Competitive Programming

Web Technologies

HTML
CSS
JavaScript
TypeScript
ReactJS
NextJS
NodeJs
Bootstrap
Tailwind CSS

Computer Science

GATE CS Notes
Operating Systems
Computer Network
Database Management System
Software Engineering
Digital Logic Design
Engineering Maths

System Design

High Level Design
Low Level Design
UML Diagrams
Interview Guide
Design Patterns
OOAD
System Design Bootcamp
Interview Questions

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).

Chemistry
Biology
Social Science
English Grammar

PostgreSQL
PL/SQL
MongoDB

Preparation Corner

Company-Wise Recruitment Process
Aptitude Preparation
Puzzles
Company-Wise Preparation

More Tutorials

Software Development
Software Testing
Product Management
Project Management
Linux
Excel
All Cheat Sheets

Courses

IBM Certification Courses
DSA and Placements
Web Development
Data Science
Programming Languages
DevOps & Cloud

Programming Languages

C Programming with Data Structures
C++ Programming Course
Java Programming Course
Python Full Course

Clouds/Devops

DevOps Engineering
AWS Solutions Architect Certification
Salesforce Certified Administrator Course

GATE 2026

GATE CS Rank Booster
GATE DA Rank Booster
GATE CS & IT Course - 2026
GATE DA Course 2026
GATE Rank Predictor

@GeeksforGeeks, Sanchhaya Education Private Limited, All rights reserved

We use cookies to ensure you have the best browsing experience on our website. By using our site, you acknowledge that you have read and understood our [Cookie Policy](#) & [Privacy Policy](#).