

Seeing the Invisible

Physics-Guided Missing-Physics Learning for PEMFC Flooding Diagnosis

Ganghyun Shin, Jongjin Jang, Geonho Shin, Jaehyeon Ryu, Seoyeon Kim, Minseo Baek, Minji Lee, Chanseo Lee, Keeyoun Yoo*

Department of Chemical and Biomolecular Engineering, Seoul National University of Science and Technology




ABSTRACT

Liquid water accumulation in porous gas diffusion layers critically limits mass transport in polymer electrolyte membrane fuel cells, yet the internal flooding state is difficult to observe directly. More broadly, physics-based multiphysics models often rely on empirical or fitted constitutive relations to represent unresolved transport behavior, making constitutive uncertainty a major source of prediction error. Here, we propose a physics-informed grey-box framework that retains the known governing equations within a finite-element solver while learning only the missing constitutive physics. A learnable gas-phase transport closure is embedded directly into the two-phase FEM model and identified from synthetic reference responses using discrete-adjoint sensitivities, without prescribing a fixed functional form in advance. After training at RH 50%, the learned constitutive law is frozen and evaluated at unseen RH 70% on two discretizations, 16×40 and 32×80 . Agreement with the synthetic reference is retained without retraining in the reported test cases, supporting transfer of the learned closure across the tested humidity and discretization changes. These synthetic tests motivate future experimental validation for flooding diagnosis and physics-guided digital twins.

INTRODUCTION

A Why flooding matters



B Challenges in flooding diagnosis


Observation	Internal liquid-water saturation $s(x,y)$ cannot be directly and continuously observed under realistic operating conditions.
Modeling	Flooding predictions depend strongly on uncertain constitutive relations such as $D_{eff}(s)$, $k_r(s)$, and $p_c(s)$.

C Existing ML vs our objective

DIFFERENT LEARNING TARGETS, NOT A BETTER/WORSE RANKING


PINN Learn PDE solution Physics is enforced mainly through residual-based training.	FNO / Neural Operator Learn solution operator Maps input fields or parameters to solution fields.	OUR APPROACH FEM + Missing Physics ML Learn the unknown physical relation Known conservation laws remain in FEM; only uncertain constitutive physics is learned.
--	--	---

METHODS




RESULTS


01 Model domain & reference conditions




02 Water accumulation grows at high current



03 Recover the missing transport relation



04 Zero-shot transfer with a frozen closure



Zero-shot transfer: the same frozen closure reproduces O_2 transport and localized water accumulation on both meshes. Reference–FEM pairs share scales; error maps show pointwise absolute differences. This tests discretization transfer, not mesh convergence. x is stretched for display. Field errors: relative nodal l_2 norm. Current error: relative absolute difference. All errors compare frozen FEM with the synthetic reference.

Conclusions

- 01 Physics-preserving learning.** Governing equations remain in FEM; only the uncertain gas-phase closure $g_\theta(\phi_g)$ is learned.
- 02 Constitutive recovery.** 2.16% relative L^2 error at epoch 179 over the evaluated support.
- 03 Frozen transfer.** Agreement at RH 70% on both 16×40 and 32×80 , without retraining.

REFERENCES [1] N. Bouziani, D. A. Ham & A. Farsi, Differentiable programming across the PDE and Machine Learning barrier, arXiv:2409.06085 (2024). [2] A. Farsi, N. Bouziani & D. A. Ham, Missing physics discovery through fully differentiable finite element-based machine learning, Scientific Reports (2026), doi:10.1038/s41598-026-65125-z. [3] M. Abdollahzadeh et al. Analysis of PEM fuel cell cathode two-dimensional modeling. Energy 68, 478–494 (2014).