Affine detector counting with asymmetric source savings

Zhongpai Gao
With AI assistance; predecessor contributions acknowledged below

14 October 2026

Canonical proof checked locally; tracker admission pending.

Abstract

We prove nonvanishing of every Dirichlet L-function in $\Re s > 21871/25000 = 0.874840$, away from the principal pole. The construction combines timbalud's asymmetric marked HIGH count and common-radical saving with DaysSky's use of Collet's initial affine moment inside the detector count. The stronger source moment increases the detector's available prime-slot width. A small change of the height junction preserves the actual source gates. We give the counting formulas, the allocation argument, and the passage from uniform scalar inequalities to the inherited analytic continuation argument. Exact rational certificates cover the continuous parameter ranges, and the complete source assembly has passed canonical comparison and all three kernels.

1 Framework and theorem

Let $\beta \le 7/8$ be the supremum of the nontrivial zero real parts in Cycle61's closed Eisenstein Hecke family. The upper bound and the transfer from this family to all Dirichlet characters are inherited from the OpenAI and Collet framework [1, 5]. Set

$$\vartheta = \frac{21871}{25000}, \quad h = \frac{1989}{2500}, \quad d_- = \frac{159}{200}, \quad d_+ = h + 10^{-10}, \quad \kappa_* = 2\vartheta - 1 = \frac{9371}{12500}, \tag{1}$$
$$\ell = \frac{11}{3} - 4\vartheta = \frac{3137}{18750}, \quad x = 1 + \ell - h = \frac{13939}{37500}, \quad y = h - 2\ell = \frac{17287}{37500}, \quad \lambda = \frac{19312}{4787}, \quad B = \frac{14200}{14361} + \frac{7}{10000}.$$

Theorem. For every modulus $q \ge 1$, every Dirichlet character χ modulo q, and every $s \in \mathbb{C}$,

$$\vartheta < \Re s, \qquad \neg(\chi = 1 \text{ and } s = 1) \quad \Longrightarrow \quad L(\chi, s) \neq 0.$$

The accompanying Lean construction proves this canonical statement.

Assume $\beta > \vartheta$, keeping β as the global supremum. The continuation contract first chooses a common error reserve $\sigma > 0$ and $\omega = (\beta - \vartheta)/2$, before quantifying over primitive targets. Constants and a small height exponent may depend on the target character; the resulting positive continuation gap must remain uniform. This order is essential when the affine moment is applied to detector characters.

2 A stronger actual source moment

Collet's initial affine moment bounds the normalized plain energy, including its selected prime factors, by

$$\mathcal{E}_\eta(U; m, s) \ll_{\eta,\varepsilon} (1+T)^{D_\eta}\, \mathcal{S}(\Phi)^2 U^{1+\varepsilon}, \qquad 2m + \lambda s \le B. \tag{2}$$

Here m is the plain length, s the selected logarithmic prime width, T controls the shifts, and $\mathcal{S}$ is a finite profile seminorm. All fixed masks, relative characters and source normalizations are those of the actual initial moment. We use (2) only in its proved window, with $s \le 1/25$ and m in the interval specified in Section 3.

The admissible choice of B is justified by the complete marked source argument of timbalud [2], not by an additional moment hypothesis. The useful common-allocation step can be seen directly. For first-allocation coordinates c, d, radical coordinate R, cancellation coordinate R_0 and retained width L_c, the actual source contains

$$F = (1-\rho)(c-d)_+ - \delta c - (1+\rho)R - E - \frac{(3c - 5d - R)_+}{6} + \rho L_c + R_0. \tag{3}$$

The radical penalty is tied to the same c, d, R; it cannot be discarded and then varied independently. Under the literal gates $R \le \min(c, d)$, $L_c \le 5(R - R_0)$, $E, R_0, c, d \ge 0$ and $1/4 \le \rho \le 2/5$, one has

$$F \le -(\delta - 3\rho/2)c - E. \tag{4}$$

Indeed, for $d \le c$, write $a = 2 - 5\rho \in [0, 1]$ and use $ad + (c/2 - d)_+ \ge ac/2$. For $c \le d$, use $R \le c$ and $1 - 5\rho/2 \ge 0$. These two cases give (4) on the whole domain.

In the marked HIGH range $17/25 \le \delta \le 7/10$, $\rho = 1 - 10\delta/[3(5 - 2\delta)] \le 103/273$; thus the first removed width has rate at least $a_0 = 519/4550$. The second allocation has the stronger direct rate δ. With M the source scale, let $u, f, v \ge 0$ be the common, residual frozen and surviving widths, and let Q be the actual remaining frequency scale. The source identities and two-stage collision accounting give

$$u + f + v \ge \frac{33M}{1000}, \qquad Q \ge \frac{1121M}{1250} - 2u.$$

Asymmetric inverse/plain selections supply the live-width payment, so that

$$a_0 u + \delta f + \frac{224}{1755} \min\left(v, \frac{13Q}{400}\right) \ \ge \ \frac{37M}{10000}. \tag{5}$$

For $v \le 13Q/400$, use the width supply and the smallest of the three rates. In the other case insert the lower bound for Q; the coefficient of u remains positive and the constant term already exceeds the right side. This proves (5) without choosing an allocation extremizer. The complementary large sixth-power tail is paid by its inherited bound. Charging at most $M/20000$ leaves $73M/20000$. These savings are retained inside the actual source moment, with moving common ideals, frozen coefficients and encoded-frequency collisions unchanged.

3 Applying that moment inside the detector

A detector fiber has inverse length r, plain length m, cutoff t, zero density δ and normalized prime amplitude $p \in [0, 1/2]$. Up to the explicitly charged mesh errors, $r + m = t$. The three relevant count exponents are

$$\begin{aligned} I(\delta, p, r) &= 1 - \delta\big(p + (1 - p)r\big), \\ A(\delta, p, m) &= 1 - 2\delta m - 2\delta p \frac{B - 2m}{\lambda}, \\ L(\delta, t) &= 1 - \delta + (5/6 - \delta)(t - 1). \end{aligned} \tag{6}$$

The affine plain branch uses (2) at width $(B - 2m)/\lambda$, with a small subtraction for rounding. It improves on the old plain width $(1 - 2m)/(6\kappa)$ on the critical fibers.

This is a legitimate instance of the *actual* source theorem. Fix a final target character ψ, and write η_j for its detector dictionary characters. Each η_j is represented as its own natural row, with the exact excluded-prime support and fixed masks. The generic paid main and error families apply to that row, and the literal energy identity identifies their energy with the fiber energy. The marked HIGH mesh cap is intersected with the original analytic mesh before the slot partition is chosen. This dictionary is finite: we take the maximum of its moment degrees and the union of its seminorm sets, then choose a common small height exponent for all its labels. The resulting exponent τ_ψ may depend on the final target ψ. No affine moment or uniform degree over infinitely many target characters is assumed.

Put

$$\begin{aligned} D(p) &= 3 - p - 4p/\lambda, \qquad & a(p) &= 2 - 4p/\lambda, \qquad & c(p) &= 1 - 2pB/\lambda, \\ J(\delta, p) &= (5/6 - \delta)D(p) + \delta(1 - p)a(p). \end{aligned}$$

Solving $I = A$ at $m = t - r$ and then balancing with L gives

$$\begin{aligned} r_N(p, t) &= \frac{a(p)t + 2pB/\lambda - p}{D(p)}, \qquad t_N = 1 + \frac{\delta(1 - p)c(p)}{J(\delta, p)}, \\ R_{\mathrm{aff}}(\delta, p) &= 1 - \delta + \frac{(5/6 - \delta)\delta(1 - p)c(p)}{J(\delta, p)}. \end{aligned} \tag{7}$$

Use this count on

$$\mathcal{R} = [89/200, 97/200] \times [199/400, 1/2], \qquad m \in [m_0, 847/2000], \quad m_0 = (B - \lambda/25)/2.$$

The exact region certificates give $D, J > 0$, $1 \le t_N \le 3/2$, $23/37 \le r_N(p, t_N) \le 1$, and $t_N - r_N(p, t_N) \ge m_0 + 10^{-5}$. For selected widths below $17/500$, the old κ-marked moment is used; the affine theorem is invoked only at or above this lower gate. For plain lengths above $847/2000$, the old marked plain branch gives the same short exponent. The original unmarked branches cover the remaining cases. Thus the complete fiber argument gives (7), with the original charged losses. The detector spacing is chosen $\varepsilon \le 10^{-5}$; the actual baseline constructor already gives $\varepsilon \le 1/200000$.

4 Closing every row and returning to the L-function

For $\kappa \in [\kappa_*, 3/4]$, write

$$C_\kappa(p) = \frac{2(1-p)(3\kappa - p)}{3\kappa(3-p) - 2p}, \qquad R_\kappa(\delta, p) = 1 - \delta + \frac{\delta C_\kappa(p)(5/6 - \delta)}{2(5/6 - \delta + \delta C_\kappa(p))}.$$

Let R^*_κ equal R_{aff} on $\mathcal{R}$ and R_κ elsewhere. The affine region count is at most the old count at κ_*, and R^*_κ is nonincreasing in p. This monotonicity is used when actual amplitude classes are transferred through the numerator masks.

The complete normalized reconstruction has row exponent

$$E_\kappa(d, C; \delta, p) = \frac{1+\delta}{2} + \frac{13h}{75} - \frac{y}{2} - \frac{\ell}{2} + \ell\delta p + d\left(\frac{1}{4} + \frac{3}{4}R^*_\kappa(\delta, p) - \frac{17}{50}\right) - \frac{\delta p C}{2}. \tag{8}$$

Its two physical capacities are

$$C_{\mathrm{new}}(d) = \frac{2y - (2-B)d}{\lambda}, \qquad C_{\mathrm{old},\kappa}(d) = \frac{\min(d, 2y - d)}{6\kappa}.$$

With (1), rational Bernstein certificates on closed rectangles, together with affine interpolation in d, establish

$$E_{\kappa_*}(d, C_{\mathrm{new}}(d); \delta, p) < \vartheta - \frac{4999}{5000000000}, \qquad d_- \le d \le d_+, \tag{9}$$

$$E_{\kappa_*}(d, C_{\mathrm{old},\kappa_*}(d); \delta, p) < \vartheta - \frac{1}{100000}, \qquad 1/200 \le d \le d_-. \tag{10}$$

Both inequalities hold for *every* $\delta \in [0, 3/4]$ and $p \in [0, 1/2]$. Below $d = 1/2$ the old saving is at least $1/1000$. For clarity, the new top row uses the affine count on $\mathcal{R}$ and the old count on its complement. The old half-height and junction rows are certified on the whole rectangle. Positive rational Bernstein coefficients certify the full polynomials, rather than samples of them; the corresponding cleared denominators are positive. The extension from h to $h + 10^{-10}$ costs less than $2 \cdot 10^{-10}$, giving (9). At d_- the new capacity exceeds the old capacity, so the new junction inherits (10).

The junction $d_- = h - 3/5000$ also respects the actual source gates. In particular the plain scale $n = 1 - y/d + t$, for $0 \le t \le 1/1000$, satisfies $41/100 < n < 264/625$; its affine capacity lies in $(17/500, 37/1000)$; and $0 \le 2y/d - 1 < 4/25$. The fixed source HIGH, LOW, amplifier and tail reserves, as well as the cube-floor payment, therefore remain available from the inherited complete estimates. These gates are checked alongside the row certificates, not inferred from their positivity.

Finally set $\kappa = 2\beta - 1$. The explicit feedback estimates strengthen (9)–(10) to

$$E_{2\beta-1}(d, C_{\mathrm{new}}; \delta, p) - \beta < -\frac{4999}{5000000000} - \frac{24}{25}(\beta - \vartheta),$$
$$E_{2\beta-1}(d, C_{\mathrm{old},2\beta-1}; \delta, p) - \beta < -\frac{1}{100000} - \frac{3}{4}(\beta - \vartheta).$$

The finite reconstruction, mesh and height costs can consequently be chosen below a fixed fraction of $\beta - \vartheta$. The unchanged source masks, prime products, transport and regularizer yield a common error reserve $\sigma > 0$. Assembly fixes this reserve and $\omega = (\beta - \vartheta)/2$ before choosing each target's height. Thus the Mellin continuation gap is

$$\varepsilon_{\mathrm{cont}} = \min\left(\frac{\beta - \vartheta}{2}, \sigma\right) > 0,$$

independent of the primitive target. Only the constants, regularizer and height choice depend on that target. The inherited continuation theorem excludes every primitive nontrivial zero in $\Re s > \beta - \varepsilon_{\mathrm{cont}}$, contradicting the global supremum. Hence $\beta \le \vartheta$, and the Dirichlet transfer proves the theorem.

5 Scope, provenance and verification

The mathematical advance is the use of a stronger, already source-paid affine moment *inside* the detector. The proof does not add the two predecessor improvements as numerical boundary gains. The first improvement changes the admissible moment; the second uses that moment in a separate counting problem. Their interaction changes the crossing formulas and the admissible prime-amplitude region, which is why the complete row certificates and source gates must be regenerated together.

The accompanying Lean construction exposes no new analytic premise in its final target. Its route is

$$\begin{aligned}\text{actual per-label affine energy} &\longrightarrow \text{actual detector count} \longrightarrow \\ \text{finite numerator and source probe} &\longrightarrow \text{canonical Dirichlet nonvanishing.}\end{aligned}$$

The final continuation contract has the order $\exists \omega, \sigma > 0\ \forall \psi\ \exists \tau_\psi$, with the displayed common gap. For each primitive target ψ, its finite detector dictionary shares τ_ψ after taking the stated finite maxima; character-dependent heights cannot shrink the continuation gap. The scalar argument above explains the new mechanism. The analytic reductions are supplied by the complete inherited source theorems.

Verification status. The canonical theorem at 21871/25000 was accepted locally for source commit `74643447f52a`. Comparator accepted the canonical statement with only `Classical.choice`, `Quot.sound` and `propext`; Lean, NanoDa and con-ron each accepted the complete proof. All 897 candidate modules were compiled from source in the parent replay. The final, metadata-only head reused an authenticated local compilation cache and ran fresh controls, canonical exports and all three kernels. Exact rational checks also cover the displayed continuous rows and physical gates. A sanitized receipt accompanies this manuscript. Official tracker admission remains pending.

Attribution. OpenAI supplies the protected 7/8 framework and library. Hailey Collet supplies Cycle25/Cycle61, the initial affine moment, and the complete physical source and continuation architecture. The weighted numerator uses Akash Levy's contribution and its inherited credits. Timbalud supplies the marked HIGH count and the asymmetric/common-radical source improvement [2]. DaysSky supplies the actual dictionary-character affine detector construction and its height order [3]. Liqsweep/blueberryvertigo also recorded the count-side affine idea and related arithmetic in PR52, whose Lean integration was unfinished [4]; that proposed endpoint is not used as a proved input. Zhongpai Gao, with AI assistance, combines the source mechanisms, chooses (1), regenerates the exact certificates and carries the changed source interfaces through the final assembly. Existing OpenAI, Collet, Levy and other predecessor notices are retained. The integration and manuscript are released under Apache-2.0, subject to the retained predecessor and dependency licenses.

References

[1] Hailey Collet, *Cycle61: all-Dirichlet boundary* 0.874861, complete source and manuscript, QuasiRiemannTracker PR42 (2026).

[2] timbalud, *Asymmetric marked counting with coupled radical allocation*, source at commit `8cb265dc3490`, QuasiRiemannTracker PR50 (2026).

[3] DaysSky, *Cycle61 with the affine moment in the detector count*, source at commit `67edfb7a1161`, QuasiRiemannTracker PR56 (2026).

[4] liqsweep / blueberryvertigo, *Collet's affine moment in the detector count*, related arithmetic proposal, QuasiRiemannTracker PR52 (2026).

[5] OpenAI, *Quasi-Riemann hypothesis formalization*, protected source snapshot `fd4aeeb2`, `https://github.com/OpenAI/math`. See the retained source notices for Akash Levy's weighted numerator and the full dependency attribution.