



# Increased Microbial Diversity and Decreased Prevalence of Common Pathogens in the Gut Microbiomes of Wild Turkeys Compared to Domestic Turkeys

Julia Craft,<sup>a</sup> Hyrum Eddington,<sup>b</sup> Nicholas D. Christman,<sup>a</sup> Weston Pryor,<sup>a</sup>  John M. Chaston,<sup>b</sup>  David L. Erickson,<sup>a</sup>  Eric Wilson<sup>a</sup>

<sup>a</sup>Department of Microbiology and Molecular Biology, Brigham Young University, Provo, Utah, USA

<sup>b</sup>Department of Plant and Wildlife Sciences, Brigham Young University, Provo, Utah, USA

**ABSTRACT** Turkeys (*Meleagris gallopavo*) provide a globally important source of protein and constitute the second most important source of poultry meat in the world. Bacterial diseases are common in commercial poultry production, causing significant production losses for farmers. Due to the increasingly recognized problems associated with large-scale/indiscriminate antibiotic use in agricultural settings, poultry producers need alternative methods to control common bacterial pathogens. In this study, we compared the cecal microbiota of wild and domestic turkeys, hypothesizing that environmental pressures faced by wild birds may select for a disease-resistant microbial community. Sequence analyses of 16S rRNA genes amplified from cecal samples indicate that free-roaming wild turkeys carry a rich and variable microbiota compared to domestic turkeys raised on large-scale poultry farms. Wild turkeys also had very low levels of *Staphylococcus*, *Salmonella*, and *Escherichia coli* compared to domestic turkeys. *E. coli* strains isolated from wild and domestic turkey cecal samples also belong to distinct phylogenetic backgrounds and differ in their propensity to carry virulence genes. *E. coli* strains isolated from factory-raised turkeys were far more likely to carry genes for capsule (*kpsII* and *kpsIII*) or siderophore (*iroN* and *fyuA*) synthesis than were those isolated from wild turkeys. These results suggest that the microbiota of wild turkeys may provide colonization resistance against common poultry pathogens.

**IMPORTANCE** Due to the increasingly recognized problems associated with antibiotic use in agricultural settings, poultry producers need alternative methods to control common bacterial pathogens. In this study, we compare the microbiota of wild and domestic turkeys. The results suggest that free-ranging wild turkeys carry a distinct microbiome compared to farm-raised turkeys. The microbiome of wild birds contains very low levels of poultry pathogens compared to that of farm-raised birds. The microbiomes of wild turkeys may be used to guide the development of new ways to control disease in large-scale poultry production.

**KEYWORDS** *Escherichia coli*, microbiome, poultry production, wild turkey, poultry pathogens

Turkeys (*Meleagris gallopavo*) evolved approximately 11 million years ago and are one of the first birds domesticated in the Americas (1–3). Although domesticated thousands of years ago, turkeys have remained generally very similar to their wild relatives until relatively recently (4, 5). In the past ~70 years, intensive selective breeding of turkeys has resulted in dramatic changes in commercially raised birds compared to their wild relatives, leading to a genome that is much less diverse than those of many other agricultural species (4). These genetic changes as well as advancements in production practices have resulted in domestic birds maturing much more quickly and reaching three times the body mass of wild birds at maturity (6). Domestic turkeys are now the

**Editor** Martha Vives, Universidad de los Andes

**Copyright** © 2022 Craft et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Eric Wilson, [ericwilson@BYU.edu](mailto:ericwilson@BYU.edu).

The authors declare no conflict of interest.

**Received** 16 July 2021

**Accepted** 1 December 2021

**Accepted manuscript posted online**

19 January 2022

**Published** 8 March 2022

second most important source of poultry in the world, with the United States producing ~250,000,000 turkeys and ~7,000,000,000 pounds of turkey meat in 2019 (7).

Relatively few studies have been published comparing the microbiomes of wild animals and their domesticated kin. However, the limited literature on this topic has overwhelmingly shown that the microbiomes of captive and wild animals vary dramatically (8–15). The observed differences in microbial communities between wild and captive animals have led to calls for more research on the microbiomes of additional wild animals (16, 17).

The gut microbiome of poultry is known to contribute to efficient growth as well as bird health (11, 18–21). The microbiome of commercially raised poultry is undoubtedly influenced by production practices such as crowded conditions, diet, and antibiotic use. Several studies have characterized the gut microbiomes of domestic turkeys in a variety of experimental and agricultural settings (20, 22–25); however, very few studies have focused on the microbiomes of wild turkeys (11).

In an effort to better characterize the potential effects of the gut microbiota on turkey health and disease, we compared the cecal microbiota from factory-raised domestic, free-ranging domestic, and free-ranging wild turkeys. Sequencing of the V4 region of the 16S rRNA gene was used to determine the abundance of multiple taxa in the ceca of individual birds within each group. Additional experiments were designed to determine the prevalence of bacterial taxa that are common pathogens of commercially raised turkeys. These studies indicate that beta diversity values within the microbiota are significantly different among factory-raised domestic turkeys, free-ranging domestic turkeys, and free-ranging wild turkeys. Several common pathogens associated with commercial poultry production (*Escherichia coli*, *Salmonella* species, and *Staphylococcus* species) were infrequent or absent in the cecal microbiota of free-ranging wild turkeys. *E. coli* strains found in wild turkeys were found to be genetically diverse and carry fewer virulence-associated genes than strains found in factory-raised birds.

## RESULTS

To explore potential differences in the microbiota of wild versus domestic turkeys, a 16S rRNA gene survey of the cecal microbiota was performed. A total of 4,070,891 bacterial reads were obtained, with an average of 53,564 reads per sample and 3,069 amplicon sequence variants (ASVs). We performed principal-coordinate analysis (PCoA) and permutational multivariate analysis of variance (PERMANOVA) of weighted UniFrac distances to compare the microbiota compositions of different flocks of turkeys. At a 13,000-read subsampling depth, PERMANOVA of UniFrac distances revealed significant differences in the microbiota of the samples within provenance and flock (Table 1). The clustering of samples on PCoA ordinations visually depicted these statistical differences, where principal coordinates 1 and 2 separated the samples into three general groups that matched the provenance of the samples when analyzed by both weighted and unweighted UniFrac distances (Fig. 1). Follow-up weighted and unweighted UniFrac analyses confirmed that there were flock-specific effects when each provenance was analyzed separately, except for birds from factory-raised flocks, analyzed by weighted UniFrac (Table 1). The finding that all flocks differed in beta diversity, except those raised in commercial production facilities, is likely a reflection of the highly standardized nature of commercial poultry production.

We also evaluated the variation in the microbiota compositions of the different flocks. At the order level, there were significant differences in the numerical density of the most abundant bacterial taxa (Fig. 2A). For example, in the factory-raised birds, *Clostridiales* were the most abundant taxon ( $71.7\% \pm 3.3\%$ ), much more abundant than in free-ranging domestic turkeys raised on hobby farms ( $33.8\% \pm 1.8\%$ ) or wild turkeys ( $18.3\% \pm 0.7\%$ ). The lower *Clostridiales* read counts in the free-ranging domestic and wild turkey flocks were largely offset by relative increases in *Bacteroidales* and *Coriobacteriales*. The abundances of these reads were all significantly different between

TABLE 1 PERMANOVA for different groups of samples<sup>a</sup>

	df	SS	MS	F	R <sup>2</sup>	p	SS	MS	F	R <sup>2</sup>	p
		Weighted Unifrac					Unweighted Unifrac				
ALL SAMPLES											
Provenance	2	1.68	0.84	55.57	0.55	<0.001	5.12	2.56	44.99	0.56	<0.001
Flock	3	0.52	0.17	11.38	0.17	<0.001	0.82	0.27	4.79	0.09	<0.001
Residuals	55	0.83	0.02		0.27		3.13	0.06		0.35	
Total	60	3.03			1.00		9.06			1.00	
WILD											
Flock	1	0.08	0.08	5.08	0.25	<0.001	0.27	0.27	5.02	0.25	<0.001
Residuals	15	0.22	0.01		0.75		0.81	0.05		0.75	
Total	16	0.30			1.00		1.08			1.00	
DOMESTIC											
Flock	1	0.38	0.38	38.71	0.63	<0.001	0.35	0.35	5.20	0.18	<0.001
Residuals	23	0.23	0.01		0.37		1.53	0.07		0.82	
Total	24	0.61			1.00		1.88			1.00	
FACTORY											
Flock	1	0.06	0.06	2.67	0.14	0.074	0.20	0.20	4.34	0.20	<0.001
Residuals	17	0.38	0.02		0.86		0.79	0.05		0.80	
Total	18	0.44			1.00		0.99			1.00	

<sup>a</sup>df, degree of freedom; SS, sum of squares; MS, mean of squares.

provenances by analysis of composition of microbiomes (ANCOM) (see Table S2 in the supplemental material). However, despite these differences in the abundances of different taxa, the variation in alpha diversity between flocks was not related to the flocks' provenance (Table 2). Therefore, key differences in numerical composition at high taxonomic levels did not necessarily reflect low-level differences in diversity.

To better understand the potential relationship between flock provenance and the carriage of potential pathogens, we next focused on the relative abundances of taxa of veterinary and medical importance by identifying ASVs that best matched known bird pathogens. V4 sequences representing *Staphylococcus* sp. were most prevalent in samples from commercially raised birds. *Staphylococcus* DNA was also detected in one flock of domestic free-ranging turkeys. Detectable levels of *Staphylococcus* DNA were





FIG 1 Principal-component analysis demonstrates that the cecal microbiota of turkeys cluster according to bird provenance. Weighted (A) and unweighted (B) UniFrac distance plots of the different flocks are colored according to the animals' provenance.



**FIG 2** (A) Taxon plot of flocks, grouped by provenance. Order-level assignments above a 2% total relative abundance are shown individually. (B to D) Relative abundances of groups of ASVs (B and C) or an individual ASV (D) from the 16S sequencing data set. Samples are grouped according to flock and colored by provenance. Red, hobby farm (HF); blue, factory farm (FF); green, wild bird (W). (B and C) Sum of reads from multiple ASVs that were each assigned to the *Staphylococcus* or *Campylobacter* genus. (D) Reads from a single ASV that could not be assigned below the *Enterobacteriaceae* family.


not found in any samples from wild birds. Similarly, *Campylobacter* DNA was identified only in factory farm-raised birds. The abundance of *Campylobacter* DNA in some birds was suggestive of heavy colonization; however, it was undetected in other birds within the same flock (Fig. 2B and C).

One limitation of our approach is that without whole-genome data, the short region that we sequenced cannot distinguish known pathogens from similar bacteria with identical sequences across the 16S V4 region. Measurable levels of the family *Enterobacteriaceae* were abundant in samples from both factory-raised flocks and one free-ranging domestic flock (Fig. 2D). The *Enterobacteriaceae* are a large family of bacteria that include *Escherichia coli* as well as other pathogens, including *Salmonella*. The V4 region of the 16S rRNA gene

**TABLE 2** Alpha diversity metrics on a per-flock basis<sup>a</sup>

Flock	Shannon	Pielou's Evenness	Faith's PD	Observed ASVs
Wild flock 1	6.03 +/- 0.09 <sup>ab</sup>	0.753 +/- 0.009 <sup>ab</sup>	17.2 +/- 0.9 <sup>a</sup>	260 +/- 13 <sup>ab</sup>
Wild flock 2	6.43 +/- 0.10 <sup>cd</sup>	0.77 +/- 0.008 <sup>ac</sup>	21.2 +/- 0.7 <sup>b</sup>	330 +/- 18 <sup>c</sup>
Domestic flock 1	6.61 +/- 0.12 <sup>ac</sup>	0.785 +/- 0.010 <sup>a</sup>	21.1 +/- 0.9 <sup>b</sup>	347 +/- 21 <sup>ad</sup>
Domestic flock 2	6.96 +/- 0.03 <sup>ac</sup>	0.81 +/- 0.007 <sup>a</sup>	22.4 +/- 0.8 <sup>a</sup>	402 +/- 20 <sup>ad</sup>
Factory flock 1	6.35 +/- 0.14 <sup>d</sup>	0.76 +/- 0.012 <sup>c</sup>	16.4 +/- 0.7 <sup>b</sup>	330 +/- 15 <sup>cd</sup>
Factory flock 2	5.80 +/- 0.18 <sup>b</sup>	0.72 +/- 0.014 <sup>b</sup>	16 +/- 0.9 <sup>a</sup>	272 +/- 22 <sup>b</sup>

<sup>a</sup>Data are shown as means +/- standard errors of the means (SEM). Different letters next to the SEM represent significant differences between flocks for each metric and were determined by a Kruskal-Wallis test. PD, phylogenetic diversity.




**FIG 3** Cecal contents of factory-raised and free-ranging domestic turkeys contain high levels of *E. coli* DNA compared to those of wild birds. Quantitative PCR was performed for the *ybbW* gene (*E. coli* specific) detected in total genomic DNA isolated from the ceca of factory-raised domestic, free-range domestic, and wild turkeys. Threshold cycle ( $C_T$ ) values from individual birds are shown on the y axis. Average  $C_T$  values of each group are indicated by a horizontal bar.  $C_T$  values of the positive control (100% *E. coli* genomic DNA) and negative control are indicated on the x axis.

does not resolve *E. coli* or *Salmonella* from other *Enterobacteriaceae*, which prevented us from estimating *E. coli* or *Salmonella* abundance in these animals through V4 sequencing alone. As *E. coli* and *Salmonella* are common pathogens in domestic poultry production, we further investigated the prevalence of these potential pathogens in wild and factory-raised turkeys.

The presence of *Salmonella* DNA was detected by PCR targeting the *Salmonella*-specific gene *invA* in total DNA isolated from cecal samples of individual birds. Of 14 samples tested from factory-raised birds, 11 tested positive for *invA*. Conversely, none of the 11 samples collected from wild birds tested positive for the presence of the *invA* gene, suggesting that factory-raised turkeys more frequently contain *Salmonella* in their digestive tracts than wild turkeys. To determine the relative abundance of *E. coli* in cecal samples, a quantitative PCR (qPCR) assay targeting the *E. coli*-specific gene *ybbW* (26) was used. Genomic *E. coli* DNA was clearly present in the total DNA samples obtained from commercially raised turkeys and hobby farm-raised turkeys. Conversely, *E. coli* DNA in samples from wild turkeys was below the limit of detection of the assay (Fig. 3). We also plated cecal samples on MacConkey agar to enrich for the growth of enteric bacteria. Although not detectable by qPCR, we were able to isolate colonies characteristic of *E. coli* from wild turkey cecal samples. Their identity as *E. coli* was subsequently verified by amplification of the *ybbW* gene. *E. coli* was readily cultured from the ceca of factory-raised turkeys. In addition to colony growth consistent with *E. coli* (pink colonies), white colonies were also observed growing on MacConkey agar. These white colonies were not studied further or collected; however, based on growth on MacConkey agar, these colonies were likely *Salmonella* or other enteric bacteria.

As we were able to isolate *E. coli* colonies from the ceca of both wild and factory-raised turkeys, we were interested in further understanding the differences that may exist between these bacterial populations. We therefore performed phylogroup analysis to compare the diversities of *E. coli* lineages that were isolated from factory-raised and wild turkeys. Of *E. coli* strains isolated from wild turkeys, 29 of 30 strains belonged to group A, B1, or E, whereas none belonged to group B2, C, or D (Fig. 4 and Data Set S3). Strains isolated from factory-raised turkeys were more diverse, with all major phylogroups being represented. Several strains (9/50) belonged to cryptic clade I or II, which have been infrequently isolated in other studies. Seven strains were classified as group B2 or D, which are lineages that are commonly associated with extraintestinal pathogenic *E. coli* strains (27, 28). These results suggested that the pathogenic potential of the *E. coli* strains present in wild turkeys may be different from that of the strains present in factory-raised domestic turkeys.

A number of virulence factors have been identified in extraintestinal pathogenic *E.*



**FIG 4** *E. coli* strains isolated from wild and factory-raised turkeys belong to distinct phylogroups. Strains were assigned to phylogroups according to Clermont PCR typing (96). Strains from factory-raised turkeys were more diverse and included those belonging to lineages traditionally thought to include pathogenic strains (lineages B2 and D).

*coli*. These include proteins essential for iron acquisition and for group 2 or group 3 capsule production (29–35). To determine if virulence-associated gene carriage differed between *E. coli* strains found in wild turkeys and those found in factory-raised turkeys, endpoint PCR was used to determine the carriage of three siderophore receptor genes (*iutA*, *iroN*, and *fyuA*) as well as the *kpsMT* genes involved in group 2 or group 3 capsule synthesis (Table 3). Nearly one-half (47%) of the *E. coli* strains isolated from wild turkeys carried the aerobactin receptor gene *iutA*. However, carriage of the salmochelin receptor gene *iroN*, the yersiniabactin receptor gene *fyuA*, or the capsule synthesis genes *kpsMT* was not observed in *E. coli* strains isolated from wild turkeys. Conversely, only 10% of strains isolated from the ceca of commercially produced turkeys contained *iutA*, whereas *iroN* was present in 22%, and *fyuA* was present in 4%. Capsule synthesis genes (*kpsMTII* or *kpsMTIII*) were present in 40% of strains isolated from factory-raised turkeys. The presence of virulence factors was not associated with any particular phylogroup, and several strains carried combinations of virulence factor genes (Table 3 and Data Set S3).

## DISCUSSION

The essential role of the gut microbiota in maintaining animal and human health has been well established (36–39). Although diet is clearly an important selector for many functional guilds of microbes within the gut, host evolutionary history is thought to be a driving factor in determining the prevalence of specific microbial operational taxonomic units (OTUs) (40). Increasingly, evidence supports the theory that many animals coevolved with their microbial symbionts, giving both the host and microbe survival advantages (41–44).

In addition to diet, the intestinal microbiome of domestic farm animals (including poultry) is likely influenced by several factors, such as past and present exposure to

**TABLE 3** Prevalence of virulence factor genes in *E. coli* strains isolated from the ceca of wild and factory-raised turkeys<sup>a</sup>

VF	No. of positive turkeys/total no. of turkeys	
	Wild	Factory raised
No VF	0/32	15/53
<i>iutA</i>	15/32	5/53
<i>iroN</i>	1/32	13/53
<i>fyuA</i>	0/32	7/53
<i>kpsII</i>	0/32	23/53
<i>kpsIII</i>	0/32	4/53
2 VFs	0/32	9/53
3 VFs	0/32	1/53
4 VFs	0/32	1/53

<sup>a</sup>VF, virulence factor.



antibiotics, exposure to the microbiome of the mother, and other microbes in their environment. The data presented here suggest that common production practices (potentially in combination with selective breeding) in modern poultry farming have resulted in a turkey microbiome in which beta diversity decreases from wild birds to free-ranging domestic birds to the highly monotaxic microbiota seen in commercially raised turkeys. The V4 sequencing results from this study are largely consistent with the results of a previous clone-based sequencing approach comparing the microbiota of wild turkeys and domestic turkeys (11).

The principal-component analysis (PCA) results presented in Fig. 1 clearly demonstrate that microbes from factory farm-raised, hobby farm-raised, and wild turkeys separate into distinct groups. In wild birds as well as hobby farm-raised birds, individual data points from different flocks were distributed throughout the groupings. However, our data from the factory farm-raised turkeys suggest that individual flocks may have perceptibly different microbiomes. Samples from factory farm flock 1 cluster separately from those of factory farm flock 2 according to the PCA (Fig. 1B). Distinct differences between factory flocks 1 and 2 are also evident in the taxonomic abundance data (Fig. 2) as well as *Staphylococcus*, *Salmonella*, and *E. coli* levels (Fig. 2 and 3).

As birds in factory flocks 1 and 2 were raised on standard commercial diets under similar housing conditions, we believe that the observed differences between these two flocks are most likely due to age differences between flock 1 (aged ~6 weeks) and flock 2 (aged ~18 weeks). Previous research has clearly demonstrated that the microbiome of developing turkeys undergoes significant changes over time (25, 45, 46). However, subtle differences in the microbiomes of commercial turkeys have also been observed on different farms (22, 45), which may be caused by variable local environmental conditions, including humans or other animals whose microbes may come into contact with the birds.

The composition of the gut microbiota in poultry likely influences a variety of beneficial characteristics, including immune system development and function (47–49). Domestic turkeys raised in commercial turkey production facilities are highly susceptible to a myriad of economically devastating bacterial, fungal, viral, and parasitic diseases (32). Previous research has shown that colonization by some commensal species of microbes prevents/inhibits colonization by pathogenic *Campylobacter*, *Staphylococcus*, and *Salmonella* (50–57) in poultry. We hypothesize that wild relatives of agriculturally important species may carry a heritable microbiome that inhibits colonization by common pathogens. Modern agricultural production practices have largely ignored the potential benefits of this natural microbiota, having instead relied on the widespread use of antibiotics to control pathogens.

While we have not yet established any specific mechanistic links between members of the normal flora and the abundance of specific pathogenic species, we observed that wild turkeys have higher levels of *Coriobacteriales* than hobby farm- or factory-raised domestic turkeys. Some *Coriobacteriales* produce hydroxysteroid dehydrogenase enzymes involved in the conversion of primary to secondary bile acids (58, 59). Bile salt conversion has demonstrated effects on the composition of the microbiome, colonization by intestinal pathogens, and immune responses in humans and livestock (58, 60–63). The growth of *Coriobacteriales* is stimulated by polyphenols found in diverse plants, and these bacteria metabolize them to phenolic compounds that have anti-inflammatory and immunomodulatory effects (59). *Coriobacteriales* are also especially prone to disruption by antibiotic treatment in mice (64). The diets of wild turkeys are free from agricultural antibiotics and likely contain diverse plant polyphenols. Whether members of this family are involved in colonization resistance to *Campylobacter*, *Salmonella*, or *E. coli* should be investigated further.

Suppression of avian-pathogenic *E. coli* in turkeys is an especially important priority for poultry producers. Therefore, it is notable that wild turkeys appeared to contain very little *E. coli* in their ceca. Furthermore, *E. coli* strains isolated from wild turkeys were dissimilar to those isolated from factory-raised birds, in terms of both phylogenetic lineage as well as the presence of specific virulence factors. Many of the *E. coli* strains isolated

from wild turkeys contained the aerobactin receptor gene. Aerobactin is a proven virulence factor in extraintestinal avian infections (33); however, its role in these strains may be related to fitness in the highly competitive environment of the wild turkey intestinal tract. The absence of capsule synthesis genes and salmochelin and yersiniabactin production from strains isolated from wild turkeys may indicate that these strains are not prone to causing bloodstream infections or colonizing other organs. It is possible that bacteriocins, prophages, contact-dependent inhibition, or type 6 secretion systems of *E. coli* lineages established within the ceca of wild turkeys exclude invasion by avian-pathogenic strains frequently found in factory-raised poultry (65–68).

Due to common poultry production practices, microbes colonizing the intestinal tract of commercially raised poultry are minimally, if at all, influenced by the microbiome of the mother. The practice of hatching surface-sterilized eggs in incubators for multiple generations of birds has surely contributed to the loss of any heritable microbial taxa that may have coevolved with the wild turkey over millennia. Consequently, domestic poultry most likely obtain their microbiota almost exclusively from the environment found in the production facilities in which they are raised. Factors influencing the microbiome of these birds are limited to eggshells, litter, feed, and water (69–71). As poultry are often grown in the same production facilities generation after generation, microbial contamination from previous generations of birds is likely a significant factor influencing the microbiome of birds raised in high-density factory farms. The taxa and gene carriage of microbes that seed the microbiome of factory farm-raised birds have also been influenced by production practices such as feeding growth-enhancing antibiotics and feeds containing metals such as copper and zinc (72, 73). These have likely skewed the gut microbiota of poultry raised in large-scale production facilities toward taxa most capable of survival in modern turkey production facilities rather than microbiota contributing to the mutual survival of the host and microbe.

The transfer of microbiota from mother to infant has been best characterized in mammals. The transfer of maternal microbes to the mammalian young begins during the birthing process and continues through nursing and social interactions (74, 75). Coprophagy is common in many animal species, including turkeys (76–78), and it is common to see turkeys consuming cecal drops of their cage mates. This innate behavior in turkeys may have evolved to enable the bird-to-bird spread of beneficial microbiota within a flock. Recent work documents that the newly hatched young of some birds readily consume cecal drops, but not normal rectal feces, of their mothers. This consumption of maternal cecal drops by chicks was observed only during a short window of time (approximately the first month of life) (79). This behavior potentially facilitates the establishment of a beneficial, heritable gut microbiome from mother to chick.

The gut microbiome is perhaps one of the most complex biological communities. As in the analysis of any biological community, it is essential to consider the effects of dominant taxa as well as taxa that may have a relatively small but potentially important role in the community. The goal of this study was to identify potential changes/differences in the microbial composition of factory-raised turkeys compared to that of their wild predecessors. The results presented here demonstrate that the overall abundance of *E. coli*, *Salmonella*, *Campylobacter*, and *Staphylococcus* in wild turkeys is much lower than the levels commonly found in commercially raised turkeys. Furthermore, *E. coli* strains occupying the intestinal tract of wild turkeys appear distinct in both lineage and carriage of common virulence factors compared to strains commonly found in commercially raised turkeys. The strong correlation between bird provenance, increased microbial diversity, and low pathogen carriage warrants further research into the potential for mining the microbiome of free-ranging wild turkeys (as well as wild relatives of other agriculturally important species) in search of therapeutics or probiotics for use in controlling pathogens common in agricultural food production.

## MATERIALS AND METHODS

**Definition of turkey groups used in this study.** The term “wild turkey” can mean both a strain of turkey as well as the lack of domestication. In this study, we define wild turkey as a population of self-



sustaining, wild, free-ranging birds. All wild turkeys sampled in this work were of the Rio Grande subspecies (*Meleagris gallopavo intermedia*) that have ranged freely for generations in the mountains of North Central Utah in the United States. Birds described as “free-range domestic turkeys” in this study are domesticated turkeys ranging freely outdoors. All free-range domestic turkeys in this study were from hobby farms where they were allowed to forage freely outdoors in both the summer and winter. The diet of all domestic free-range turkeys was supplemented with commercial poultry food by their owners. The term “factory-raised domestic turkey” refers to turkeys raised in commercial turkey production facilities. All turkeys in this group were of the Broad Breasted White variety. Although these factory-raised birds may fit the legal definition of “free-range” by virtue of their caging conditions, they were not considered free-ranging for the purposes of this study.

**Collection of cecal samples.** Some birds, including turkeys and chickens, produce two distinctly different types of feces. Cecal drops are a type of feces that the bird periodically excretes directly from the intestinal cecum (80). Previous work has demonstrated that the cecum contains the greatest microbial diversity found in the intestinal tract of poultry (81, 82). Additionally, the microbiota found in cecal drops is highly reflective of the microbiota found in cecal contents collected following sacrifice of the bird (83). The collection of cecal drops, which are easily distinguishable from normal feces, enables a simple, non-invasive method of obtaining a clear view of the cecal microbiota and eliminates the need to sacrifice (or even to come into contact with) study animals.

In this study, all samples were of cecal origin. Cecal drops from wild and free-ranging domestic turkeys were collected during the winter months following snowstorms. Sample collection immediately following snowstorms ensured that only fresh samples were collected and that the sample remained relatively uncontaminated by bacteria from the soil or other environmental sources. Cecal contents from one flock of factory-raised turkeys were collected from a turkey processing facility postmortem. Cecal drops from a second commercially raised flock were collected from the floor of the production facility. Sampling sites, bird age, and other details of sample origin are listed in Fig. S1 in the supplemental material.

**DNA preparation.** Following sample collection, all cecal contents were kept frozen until DNA isolation. DNA used for V4 sequencing was extracted from each sample using the Zymo Quick-DNA fecal/soil microbe 96 kit (catalog number D6011) according to the manufacturer’s instructions, including a bead homogenization step using a 2010 Geno/Grinder (Spex, Metuchen, NJ) at 1,750 rpm for 10 min. DNA was prepared for 16S rRNA gene V4 region sequencing based on an established protocol, with minor deviations (84). First, the V4 region of the 16S rRNA gene was amplified individually from each sample with the AccuPrime Pfx enzyme (Thermo Fisher Scientific, Waltham, MA, USA) in 20- $\mu$ L volumes using a subset of the exact primer sequences described previously (84). PCR amplicons were normalized using the SequalPrep normalization kit (Applied Biosystems, Waltham, MA, USA) and pooled into groups of 96 reaction mixtures, and fragments in the range of 250 to 450 bp were purified using a BluePippin (Sage Science, Beverly, MA) selection step. Equimolar normalization of each pool and sequencing were performed at the BYU DNA Sequencing center on a partial 2-by-250 lane (v2) of a HiSeq 2500 platform (Illumina, Inc., San Diego, CA). Laser complexity was ensured by including at least 10% of each lane with shotgun sequencing libraries for other bacterial genomes.

**Sequence analysis.** Sample reads were demultiplexed on the Illumina platform and analyzed using QIIME2 (85, 86) and R. Briefly, reads were trimmed to maximize the quality scores of each nucleotide position. DADA2 (87) was used to denoise, dereplicate, and call amplicon sequence variants (ASVs), and taxonomy was assigned to the ASVs using GreenGenes classifier 13\_8\_99 (88). ASV tables were filtered to 13,000 reads per sample, and differences between groups were determined by PERMANOVA (89) of weighted and unweighted UniFrac distances (90, 91). To permit calculating UniFrac distances, we built a phylogenetic tree with fasttree2 (92) based on mafft alignment (93). Differences in OTU abundances between samples were determined using ANCOM (94). Abundances of individual OTUs were manually analyzed based on the taxonomic assignments, which were assigned to OTUs using QIIME2 q2-feature-classifier (95). Alpha diversity metrics were defined using QIIME2, and differences in alpha diversity metrics between sampling locations were determined by a Kruskal-Wallis test.

**Determination of relative *E. coli* DNA levels in cecal samples.** A qPCR-based assay was designed to estimate the relative abundance of *E. coli* DNA in each cecal sample based on the detection of the *ybbW* gene, which is found exclusively in *E. coli* (26). For qPCR experiments, total DNA was isolated from cecal samples using the Qiagen blood and tissue DNA kit as directed by the manufacturer. DNA samples were diluted to a concentration of 100 ng/ $\mu$ L. IDT PrimeTime gene expression master mix was used in all qPCR assays. Thermocycling was performed as suggested by the manufacturer (40 cycles of 95°C of denaturation for 15 s followed by 57°C of annealing/amplification for 1 min) using an ABI StepOnePlus real-time PCR system. The qPCR primers and probe were designed using IDT PrimerQuest and manufactured by Integrated DNA Technologies. Primer and probe sets used in this study are listed in Table 4. The efficiency and reproducibility of amplification were verified by generating a standard curve using doubling dilutions of positive-control DNA. Negative controls consisted of reaction mixtures with DNA elution buffer rather than DNA. Each sample was tested in duplicate. Purified DNA from pooled *E. coli* strains was used as a positive control.

**Isolation and genotyping of *E. coli* strains.** *E. coli* strains present in cecal samples were isolated by homogenizing a portion of the sample in sterile phosphate-buffered saline (PBS) and plating the sample onto MacConkey agar, followed by growth at 37°C for 24 h. Colonies with characteristic *E. coli* morphology were then restreaked and verified as *E. coli* by PCR targeting the *ybbW* gene. Total DNA was isolated from individual colonies using a minigenomic DNA kit for blood and cultured cells (IBI Scientific). Putative *E. coli* strains were assigned to phylogroups using the Clermont quadruplex assay, with additional PCR tests to distinguish group C or group E when warranted, as previously described (96).

**TABLE 4** Primers used in this study

Primer or probe	Sequence	Target, expected size (bp)
invA 1F invA 1R	GTGAAATTATCGCCACGTTTCGGGCAA TCATCGCACCGTCAAAGGAACC	<i>invA</i> , 284
iutA F iutA R	CTGCAGTACTCCGATCGGCTG TGGTTGGAGGTAAAGCGCTCATG	<i>iutA</i> , 470
iroN R iroN F	TGTCGGTACAGGCGGTTTCGTC CTCTGGTGGTGAAGCCACC	<i>iroN</i> , 814
fyuA F fyuA R	ACGGCTTTATCCTCTGGCCTTGG TGAAAACCCAGTCATCGGTGG	<i>fyuA</i> , 877
kpsII F kpsII R	GCGCATTTGCTGATACTGTTG AGGTAGTTCAGACTCACACCT	<i>kpsMTII</i> , 581
kpsIII F kpsIII R	TCCTCTTGCTACTATCCCCCT AAGGCGTATCCATCCCTCCTAAC	<i>kpsMIII</i> , 390
chuA.1b chuA.2	ATGGTACCGGACGAACCAAC TGCCGCCAGTACCAAAGACA	<i>chuA</i> , 288
AceK.f ArpA1.r	AACGCTATTCGCCAGCTTGC TCTCCCCATACCGTACGCTA	<i>arpA</i> , 400
yjaA.1b yjaA.2b	CAAACGTGAAGTGTACAGGAG AATGCGTTCTCAACCTGTG	<i>yjaA</i> , 211
TspE4C2.1b TspE4C2.2b	CACTATTCGTAAGGTCATCC AGTTTATCGCTGCGGGTCGC	<i>TspE4.C2</i> , 152
ArpAgpE.f ArpAgpE.r	GATTCCATCTTGTCAAAATATGCC GAAAAGAAAAGAATTCCCAAGAG	Group E <i>arpA</i> , 301
trpAgpC.1 trpAgpC.2	AGTTTTATGCCAGTGCAGG TCTGCGCCGGTCACGCC	Group C <i>trpA</i> , 219
ybbW 1F ybbW 1R	TGATTGGCAAATCTGGCCG CGTTGACCAGCCAGAAGATTAAG	qPCR of <i>ybbW</i>
ybbW Probe	56-FAM/AAGCCCGGT/ZEN/AGAGAAAGGCCTAAC/3IABkFQ <sup>a</sup>	Probe for qPCR of <i>ybbW</i>

<sup>a</sup>FAM, carboxyfluorescein; 3IABkFQ, 3' Iowa Black FQ.

The presence or absence of genes associated with the virulence of avian-pathogenic *E. coli* (*iutA*, *iss*, *iroN*, *fyuA*, *kpsMTII*, and *kpsMIII*) was determined by PCR. These reactions were performed using approximately 100 ng genomic DNA as the template and 20 pmol of each primer in OneTaq master mix. The conditions were 94°C for 3 min followed by 30 cycles of 94°C for 15 s, 57°C for 15 s, and 68°C for 45 s and a final extension step at 68°C for 5 min. The *fyuA*, *kpsII*, and *kpsIII* reactions were multiplexed, while the *iutA*, *iroN*, and *iutA* reactions were run individually. Primers used for these reactions are listed in Table 4.

**Detection of *Salmonella* DNA in cecal samples.** The presence or absence of *Salmonella* sp. in cecal sample DNA was determined using a semiquantitative PCR assay using primers targeting the *invA* gene (Table 4), which has previously been demonstrated to specifically detect most *Salmonella* strains (97). The conditions used for these PCR tests were identical to the ones for the *E. coli* virulence genotyping described above, except that the annealing temperature was 56°C.

**Data availability.** Sequences were deposited to the National Center for Biotechnology Information Sequence Read Archive under BioProject accession number [PRJNA786944](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA786944).

## SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**SUPPLEMENTAL FILE 1**, XLSX file, 0.01 MB.

**SUPPLEMENTAL FILE 2**, XLSX file, 0.01 MB.

**SUPPLEMENTAL FILE 3**, XLSX file, 0.01 MB.

## REFERENCES

- Speller CF, Kemp BM, Wyatt SD, Monroe C, Lipe WD, Arndt UM, Yang DY. 2010. Ancient mitochondrial DNA analysis reveals complexity of indigenous North American turkey domestication. *Proc Natl Acad Sci U S A* 107: 2807–2812. <https://doi.org/10.1073/pnas.0909724107>.
- Guan X, Silva P, Gyenai KB, Xu J, Geng T, Tu Z, Samuels DC, Smith EJ. 2009. The mitochondrial genome sequence and molecular phylogeny of the turkey, *Meleagris gallopavo*. *Anim Genet* 40:134–141. <https://doi.org/10.1111/j.1365-2052.2008.01810.x>.
- Thornton EK, Emery KF, Steadman DW, Speller C, Matheny R, Yang D. 2012. Earliest Mexican turkeys (*Meleagris gallopavo*) in the Maya region: implications for pre-Hispanic animal trade and the timing of turkey domestication. *PLoS One* 7:e42630. <https://doi.org/10.1371/journal.pone.0042630>.
- Aslam ML, Bastiaansen JWM, Elferink MG, Megens H-J, Crooijmans RPMA, Blomberg LA, Fleischer RC, Van Tassell CP, Sonstegard TS, Schroeder SG, Groenen MAM, Long JA. 2012. Whole genome SNP discovery and analysis of genetic diversity in turkey (*Meleagris gallopavo*). *BMC Genomics* 13: 391. <https://doi.org/10.1186/1471-2164-13-391>.
- Canales Vergara AM, Landi V, Delgado Bermejo JV, Martinez A, Cervantes Acosta P, Pons Barro A, Bigi D, Sponenberg P, Helal M, Hossein Banabazi M, Camacho Vallejo ME. 2019. Tracing worldwide turkey genetic diversity using D-loop sequence mitochondrial DNA analysis. *Animals (Basel)* 9: 897. <https://doi.org/10.3390/ani9110897>.
- Stover KK, Weinreich DM, Roberts TJ, Brainerd EL. 2018. Patterns of musculoskeletal growth and dimensional changes associated with selection and developmental plasticity in domestic and wild strain turkeys. *Ecol Evol* 8:3229–3239. <https://doi.org/10.1002/ece3.3881>.
- USDA. 2021. Turkey sector: background & statistics. USDA, Washington, DC.
- Colston TJ, Jackson CR. 2016. Microbiome evolution along divergent branches of the vertebrate tree of life: what is known and unknown. *Mol Ecol* 25:3776–3800. <https://doi.org/10.1111/mec.13730>.
- Clayton JB, Vangay P, Huang H, Ward T, Hillmann BM, Al-Ghalith GA, Travis DA, Long HT, Tuan BV, Minh VV, Cabana F, Nadler T, Toddes B, Murphy T, Glander KE, Johnson TJ, Knights D. 2016. Captivity humanizes the primate microbiome. *Proc Natl Acad Sci U S A* 113:10376–10381. <https://doi.org/10.1073/pnas.1521835113>.
- McKenzie VJ, Song SJ, Delsuc F, Prest TL, Oliverio AM, Korpita TM, Alexiev A, Amato KR, Metcalf JL, Kowalewski M, Avenant NL, Link A, Di Fiore A, Seguin-Orlando A, Feh C, Orlando L, Mendelson JR, Sanders J, Knight R. 2017. The effects of captivity on the mammalian gut microbiome. *Integr Comp Biol* 57:690–704. <https://doi.org/10.1093/icb/ixc090>.
- Scupham AJ, Patton TG, Bent E, Bayles DO. 2008. Comparison of the cecal microbiota of domestic and wild turkeys. *Microb Ecol* 56:322–331. <https://doi.org/10.1007/s00248-007-9349-4>.
- Schreuder J, Velkers FC, Bouwstra RJ, Beerens N, Stegeman JA, de Boer WF, van Hooft P, Elbers ARW, Bossers A, Jurburg SD. 2020. An observational field study of the cloacal microbiota in adult laying hens with and without access to an outdoor range. *Anim Microbiome* 2:28. <https://doi.org/10.1186/s42523-020-00044-6>.
- Cui Y, Wang Q, Liu S, Sun R, Zhou Y, Li Y. 2017. Age-related variations in intestinal microflora of free-range and caged hens. *Front Microbiol* 8: 1310. <https://doi.org/10.3389/fmicb.2017.01310>.
- Hubert SM, Al-Ajeeli M, Bailey CA, Athrey G. 2019. The role of housing environment and dietary protein source on the gut microbiota of chicken. *Animals (Basel)* 9:1085. <https://doi.org/10.3390/ani9121085>.
- Souza MR, Moreira JL, Barbosa FH, Cerqueira MM, Nunes AC, Nicoli JR. 2007. Influence of intensive and extensive breeding on lactic acid bacteria isolated from *Gallus gallus domesticus* ceca. *Vet Microbiol* 120: 142–150. <https://doi.org/10.1016/j.vetmic.2006.10.019>.
- Pascoe EL, Hauffe HC, Marchesi JR, Perkins SE. 2017. Network analysis of gut microbiota literature: an overview of the research landscape in non-human animal studies. *ISME J* 11:2644–2651. <https://doi.org/10.1038/ismej.2017.133>.
- Hird SM. 2017. Evolutionary biology needs wild microbiomes. *Front Microbiol* 8:725. <https://doi.org/10.3389/fmicb.2017.00725>.
- Brislin JT, Gong J, Sharif S. 2008. Interactions between commensal bacteria and the gut-associated immune system of the chicken. *Anim Health Res Rev* 9:101–110. <https://doi.org/10.1017/S146625230800145X>.
- Yeoman CJ, Chia N, Jeraldo P, Sipos M, Goldenfeld ND, White BA. 2012. The microbiome of the chicken gastrointestinal tract. *Anim Health Res Rev* 13:89–99. <https://doi.org/10.1017/S1466252312000138>.
- Danzeisen JL, Calvert AJ, Noll SL, McComb B, Sherwood JS, Logue CM, Johnson TJ. 2013. Succession of the turkey gastrointestinal bacterial microbiome related to weight gain. *PeerJ* 1:e237. <https://doi.org/10.7717/peerj.237>.
- Wei S, Gutek A, Lilburn M, Yu Z. 2013. Abundance of pathogens in the gut and litter of broiler chickens as affected by bacitracin and litter management. *Vet Microbiol* 166:595–601. <https://doi.org/10.1016/j.vetmic.2013.06.006>.
- Taylor KJM, Ngunjiri JM, Abundo MC, Jang H, Elash M, Ghorbani A, KC M, Weber BP, Johnson TJ, Lee C-W. 2020. Respiratory and gut microbiota in commercial turkey flocks with disparate weight gain trajectories display differential compositional dynamics. *Appl Environ Microbiol* 86:e00431–20. <https://doi.org/10.1128/AEM.00431-20>.
- Danzeisen JL, Clayton JB, Huang H, Knights D, McComb B, Hayer SS, Johnson TJ. 2015. Temporal relationships exist between cecum, ileum, and litter bacterial microbiomes in a commercial turkey flock, and subtherapeutic penicillin treatment impacts ileum bacterial community establishment. *Front Vet Sci* 2:56. <https://doi.org/10.3389/fvets.2015.00056>.
- Johnson TA, Sylte MJ, Looft T. 2019. In-feed bacitracin methylene disalicylate modulates the turkey microbiota and metabolome in a dose-dependent manner. *Sci Rep* 9:8212. <https://doi.org/10.1038/s41598-019-44338-5>.
- Wilkinson TJ, Cowan AA, Vallin HE, Onime LA, Oyama LB, Cameron SJ, Gonot C, Moorby JM, Waddams K, Theobald VJ, Leemans D, Bowra S, Nixey C, Huws SA. 2017. Characterization of the microbiome along the gastrointestinal tract of growing turkeys. *Front Microbiol* 8:1089. <https://doi.org/10.3389/fmicb.2017.01089>.
- Walker DI, McQuillan J, Taiwo M, Parks R, Stenton CA, Morgan H, Mowlem MC, Lees DN. 2017. A highly specific *Escherichia coli* qPCR and its comparison with existing methods for environmental waters. *Water Res* 126: 101–110. <https://doi.org/10.1016/j.watres.2017.08.032>.
- Picard B, Garcia JS, Gouriou S, Duriez P, Brahimi N, Bingen E, Elion J, Denamur E. 1999. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect Immun* 67:546–553. <https://doi.org/10.1128/IAI.67.2.546-553.1999>.
- Johnson JR, Stell AL. 2000. Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J Infect Dis* 181:261–272. <https://doi.org/10.1086/315217>.
- Dziva F, Stevens MP. 2008. Colibacillosis in poultry: unravelling the molecular basis of virulence of avian pathogenic *Escherichia coli* in their natural hosts. *Avian Pathol* 37:355–366. <https://doi.org/10.1080/03079450802216652>.
- Magieiros L, Meric G, Bayliss SC, Pensar J, Pascoe B, Mourkas E, Calland JK, Yahara K, Murray S, Wilkinson TS, Williams LK, Hitchings MD, Porter J, Kemmett K, Feil EJ, Jolley KA, Williams NJ, Corander J, Sheppard SK. 2021. Genome evolution and the emergence of pathogenicity in avian *Escherichia coli*. *Nat Commun* 12:765. <https://doi.org/10.1038/s41467-021-20988-w>.
- La Ragione RM, Woodward MJ. 2002. Virulence factors of *Escherichia coli* serotypes associated with avian colisepticaemia. *Res Vet Sci* 73:27–35. [https://doi.org/10.1016/S0034-5288\(02\)00075-9](https://doi.org/10.1016/S0034-5288(02)00075-9).
- Swayne DE, Boulianne M, Logue CM, McDougald LR, Nair V, Suarez DL (ed). 2020. Diseases of poultry, 14th ed. Wiley-Blackwell, Hoboken, NJ.
- Dozois CM, Daigle F, Curtiss R, III. 2003. Identification of pathogen-specific and conserved genes expressed in vivo by an avian pathogenic *Escherichia coli* strain. *Proc Natl Acad Sci U S A* 100:247–252. <https://doi.org/10.1073/pnas.232686799>.
- Collingwood C, Kemmett K, Williams N, Wigley P. 2014. Is the concept of avian pathogenic *Escherichia coli* as a single pathotype fundamentally flawed? *Front Vet Sci* 1:5. <https://doi.org/10.3389/fvets.2014.00005>.
- Antao EM, Glodde S, Li G, Sharifi R, Homeier T, Laturnus C, Diehl I, Bethe A, Philipp HC, Preisinger R, Wieler LH, Ewers C. 2008. The chicken as a natural model for extraintestinal infections caused by avian pathogenic *Escherichia coli* (APEC). *Microb Pathog* 45:361–369. <https://doi.org/10.1016/j.micpath.2008.08.005>.
- Clemente JC, Ursell LK, Parfrey LW, Knight R. 2012. The impact of the gut microbiota on human health: an integrative view. *Cell* 148:1258–1270. <https://doi.org/10.1016/j.cell.2012.01.035>.
- Sekirov I, Russell SL, Antunes LC, Finlay BB. 2010. Gut microbiota in health and disease. *Physiol Rev* 90:859–904. <https://doi.org/10.1152/physrev.00045.2009>.
- Peixoto RS, Harkins DM, Nelson KE. 2021. Advances in microbiome research for animal health. *Annu Rev Anim Biosci* 9:289–311. <https://doi.org/10.1146/annurev-animal-091020-075907>.
- Waite DW, Taylor MW. 2014. Characterizing the avian gut microbiota: membership, driving influences, and potential function. *Front Microbiol* 5:223. <https://doi.org/10.3389/fmicb.2014.00223>.

40. Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, Ley RE, Farnleitner AH. 2019. Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. *Nat Commun* 10:2200. <https://doi.org/10.1038/s41467-019-10191-3>.
41. Moeller AH, Li Y, Mpoudi Ngole E, Ahuka-Mundede S, Lonsdorf EV, Pusey AE, Peeters M, Hahn BH, Ochman H. 2014. Rapid changes in the gut microbiome during human evolution. *Proc Natl Acad Sci U S A* 111: 16431–16435. <https://doi.org/10.1073/pnas.1419136111>.
42. Moeller AH, Caro-Quintero A, Mjunga D, Georgiev AV, Lonsdorf EV, Muller MN, Pusey AE, Peeters M, Hahn BH, Ochman H. 2016. Cospeciation of gut microbiota with hominids. *Science* 353:380–382. <https://doi.org/10.1126/science.aaf3951>.
43. Groussin M, Mazel F, Alm EJ. 2020. Co-evolution and co-speciation of host-gut bacteria systems. *Cell Host Microbe* 28:12–22. <https://doi.org/10.1016/j.chom.2020.06.013>.
44. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, Schlegel ML, Tucker TA, Schrenzel MD, Knight R, Gordon JL. 2008. Evolution of mammals and their gut microbes. *Science* 320:1647–1651. <https://doi.org/10.1126/science.1155725>.
45. Adhikari B, Tellez-Isaías G, Jiang T, Wooming B, Kwon YM. 2020. Comprehensive survey of the litter bacterial communities in commercial turkey farms. *Front Vet Sci* 7:596933. <https://doi.org/10.3389/fvets.2020.596933>.
46. Scupham AJ. 2007. Succession in the intestinal microbiota of preadolescent turkeys. *FEMS Microbiol Ecol* 60:136–147. <https://doi.org/10.1111/j.1574-6941.2006.00245.x>.
47. Mwangi WN, Beal RK, Powers C, Wu X, Humphrey T, Watson M, Bailey M, Friedman A, Smith AL. 2010. Regional and global changes in TCR $\alpha$ -beta T cell repertoires in the gut are dependent upon the complexity of the enteric microflora. *Dev Comp Immunol* 34:406–417. <https://doi.org/10.1016/j.dci.2009.11.009>.
48. van Veelen HPJ, Falcao Salles J, Matson KD, van der Velde M, Tieleman BL. 2020. Microbial environment shapes immune function and cloacal microbiota dynamics in zebra finches *Taeniopygia guttata*. *Anim Microbiome* 2:21. <https://doi.org/10.1186/s42523-020-00039-3>.
49. McFall-Ngai M. 2007. Adaptive immunity: care for the community. *Nature* 445:153. <https://doi.org/10.1038/445153a>.
50. Cole K, Farnell MB, Donoghue AM, Stern NJ, Svetoch EA, Eruslanov BN, Volodina LI, Kovalev YN, Perelygin VV, Mitsevich EV, Mitsevich IP, Levchuk VP, Pokhilenko VD, Borzenkov VN, Svetoch OE, Kudryavtseva TY, Reyes-Herrera I, Blore PJ, Solis de los Santos F, Donoghue DJ. 2006. Bacteriocins reduce *Campylobacter* colonization and alter gut morphology in turkey poults. *Poult Sci* 85:1570–1575. <https://doi.org/10.1093/ps/85.9.1570>.
51. Stern NJ, Svetoch EA, Eruslanov BV, Kovalev YN, Volodina LI, Perelygin VV, Mitsevich EV, Mitsevich IP, Levchuk VP. 2005. *Paenibacillus polymyxa* purified bacteriocin to control *Campylobacter jejuni* in chickens. *J Food Prot* 68:1450–1453. <https://doi.org/10.4315/0362-028X-68.7.1450>.
52. Stern NJ, Svetoch EA, Eruslanov BV, Perelygin VV, Mitsevich EV, Mitsevich IP, Pokhilenko VD, Levchuk VP, Svetoch OE, Seal BS. 2006. Isolation of a *Lactobacillus salivarius* strain and purification of its bacteriocin, which is inhibitory to *Campylobacter jejuni* in the chicken gastrointestinal system. *Antimicrob Agents Chemother* 50:3111–3116. <https://doi.org/10.1128/AAC.00259-06>.
53. Scupham AJ, Jones JA, Rettedal EA, Weber TE. 2010. Antibiotic manipulation of the intestinal microbiota to identify microbes associated with *Campylobacter jejuni* exclusion in poultry. *Appl Environ Microbiol* 76:8026–8032. <https://doi.org/10.1128/AEM.00678-10>.
54. Scupham AJ. 2009. *Campylobacter* colonization of the turkey intestine in the context of microbial community development. *Appl Environ Microbiol* 75:3564–3571. <https://doi.org/10.1128/AEM.01409-08>.
55. Nicoll TR, Jensen MM. 1987. Staphylococcosis of turkeys. 5. Large-scale control programs using bacterial interference. *Avian Dis* 31:85–88. <https://doi.org/10.2307/1590778>.
56. Wilkinson DM, Jensen MM. 1987. Staphylococcosis of turkeys. 4. Characterization of a bacteriocin produced by an interfering *Staphylococcus*. *Avian Dis* 31:80–84. <https://doi.org/10.2307/1590777>.
57. Bielke LR, Elwood AL, Donoghue DJ, Donoghue AM, Newberry LA, Neighbor NK, Hargis BM. 2003. Approach for selection of individual enteric bacteria for competitive exclusion in turkey poults. *Poult Sci* 82: 1378–1382. <https://doi.org/10.1093/ps/82.9.1378>.
58. Ridlon JM, Kang DJ, Hylemon PB. 2006. Bile salt biotransformations by human intestinal bacteria. *J Lipid Res* 47:241–259. <https://doi.org/10.1194/jlr.R500013-JLR200>.
59. Rodriguez-Daza MC, Roquim M, Dudonne S, Pilon G, Levy E, Marette A, Roy D, Desjardins Y. 2020. Berry polyphenols and fibers modulate distinct microbial metabolic functions and gut microbiota enterotype-like clustering in obese mice. *Front Microbiol* 11:2032. <https://doi.org/10.3389/fmicb.2020.02032>.
60. Gaboriaud P, Sadrin G, Guitton E, Fort G, Niepceon A, Lallier N, Rossignol C, Larcher T, Sausset A, Guabiraba R, Silvestre A, Lacroix-Lamande S, Schouler C, Laurent F, Bussiere FI. 2020. The absence of gut microbiota alters the development of the apicomplexan parasite *Eimeria tenella*. *Front Cell Infect Microbiol* 10:632556. <https://doi.org/10.3389/fcimb.2020.632556>.
61. McKenney PT, Yan J, Vaubourgeix J, Becattini S, Lampen N, Motzer A, Larson PJ, Dannaoui D, Fujisawa S, Xavier JB, Pamer EG. 2019. Intestinal bile acids induce a morphotype switch in vancomycin-resistant *Enterococcus* that facilitates intestinal colonization. *Cell Host Microbe* 25:695–705.e5. <https://doi.org/10.1016/j.chom.2019.03.008>.
62. Zhang L, Wu W, Lee YK, Xie J, Zhang H. 2018. Spatial heterogeneity and co-occurrence of mucosal and luminal microbiome across swine intestinal tract. *Front Microbiol* 9:48. <https://doi.org/10.3389/fmicb.2018.00048>.
63. van Best N, Rolle-Kampczyk U, Schaap FG, Basic M, Olde Damink SWM, Bleich A, Savelkoul PHM, von Bergen M, Penders J, Hornef MW. 2020. Bile acids drive the newborn's gut microbiota maturation. *Nat Commun* 11: 3692. <https://doi.org/10.1038/s41467-020-17183-8>.
64. Zákostelská Z, Málková J, Klimešová K, Rossmann P, Hornová M, Novosádová I, Stehlíková Z, Kostovčík M, Hudcovic T, Štěpánková R, Jůzlová K, Hercogová J, Tlaskalová-Hogenová H, Kverka M. 2016. Intestinal microbiota promotes psoriasis-like skin inflammation by enhancing Th17 response. *PLoS One* 11: e0159539. <https://doi.org/10.1371/journal.pone.0159539>.
65. Aoki SK, Pamma R, Hernday AD, Bickham JE, Braaten BA, Low DA. 2005. Contact-dependent inhibition of growth in *Escherichia coli*. *Science* 309: 1245–1248. <https://doi.org/10.1126/science.1115109>.
66. Journet L, Cascales E. 2016. The type VI secretion system in *Escherichia coli* and related species. *EcoSal Plus* 7:ESP-0009-2015. <https://doi.org/10.1128/ecosalplus.ESP-0009-2015>.
67. Cascales E, Buchanan SK, Duche D, Kleantous C, Lloubes R, Postle K, Riley M, Slatin S, Cavard D. 2007. Colicin biology. *Microbiol Mol Biol Rev* 71:158–229. <https://doi.org/10.1128/MMBR.00036-06>.
68. Da Re S, Valle J, Charbonnel N, Beloin C, Latour-Lambert P, Faure P, Turlin E, Le Bouguenec C, Renaud-Mongenie G, Forestier C, Ghigo JM. 2013. Identification of commensal *Escherichia coli* genes involved in biofilm resistance to pathogen colonization. *PLoS One* 8:e61628. <https://doi.org/10.1371/journal.pone.0061628>.
69. Stanley D, Geier MS, Hughes RJ, Denman SE, Moore RJ. 2013. Highly variable microbiota development in the chicken gastrointestinal tract. *PLoS One* 8:e84290. <https://doi.org/10.1371/journal.pone.0084290>.
70. Maki JJ, Bobeck EA, Sylte MJ, Looft T. 2020. Eggshell and environmental bacteria contribute to the intestinal microbiota of growing chickens. *J Anim Sci Biotechnol* 11:60. <https://doi.org/10.1186/s40104-020-00459-w>.
71. Maki JJ, Klima CL, Sylte MJ, Looft T. 2019. The microbial pecking order: utilization of intestinal microbiota for poultry health. *Microorganisms* 7:376. <https://doi.org/10.3390/microorganisms7100376>.
72. Olukosi OA, van Kuijck S, Han Y. 2018. Copper and zinc sources and levels of zinc inclusion influence growth performance, tissue trace mineral content, and carcass yield of broiler chickens. *Poult Sci* 97:3891–3898. <https://doi.org/10.3382/ps/pey247>.
73. Argudin MA, Deplano A, Meghraoui A, Dodemont M, Heinrichs A, Denis O, Nonhoff C, Roisin S. 2017. Bacteria from animals as a pool of antimicrobial resistance genes. *Antibiotics (Basel)* 6:12. <https://doi.org/10.3390/antibiotics6020012>.
74. Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer N, Knight R. 2010. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci U S A* 107:11971–11975. <https://doi.org/10.1073/pnas.1002601107>.
75. Mueller NT, Bakacs E, Combellick J, Grigoryan Z, Dominguez-Bello MG. 2015. The infant microbiome development: mom matters. *Trends Mol Med* 21:109–117. <https://doi.org/10.1016/j.molmed.2014.12.002>.
76. Soave O, Brand CD. 1991. Coprophagy in animals: a review. *Cornell Vet* 81:357–364.
77. Pan D, Yu Z. 2014. Intestinal microbiome of poultry and its interaction with host and diet. *Gut Microbes* 5:108–119. <https://doi.org/10.4161/gmic.26945>.
78. Kers JG, Velkers FC, Fischer EAJ, Hermes GDA, Stegeman JA, Smidt H. 2018. Host and environmental factors affecting the intestinal microbiota in chickens. *Front Microbiol* 9:235. <https://doi.org/10.3389/fmicb.2018.00235>.



79. Kobayashi A, Tsuchida S, Ueda A, Yamada T, Murata K, Nakamura H, Ushida K. 2019. Role of coprophagy in the cecal microbiome development of an herbivorous bird Japanese rock ptarmigan. *J Vet Med Sci* 81: 1389–1399. <https://doi.org/10.1292/jvms.19-0014>.
80. Fenna L, Boag DA. 1974. Filling and emptying of the galliform caecum. *Can J Zool* 52:537–540. <https://doi.org/10.1139/z74-067>.
81. Barnes EM, Mead GC, Barnum DA, Harry EG. 1972. The intestinal flora of the chicken in the period 2 to 6 weeks of age, with particular reference to the anaerobic bacteria. *Br Poult Sci* 13:311–326. <https://doi.org/10.1080/00071667208415953>.
82. Bjerrum L, Engberg RM, Leser TD, Jensen BB, Finster K, Pedersen K. 2006. Microbial community composition of the ileum and cecum of broiler chickens as revealed by molecular and culture-based techniques. *Poult Sci* 85:1151–1164. <https://doi.org/10.1093/ps/85.7.1151>.
83. Pauwels J, Taminiau B, Janssens GP, De Beenhouwer M, Delhalle L, Daube G, Coopman F. 2015. Cecal drop reflects the chickens' cecal microbiome, fecal drop does not. *J Microbiol Methods* 117:164–170. <https://doi.org/10.1016/j.mimet.2015.08.006>.
84. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* 79:5112–5120. <https://doi.org/10.1128/AEM.01043-13>.
85. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodriguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, et al. 2019. Author correction: Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 37:1091. <https://doi.org/10.1038/s41587-019-0252-6>.
86. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodriguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, et al. 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 37:852–857. <https://doi.org/10.1038/s41587-019-0209-9>.
87. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods* 13:581–583. <https://doi.org/10.1038/nmeth.3869>.
88. McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL, Knight R, Hugenholtz P. 2012. An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* 6:610–618. <https://doi.org/10.1038/ismej.2011.139>.
89. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Wagner H. 2018. vegan: community ecology package.
90. Lozupone CA, Hamady M, Kelley ST, Knight R. 2007. Quantitative and qualitative beta diversity measures lead to different insights into factors that structure microbial communities. *Appl Environ Microbiol* 73:1576–1585. <https://doi.org/10.1128/AEM.01996-06>.
91. Lozupone C, Knight R. 2005. UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microbiol* 71:8228–8235. <https://doi.org/10.1128/AEM.71.12.8228-8235.2005>.
92. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS One* 5:e9490. <https://doi.org/10.1371/journal.pone.0009490>.
93. Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30:3059–3066. <https://doi.org/10.1093/nar/gkf436>.
94. Mandal S, Van Treuren W, White RA, Eggesbo M, Knight R, Peddada SD. 2015. Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microb Ecol Health Dis* 26:27663. <https://doi.org/10.3402/mehd.v26.27663>.
95. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA, Gregory Caporaso J. 2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 6:90. <https://doi.org/10.1186/s40168-018-0470-z>.
96. Clermont O, Christenson JK, Denamur E, Gordon DM. 2013. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep* 5:58–65. <https://doi.org/10.1111/1758-2229.12019>.
97. Rahn K, De Grandis SA, Clarke RC, McEwen SA, Galan JE, Ginocchio C, Curtiss R, III, Gyles CL. 1992. Amplification of an *invA* gene sequence of *Salmonella typhimurium* by polymerase chain reaction as a specific method of detection of *Salmonella*. *Mol Cell Probes* 6:271–279. [https://doi.org/10.1016/0890-8508\(92\)90002-F](https://doi.org/10.1016/0890-8508(92)90002-F).