250 results for “(survey OR review) AND (wildfire OR “wild fire” OR “wildland fire” OR “wild land fire” OR “forest fire” OR bushfire) AND (propagation OR spread OR behavior OR dynamics OR trajectory) AND (prediction OR forecast OR simulation OR model OR management)”

Fields of Study | Date Range | Has PDF | More Filters | Filters | Sort by Relevance

Ensemble Modeling of CMEs Using the WSA–ENLIL+Cone Model

M. L. Mays | M. L. Mays | +12 authors | M. Kuznetsova

Physics, Environmental Science • 16 April 2015

Ensemble **modeling** of coronal mass ejections (CMEs) provides a probabilistic **forecast** of CME arrival time that includes an estimation of arrival-time uncertainty from the **spread** and distribution of… Expand

196 | [PDF] | Springer Nature | Save | Cite

Prediction modelling of riverine landscape dynamics in the context of sustainable management of floodplain: a Geospatial approach

Nasibul Alam | Subrata Saha | Srimanta Gupta | S. Chakraborty

Environmental Science, Geography • Ann. GIS • 5 January 2021

TLDR If the present rate of **land**use/**land**cover trend persists in the study stretch of Ganges river without appropriate sustainable development practice, severe floodplain degradation will ensue, accordingto the **prediction** of future **forecast**. Expand

55 | PDF | Taylor & Francis | Save | Cite

Simulation and Prediction of Category 4 and 5 Hurricanes in the High-Resolution GFDL HiFLOR Coupled Climate Model

H. Murakami | G. Vecchi | +8 authors | F. Zeng

Environmental Science • 7 December 2015

AbstractA new high-resolution Geophysical Fluid **Dynamics** Laboratory (GFDL) coupled **model** [the High-Resolution **Forecast**-Oriented Low Ocean Resolution (FLOR) **model** (HiFLOR)] has been developed and used… Expand

199 | PDF | Publisher | Save | Cite

Forecasting the Spread of COVID-19 in Kuwait Using Compartmental and Logistic Regression Models

A. Almeshal | A. Almazrouee | Mohammad R. Alenizi | S. Alhajeri

Mathematics, Environmental Science · Applied Sciences · 14 May 2020

TLDR A forecast of the COVID-19 epidemic size in Kuwait based on the confirmed data shows that a high contact rate among the population implies that the epidemic peak value is yet to be reached and that more strict intervention measures must be incorporated. Expand

77 [PDF] PDF Save Cite

LSTM-based Flight Trajectory Prediction

Zhiyuan Shi | Min Xu | Q. Pan | Bing Yan | Haimin Zhang

Engineering, Computer Science · IEEE International Joint Conference on Neural… · 1 July 2018

TLDR This paper proposes a flight trajectory prediction model based on a Long Short-Term Memory (LSTM) network that outperforms the existing models (MM and wMM) and provides a strong basis for abnormal detection and decision-making. Expand

115 PDF IEEE Save Cite

Short to Long-Term Forecasting of River Flows by Heuristic Optimization Algorithms Hybridized with ANFIS

Hossien Riahi-Madvar | Majid Dehghani | R. Memarzadeh | Bahram Gharabaghi

Environmental Science, Engineering · Water resources management · 11 February 2021

TLDR This study trained and tested the performance of several evolutionary algorithms, including Fire-Fly Algorithm(FFA), Genetic Algorithm (GA), Grey Wolf Optimization (GWO), Particle Swarmoptimization (PSO), and Differential Evolution (DE) hybridized with ANFIS, and indicated that all developed hybrid algorithms significantly outperformed the traditional ANF IS model performance for all prediction horizons. Expand

39 Springer Nature Save Cite

Current status of ENSO prediction skill in coupled ocean–atmosphere models

E. Jin | J. Kinter | +9 authors | T. Yamagata

Environmental Science · 25 April 2008

The overall skill of ENSO **prediction** in retrospective **forecasts** made with ten different coupled GCMs is investigated. The coupled GCM datasets of the APCC/CliPAS and DEMETER projects are used for… Expand

496 PDF Springer Nature Save Cite

Real-time numerical forecast of global epidemic spreading: case study of 2009 A/H1N1pdm

M. Tizzoni P. Bajardi +6 authors Alessandro Vespignani

Environmental Science, Medicine · BMC Medicine · 13 December 2012

TLDR The results show that large-scale **models** can be used to provide valuable real-time **forecasts** of influenza **spreading**, but they require high-performance computing. Expand

276 PDF Springer Nature Save Cite

Forecasting species ranges by statistical estimation of ecological niches and spatial population dynamics

J. Pagel F. Schurr Environmental Science · 1 February 2012

Aim The study and **prediction** of species–environment relationships is currently mainly based on species distribution **models**. These purely correlative **models** neglect spatial population **dynamics** and… Expand

249 Publisher Save Cite

A Data-driven Model for Interaction-Aware Pedestrian Motion Prediction in Object Cluttered Environments

Mark Pfeiffer Giuseppe Paolo H. Sommer Juan I. Nieto R. Siegwart César Cadena

Computer Science, Engineering · IEEE International Conference on Robotics and… · 25 September 2017

TLDR This paper introduces a new **model** based on Long-Short Term Memory (LSTM) neural networks, which is able to learn human motion **behavior** from demonstrated data **and** is the first approach using LSTMs, that incorporates both static obstacles **and** surrounding pedestrians **for trajectory forecasting**. Expand

106 [PDF] IEEE Save Cite

Stay Connected With Semantic Scholar

Your E-mail Address

Sign Up

What Is Semantic Scholar?

Semantic Scholar is a free, AI-powered research tool for scientific literature, based at Ai2.

Learn More

About

About Us
Publishers
Blog
Ai2 Careers

Product

Product Overview
Semantic Reader
Scholar's Hub
Beta Program
Release Notes

API

API Overview
API Tutorials
API Documentation
API Gallery

Research

Publications
Research Careers
Resources

Help

FAQ
Librarians
Tutorials
Contact

Proudly built by Ai2

Collaborators & Attributions • Terms of Service • Privacy Policy • API License Agreement

Ai2