

Unit 6: Introduction to linear regression

2. Outliers and inference for regression

GOVT 3990 - Spring 2020

Cornell University

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

Uncertainty of predictions

- ▶ Regression models are useful for making predictions for new observations not include in the original dataset.

Uncertainty of predictions

- ▶ Regression models are useful for making predictions for new observations not include in the original dataset.
- ▶ If the model is good, the predictions should be close to the true value of the response variable for this observation, however it may not be exact, i.e. \hat{y} might be different than y .

Uncertainty of predictions

- ▶ Regression models are useful for making predictions for new observations not include in the original dataset.
- ▶ If the model is good, the predictions should be close to the true value of the response variable for this observation, however it may not be exact, i.e. \hat{y} might be different than y .
- ▶ With any prediction we can (and should) also report a measure of uncertainty of the prediction.

Prediction intervals for specific predicted values

A *prediction interval* for y for a given x^* is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where s is the standard deviation of the residuals, and x^* is a new observation.

Prediction intervals for specific predicted values

A *prediction interval* for y for a given x^* is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where s is the standard deviation of the residuals, and x^* is a new observation.

- Interpretation: We are XX% confident that \hat{y} for given x^* is within this interval.

Prediction intervals for specific predicted values

A *prediction interval* for y for a given x^* is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where s is the standard deviation of the residuals, and x^* is a new observation.

- ▶ Interpretation: We are XX% confident that \hat{y} for given x^* is within this interval.
- ▶ The width of the prediction interval for \hat{y} increases as
 - x^* moves away from the center
 - s (the variability of residuals), i.e. the scatter, increases

Prediction intervals for specific predicted values

A *prediction interval* for y for a given x^* is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where s is the standard deviation of the residuals, and x^* is a new observation.

- Interpretation: We are XX% confident that \hat{y} for given x^* is within this interval.
- The width of the prediction interval for \hat{y} increases as
 - x^* moves away from the center
 - s (the variability of residuals), i.e. the scatter, increases
- Prediction level: If we repeat the study of obtaining a regression data set many times, each time forming a XX% prediction interval at x^* , and wait to see what the future

Calculating the prediction interval

By hand:

Don't worry about it...

Calculating the prediction interval

By hand:

Don't worry about it...

In R:

```
# predict  
predict(m_mur_pov, newdata, interval = "prediction", level = 0.95)
```

Calculating the prediction interval

By hand:

Don't worry about it...

In R:

```
# predict  
predict(m_mur_pov, newdata, interval = "prediction", level = 0.95)
```

	fit	lwr	upr
1	21.28663	9.418327	33.15493

Calculating the prediction interval

By hand:

Don't worry about it...

In R:

```
# predict  
predict(m_mur_pov, newdata, interval = "prediction", level = 0.95)
```

	fit	lwr	upr
1	21.28663	9.418327	33.15493

We are 95% confident that the annual murders per million for a county with 20% poverty rate is between 9.52 and 33.15.

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

(1) R^2 assesses model fit – higher the better

- ▶ R^2 : percentage of variability in y explained by the model.

(1) R^2 assesses model fit – higher the better

- ▶ R^2 : percentage of variability in y explained by the model.
- ▶ For single predictor regression: R^2 is the square of the correlation coefficient, R .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

(1) R^2 assesses model fit – higher the better

- ▶ R^2 : percentage of variability in y explained by the model.
- ▶ For single predictor regression: R^2 is the square of the correlation coefficient, R .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

- ▶ For all regression: $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual_murders_per_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

(1) R^2 assesses model fit – higher the better

- For all regression: $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual_murders_per_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}}$$

(1) R^2 assesses model fit – higher the better

- For all regression: $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual_murders_per_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{SS_{reg}}{SS_{tot}}$$

(1) R^2 assesses model fit – higher the better

- For all regression: $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual_murders_per_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{SS_{reg}}{SS_{tot}} = \frac{1308.34}{1308.34 + 546.86}$$

(1) R^2 assesses model fit – higher the better

- For all regression: $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table


Response: annual_murders_per_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{SS_{reg}}{SS_{tot}} = \frac{1308.34}{1308.34 + 546.86} = \frac{1308.34}{1855.2} \approx 0.71$$

Your turn


R^2 for the regression model for predicting annual murders per million based on percentage living in poverty is roughly 71%. Which of the following is the correct interpretation of this value?



- (a) 71% of the variability in percentage living in poverty is explained by the model.
- (b) 84% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.
- (c) 71% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.
- (d) 71% of the time percentage living in poverty predicts murder rates accurately.

Your turn

R^2 for the regression model for predicting annual murders per million based on percentage living in poverty is roughly 71%. Which of the following is the correct interpretation of this value?



- (a) 71% of the variability in percentage living in poverty is explained by the model.
- (b) 84% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.
- (c) *71% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.*
- (d) 71% of the time percentage living in poverty predicts murder rates accurately.

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

Inference for regression uses the t -distribution

- ▶ Use a T distribution for inference on the slope, with degrees of freedom $n - 2$
 - Degrees of freedom for the slope(s) in regression is $df = n - k - 1$ where k is the number of slopes being estimated in the model.

Inference for regression uses the t -distribution

- ▶ Use a T distribution for inference on the slope, with degrees of freedom $n - 2$
 - Degrees of freedom for the slope(s) in regression is $df = n - k - 1$ where k is the number of slopes being estimated in the model.
- ▶ Hypothesis testing for a slope: $H_0 : \beta_1 = 0$;
 $H_A : \beta_1 \neq 0$
 - $T_{n-2} = \frac{b_1 - 0}{SE_{b_1}}$
 - p-value = P(observing a slope at least as different from 0 as the one observed if in fact there is no relationship between x and y)

Inference for regression uses the t -distribution

- ▶ Use a T distribution for inference on the slope, with degrees of freedom $n - 2$
 - Degrees of freedom for the slope(s) in regression is $df = n - k - 1$ where k is the number of slopes being estimated in the model.
- ▶ Hypothesis testing for a slope: $H_0 : \beta_1 = 0$;
 $H_A : \beta_1 \neq 0$
 - $T_{n-2} = \frac{b_1 - 0}{SE_{b_1}}$
 - p-value = P(observing a slope at least as different from 0 as the one observed if in fact there is no relationship between x and y)
- ▶ Confidence intervals for a slope:
 - $b_1 \pm T_{n-2}^* SE_{b_1}$
 - In R:

```
confint(m_mur_pov, level = 0.95)
```

Inference for regression uses the t -distribution

► Confidence intervals for a slope:

- $b_1 \pm T_{n-2}^* SE_{b_1}$
- In R:

```
confint(m_mur_pov, level = 0.95)
```

	2.5 %	97.5 %
(Intercept)	-46.265631	-13.536694
perc_pov	1.740003	3.378776

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

Conditions for regression

Important regardless of doing inference

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

Conditions for regression

Important regardless of doing inference

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

Important for inference

- ▶ Nearly normally distributed residuals → histogram or normal probability plot of residuals

Conditions for regression

Important regardless of doing inference

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

Important for inference

- ▶ Nearly normally distributed residuals → histogram or normal probability plot of residuals
- ▶ Constant variability of residuals (*homoscedasticity*) → no fan shape in the residuals plot

Conditions for regression

Important regardless of doing inference

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

Important for inference


- ▶ Nearly normally distributed residuals → histogram or normal probability plot of residuals
- ▶ Constant variability of residuals (*homoscedasticity*) → no fan shape in the residuals plot
- ▶ Independence of residuals (and hence observations) → depends on data collection method, often violated for time-series data

Checking conditions

Your turn

What condition is this linear model obviously and definitely violating?

- (a) Linear relationship
- (b) Non-normal residuals
- (c) Constant variability
- (d) Independence of observations




Checking conditions

Your turn

What condition is this linear model obviously and definitely violating?

- (a) Linear relationship
- (b) Non-normal residuals
- (c) *Constant variability*
- (d) Independence of observations




Checking conditions

Your turn

What condition is this linear model obviously and definitely violating?

- (a) Linear relationship
- (b) Non-normal residuals
- (c) Constant variability
- (d) Independence of observations




Checking conditions

Your turn

What condition is this linear model obviously and definitely violating?

- (a) *Linear relationship*
- (b) Non-normal residuals
- (c) Constant variability
- (d) Independence of observations



Outline

1. Housekeeping


2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary


Type of outlier determines how it should be handled

- ▶ *Leverage* point is away from the cloud of points horizontally, does not necessarily change the slope
- ▶ *Influential* point changes the slope (most likely also has high leverage) – run the regression with and without that point to determine



Type of outlier determines how it should be handled

- ▶ *Leverage* point is away from the cloud of points horizontally, does not necessarily change the slope
- ▶ *Influential* point changes the slope (most likely also has high leverage) – run the regression with and without that point to determine
- ▶ *Outlier* is an unusual point without these special characteristics (this one likely affects the intercept only)
- ▶ If clusters (groups of points) are apparent in the data, it might be worthwhile to model the groups separately.



Application exercise: 6.2 Linear regression

See course website for details

Outline

1. Housekeeping

2. Main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled

3. Summary

Summary of main ideas

1. Predicted values also have uncertainty around them
2. R^2 assesses model fit – higher the better
3. Inference for regression uses the t -distribution
4. Conditions for regression
5. Type of outlier determines how it should be handled