

# Is essentiality of genes conserved in Enterobacteriaceae?

Fatemeh Ashari-Ghomi <sup>\*</sup>, Paul P. Gardner <sup>\*</sup> and Lars Barquist <sup>†</sup>

<sup>\*</sup>University of Canterbury, and <sup>†</sup>Wurzburg University

Submitted to Proceedings of the National Academy of Sciences of the United States of America

essentiality | conservation | transposon insertion

## Introduction

**S**tudying the essentiality of genes helps with identifying the fundamental processes necessary for cell viability [1]. So far, scientists have studied the essential genes in organisms from different domains of life [2]. The results have led to new insights for developing new antibiotics that target essential genes of pathogenic bacteria [3, 4] and synthesising new genomes [5, 6]. Researchers have used different methods for studying the essentiality of genes in prokaryotes. Baba et al. [7] have made a library of single gene deletions using phage lambda Red recombination system to screen essential genes while another group have used antisense RNA knockdowns for this purpose [8]. Another method that is widely used due to its simplicity and accuracy is transposon mutagenesis along with high-throughput sequencing [9, 10, 11, 12, 13, 14, 15]. In this method, pools of single insertion mutants are constructed using transposon mutagenesis and the effect of each mutation on the survival of mutants is evaluated by sequencing the survivors [16]. This can lead to the identification of essential genes.

Although the essentiality of genes has been studied in a variety of organisms, there is still room to study the evolutionary conservation of essentiality. Barquist et al. [17] have used transposon-directed insertion-site sequencing to study the differentiation of the essentiality of genes in *Salmonella* serovars Typhi and Typhimurium which has led to divergence in their pathogenicity and host ranges. We extend this research by studying 13 bacterial strains from Enterobacteriaceae. These strains are depicted in Fig. 1.

Enterobacteriaceae is a family that includes bacteria with different host ranges and pathogenicity found in soil, water, plants, animals and humans [18]. In humans, various strains from this family can cause diarrhoea, septicaemia, urinary tract infection, meningitis, respiratory disease, and wound and burn infection [18]. Besides, they can infect poultry and livestock and cause financial losses for farmers [18]. Here, we perform a transposon-directed insertion-site sequencing experiment to study the conservation of essentiality of genes in strains from 5 different species in this family.

{A summary of what we have done}

## Transposon mutagenesis

Strains.

Transposon insertion experiment.

Essentiality.

## Are transposons biased towards certain positions?

Different articles have reported biases in transposon mutagenesis [17, 15]. We have performed a thorough study of these biases. To study the bias towards the position of the genes,

we plotted the insertion index for each gene versus the distance of the gene from the origin of replication normalised by the length of the genome. Fig. 2 shows the results. The red line is a loess curve that has been fitted to the data when the smoothness parameter equals 0.2. The figure indicates that the insertion indices decrease when the genes are located further from the origin of replication. A possible explanation for this phenomenon is that the bacteria were under replication while being infected with transposons. Therefore, the number of gene copies close to the origin of replication was greater and more insertions have occurred in these genes.

We have tested whether our transposons are biased towards certain motifs. For this, we have generated a logo from 10 nucleotides flanking the 100 top most frequent insertion sites in each genome. The results are depicted in Fig. 3. The results show a slight bias towards certain combinations of bases. In addition, we have investigated if the G-C content of genes can change the number of insertions by plotting the number of G-C bases in a gene normalised by the length of the gene versus insertion index. The red lines show the loess curve when the smoothness parameter is 0.2. {To be completed}

The other bias that we have considered is the bias towards certain locations in a gene. We have divided every gene into 100 bins and calculated the mean insertion index for each bin. Fig. 5a shows almost no bias towards any location. We have also divided our genes into three groups - essential genes, non-essential genes, and beneficial losses - and studied the bias in each group. The results imply that the number of insertions in the internal region of the essential genes is outnumbered by the number of insertions in the 5' and 3' ends while it is the opposite in beneficial losses. The case for the non-essential genes is similar to the average (Fig. 5a). High number of insertions at the 3' end of essential genes implies that the functional part of the genes are located before the insertions. On the other hand, high number of insertions at the 5' end of the essential genes indicates there might be alternative start codons in the 5' end or it might be because of alignment errors. {To be tested}

## Gene classes

Genus specific.

Single copy.

Reserved for Publication Footnotes

## Multi-copy.

### Evolution of essentiality

- UpSetR results (Fig. 6)
- Stringent
- Dollo law
- Ancestral insertion index

## Case study of genes

### Core genes.

### Accessory genes. ACKNOWLEDGMENTS.

1. Mario Juhas, Leo Eberl, and John I. Glass. Essence of life: essential genes of minimal genomes. 21(10):562–568.
2. Hao Luo, Yan Lin, Feng Gao, Chun-Ting Zhang, and Ren Zhang. DEG 10, an update of the database of essential genes that includes both protein-coding genes and noncoding genomic elements. 42:D574–D580.
3. Anne E. Clatworthy, Emily Pierson, and Deborah T. Hung. Targeting virulence: a new paradigm for antimicrobial therapy. 3(9):541–548.
4. Jason M. Peters, Alexandre Colavin, Handuo Shi, Tomasz L. Czarny, Matthew H. Larson, Spencer Wong, John S. Hawkins, Candy H. S. Lu, Byoung-Mo Koo, Elizabeth Marta, Anthony L. Shiver, Evan H. Whitehead, Jonathan S. Weissman, Eric D. Brown, Lei S. Qi, Kerwyn Casey Huang, and Carol A. Gross. A comprehensive, CRISPR-based functional analysis of essential genes in bacteria. 165(6):1493–1506.
5. Clyde A. Hutchison, Scott N. Peterson, Steven R. Gill, Robin T. Cline, Owen White, Claire M. Fraser, Hamilton O. Smith, and J. Craig Venter. Global transposon mutagenesis and a minimal mycoplasma genome. 286(5447):2165–2169.
6. Clyde A. Hutchison, Ray-Yuan Chuang, Vladimir N. Noskov, Nacyra Assad-Garcia, Thomas J. Deerinck, Mark H. Ellisman, John Gill, Krishna Kannan, Bogumil J. Karas, Li Ma, James F. Pelletier, Zhi-Qing Qi, R. Alexander Richter, Elizabeth A. Strychalski, Lijie Sun, Yo Suzuki, Billyana Tsvetanova, Kim S. Wise, Hamilton O. Smith, John I. Glass, Chuck Merryman, Daniel G. Gibson, and J. Craig Venter. Design and synthesis of a minimal bacterial genome. 351(6280):aad6253.
7. Tomoya Baba, Takeshi Ara, Miki Hasegawa, Yuki Takai, Yoshiko Okumura, Miki Baba, Kirill A Datsenko, Masaru Tomita, Barry L Wanner, and Hirotsada Mori. Construction of escherichia coli k-12 in-frame, single-gene knockout mutants: the keio collection. 2:2006.0008.
8. H. Howard Xu, John D. Trawick, Robert J. Haselbeck, R. Allyn Forsyth, Robert T. Yamamoto, Rich Archer, Joe Patterson, Molly Allen, Jamie M. Froelich, Ian Taylor, Danny Nakaji, Randy Maile, G. C. Kedar, Marshall Pilcher, Vickie Brown-Driver, Melissa McCarthy, Amy Files, David Robbins, Paula King, Susan Sillaots, Cheryl Malone, Carlos S. Zamudio, Terry Roemer, Liangsu Wang, Philip J. Youngman, and Daniel Wall. Staphylococcus aureus TargetArray: comprehensive differential essential gene expression as a mechanistic tool to profile antibacterials. 54(9):3659–3670.
9. Jeffrey D. Gawronski, Sandy M. S. Wong, Georgia Giannoukos, Doyle V. Ward, and Brian J. Akerley. Tracking insertion mutants within libraries by deep sequencing and a genome-wide screen for haemophilus genes required in the lung. 106(38):16422–16427.
10. Tim van Opijnen, Kip L. Bodi, and Andrew Camilli. Tn-seq: high-throughput parallel sequencing for fitness and genetic interaction studies in microorganisms. 6(10):767–772.
11. Gemma C. Langridge, Minh-Duy Phan, Daniel J. Turner, Timothy T. Perkins, Leopold Parts, Jana Haase, Ian Charles, Duncan J. Maskell, Sarah E. Peters, Gordon Dougan, John Wain, Julian Parkhill, and A. Keith Turner. Simultaneous assay of every salmonella typhi gene using one million transposon mutants. 19(12):2308–2316.
12. Beat Christen, Eduardo Abeliuk, John M Collier, Virginia S Kalogeraki, Ben Passarelli, John A Collier, Michael J Fero, Harley H McAdams, and Lucy Shapiro. The essential genome of a bacterium. 7:528.
13. Andrew L. Goodman, Meng Wu, and Jeffrey I. Gordon. Identifying microbial fitness determinants by insertion sequencing using genome-wide transposon mutant libraries. 6(12):1969–1980.
14. Kelly M. Wetmore, Morgan N. Price, Robert J. Waters, Jacob S. Lamson, Jennifer He, Cindi A. Hoover, Matthew J. Blow, James Bristow, Gareth Butland, Adam P. Arkin, and Adam Deutschbauer. Rapid quantification of mutant fitness in diverse bacteria by sequencing randomly bar-coded transposons. 6(3):e00306–15.
15. Benjamin E. Rubin, Kelly M. Wetmore, Morgan N. Price, Spencer Diamond, Ryan K. Shultzaberger, Laura C. Lowe, Genevieve Curtin, Adam P. Arkin, Adam Deutschbauer, and Susan S. Golden. The essential gene set of a photosynthetic organism. 112(48):E6634–E6643.
16. Lars Barquist, Christine J. Boinett, and Amy K. Cain. Approaches to querying bacterial genomes with transposon-insertion sequencing. 10(7):1161–1169.
17. Lars Barquist, Gemma C. Langridge, Daniel J. Turner, Minh-Duy Phan, A. Keith Turner, Alex Bateman, Julian Parkhill, John Wain, and Paul P. Gardner. A comparison of dense transposon insertion libraries in the salmonella serovars typhi and typhimurium. page gkt148.
18. Don J. Brenner and Noel R. Krieg. Bergey’s Manual of Systematic Bacteriology: Volume Two: The Proteobacteria. Springer Science & Business Media.
19. Adrian M. Altenhoff, Brigitte Boeckmann, Salvador Capella-Gutierrez, Daniel A. Dalquen, Todd DeLuca, Kristoffer Forslund, Jaime Huerta-Cepas, Benjamin Linard, Ccile Pereira, Leszek P. Pryszcz, Fabian Schreiber, Alan Sousa da Silva, Damian Szklarczyk, Clment-Marie Train, Peer Bork, Odile Lecompte, Christian von Mering, Ioannis Xenarios, Kimmen Sjlinder, Lars Juhl Jensen, Maria J. Martin, Matthieu Muffato, Quest for Orthologs Consortium, Toni Gabaldn, Suzanna E. Lewis, Paul D. Thomas, Erik Sonnhammer, and Christophe Dessimoz. Standardized benchmarking in the quest for orthologs. 13(5):425–430.
20. Fabian Schreiber and Erik L. L. Sonnhammer. Hieranoid: hierarchical orthology inference. 425(11):2072–2081.
21. Jaina Mistry, Robert D. Finn, Sean R. Eddy, Alex Bateman, and Marco Punta. Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. 41(12):e121–e121.
22. Alexandros Stamatakis. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. 30(9):1312–1313.
23. Aaron E. Darling, Guillaume Jospin, Eric Lowe, Frederick A. Matsen, Holly M. Bik, and Jonathan A. Eisen. PhyloSift: phylogenetic analysis of genomes and metagenomes. 2:e243.




Fig. 1: The species tree containing the 13 strains under study and *Escherichia coli* K-12 MG1655 studied in Keio collection [7]. We have generated the tree by running RAxML [22] on Phylosift [23] amino acid markers.




Fig. 2: The plots show the position of the genes within the genome (normalised by the lengths of the genomes) versus the insertion indices of the genes




Fig. 3: We have generated the logos from 10 nucleotides flanking the 100 top most frequent insertion sites.




Fig. 4: The plots show the G-C contents of the genes (normalised by the lengths of the genes) against their insertion indices




Fig. 5: We have divided our genes into 3 segments: 5% of the genes on the 5' end, 20% of the genes on the 3' end, and the rest in the middle.




Fig. 6: The first figure shows the number of core genes between each group of species and the second figure shows the number of core essential genes.