

Is essentiality of genes conserved in Enterobacteriaceae?

Fatemeh Ashari-Ghomi ^{*}, Paul P. Gardner ^{*} and Lars Barquist [†]

^{*}University of Canterbury, and [†]Wurzburg University

Submitted to Proceedings of the National Academy of Sciences of the United States of America

essentiality | conservation | transposon insertion

Introduction

Studying the essentiality of genes helps with identifying the fundamental processes necessary for cell viability [1]. So far, scientists have studied the essential genes in organisms from different domains of life [2]. The results have led to new insights for developing new antibiotics that target essential genes of pathogenic bacteria [3, 4] and synthesising new genomes [5, 6]. Researchers have used different methods for studying the essentiality of genes in prokaryotes. Baba et al. [7] have made a library of single gene deletions using phage lambda Red recombination system to screen essential genes while another group have used antisense RNA knockdowns for this purpose [8]. Another method that is widely used due to its simplicity and accuracy is transposon mutagenesis along with high-throughput sequencing [9, 10, 11, 12, 13, 14, 15]. In this method, pools of single insertion mutants are constructed using transposon mutagenesis and the effect of each mutation on the survival of mutants is evaluated by sequencing the survivors [16]. This can lead to the identification of essential genes.

Although the essentiality of genes has been studied in a variety of organisms, there is still room to study the evolutionary conservation of essentiality. Barquist et al. [17] have used transposon-directed insertion-site sequencing to study the differentiation of the essentiality of genes in *Salmonella* serovars Typhi and Typhimurium which has led to divergence in their pathogenicity and host ranges. We extend this research by studying 13 bacterial strains from Enterobacteriaceae. These strains are depicted in Fig. 1.

Enterobacteriaceae is a family that includes bacteria with different host ranges and pathogenicity found in soil, water, plants, animals and humans [18]. In humans, various strains from this family can cause diarrhoea, septicaemia, urinary tract infection, meningitis, respiratory disease, and wound and burn infection [18]. Besides, they can infect poultry and live-stock and cause financial losses for farmers [18]. Here, we perform a transposon-directed insertion-site sequencing experiment to study the conservation of essentiality of genes in strains from 5 different species in this family.

{A summary of what we have done}

1. Mario Juhas, Leo Eberl, and John I. Glass. Essence of life: essential genes of minimal genomes. 21(10):562–568.
2. Hao Luo, Yan Lin, Feng Gao, Chun-Ting Zhang, and Ren Zhang. DEG 10, an update of the database of essential genes that includes both protein-coding genes and noncoding genomic elements. 42:D574–D580.
3. Anne E. Clatworthy, Emily Pierson, and Deborah T. Hung. Targeting virulence: a new paradigm for antimicrobial therapy. 3(9):541–548.
4. Jason M. Peters, Alexandre Colavin, Handuo Shi, Tomasz L. Czarny, Matthew H. Larson, Spencer Wong, John S. Hawkins, Candy H. S. Lu, Byoung-Mo Koo, Elizabeth Marta, Anthony L. Shiver, Evan H. Whitehead, Jonathan S. Weissman, Eric D. Brown, Lei S. Qi, Kerwyn Casey Huang, and Carol A. Gross. A comprehensive, CRISPR-based functional analysis of essential genes in bacteria. 165(6):1493–1506.

Are transposons biased towards certain positions?

- There is a bias towards the position of the gene (Fig. 2).
- There is a negligible bias towards certain motifs (Fig. 3)
- The G-C bias needs to be studied further (Fig. 4)
- The number of insertions on the 3 and 5 ends is more than the internal region in essential genes and less than the internal region in beneficial losses. (Fig. 5)

Gene classes

Genus specific.

Single copy.

Multi-copy.

Evolution of essentiality

- UpSetR results (Fig. 6)
- Stringent
- Dollo law
- Ancestral insertion index

Case study of genes

Core genes.

Accessory genes.

Discussion

Materials and Methods

Clustering orthologous and paralogous genes. To study the essentiality of genes in 12 our strains, We needed to cluster sets of orthologous genes in these strain. Plenty of methods are proposed for this purpose. Altenhoff et al. have compared 15 of these methods [19] and shown that Hieranoid [20] is among three methods that keep a balance between precision and recall. We have used Hieranoid to cluster the sets of orthologous genes. In addition, we intended to study the essentiality of genes in paralogous genes. For this, we have developed a program that clusters all homologous genes using Jackhmmer from HMMER3 package [21].

ACKNOWLEDGMENTS.

—5—Clyde A. Hutchison, Scott N. Peterson, Steven R. Gill, Robin T. Cline, Owen White, Reserved for Publication Footnotes

- Claire M. Fraser, Hamilton O. Smith, and J. Craig Venter. Global transposon mutagenesis and a minimal mycoplasma genome. 286(5447):2165–2169.
6. Clyde A. Hutchison, Ray-Yuan Chuang, Vladimir N. Noskov, Nacyra Assad-Garcia, Thomas J. Deerinck, Mark H. Ellisman, John Gill, Krishna Kannan, Bogumil J. Karas, Li Ma, James F. Pelletier, Zhi-Qing Qi, R. Alexander Richter, Elizabeth A. Strychalski, Lijie Sun, Yo Suzuki, Billyana Tsvetanova, Kim S. Wise, Hamilton O. Smith, John I. Glass, Chuck Merryman, Daniel G. Gibson, and J. Craig Venter. Design and synthesis of a minimal bacterial genome. 351(6280):aad6253.
7. Tomoya Baba, Takeshi Ara, Miki Hasegawa, Yuki Takai, Yoshiko Okumura, Miki Baba, Kirill A Datsenko, Masaru Tomita, Barry L Wanner, and Hirota Mori. Construction of *escherichia coli* k-12 in-frame, single-gene knockout mutants: the keio collection. 2:2006.0008.
8. H. Howard Xu, John D. Trawick, Robert J. Haselbeck, R. Allyn Forsyth, Robert T. Yamamoto, Rich Archer, Joe Patterson, Molly Allen, Jamie M. Froelich, Ian Taylor, Danny Nakaji, Randy Maile, G. C. Kedar, Marshall Pilcher, Vickie Brown-Driver, Melissa McCarthy, Amy Files, David Robbins, Paula King, Susan Sillaots, Cheryl Malone, Carlos S. Zamudio, Terry Roemer, Liangsu Wang, Philip J. Youngman, and Daniel Wall. *Staphylococcus aureus* TargetArray: comprehensive differential essential gene expression as a mechanistic tool to profile antibacterials. 54(9):3659–3670.
9. Jeffrey D. Gawronski, Sandy M. S. Wong, Georgia Giannoukos, Doyle V. Ward, and Brian J. Akerley. Tracking insertion mutants within libraries by deep sequencing and a genome-wide screen for *haemophilus* genes required in the lung. 106(38):16422–16427.
10. Tim van Opijnen, Kip L. Bodi, and Andrew Camilli. Tn-seq: high-throughput parallel sequencing for fitness and genetic interaction studies in microorganisms. 6(10):767–772.
11. Gemma C. Langridge, Minh-Duy Phan, Daniel J. Turner, Timothy T. Perkins, Leopold Parts, Jana Haase, Ian Charles, Duncan J. Maskell, Sarah E. Peters, Gordon Dougan, John Wain, Julian Parkhill, and A. Keith Turner. Simultaneous assay of every *salmonella typhi* gene using one million transposon mutants. 19(12):2308–2316.
12. Beat Christen, Eduardo Abeliuk, John M Collier, Virginia S Kalogeraki, Ben Passarelli, John A Collier, Michael J Fero, Harley H McAdams, and Lucy Shapiro. The essential genome of a bacterium. 7:528.
13. Andrew L. Goodman, Meng Wu, and Jeffrey I. Gordon. Identifying microbial fitness determinants by insertion sequencing using genome-wide transposon mutant libraries. 6(12):1969–1980.
14. Kelly M. Wetmore, Morgan N. Price, Robert J. Waters, Jacob S. Lamson, Jennifer He, Cindi A. Hoover, Matthew J. Blow, James Bristow, Gareth Butland, Adam P. Arkin, and Adam Deutschbauer. Rapid quantification of mutant fitness in diverse bacteria by sequencing randomly bar-coded transposons. 6(3):e00306–15.
15. Benjamin E. Rubin, Kelly M. Wetmore, Morgan N. Price, Spencer Diamond, Ryan K. Shultzaberger, Laura C. Lowe, Genevieve Curtin, Adam P. Arkin, Adam Deutschbauer, and Susan S. Golden. The essential gene set of a photosynthetic organism. 112(48):E6634–E6643.
16. Lars Barquist, Christine J. Boinett, and Amy K. Cain. Approaches to querying bacterial genomes with transposon-insertion sequencing. 10(7):1161–1169.
17. Lars Barquist, Gemma C. Langridge, Daniel J. Turner, Minh-Duy Phan, A. Keith Turner, Alex Bateman, Julian Parkhill, John Wain, and Paul P. Gardner. A comparison of dense transposon insertion libraries in the *salmonella* serovars typhi and typhimurium. page gkt148.
18. Don J. Brenner and Noel R. Krieg. *Bergey's Manual of Systematic Bacteriology: Volume Two: The Proteobacteria*. Springer Science & Business Media.
19. Adrian M. Altenhoff, Brigitte Boeckmann, Salvador Capella-Gutierrez, Daniel A. Dalquen, Todd DeLuca, Kristoffer Forslund, Jaime Huerta-Cepas, Benjamin Linard, Cile Pereira, Leszek P. Pryszcz, Fabian Schreiber, Alan Sousa da Silva, Damian Szklarczyk, Clment-Marie Train, Peer Bork, Odile Lecompte, Christian von Mering, Ioannis Xenarios, Kimmen Sjlinder, Lars Juhl Jensen, Maria J. Martin, Matthieu Muffato, Quest for Orthologs Consortium, Toni Gabaldn, Suzanna E. Lewis, Paul D. Thomas, Erik Sonnhammer, and Christophe Dessimoz. Standardized benchmarking in the quest for orthologs. 13(5):425–430.
20. Fabian Schreiber and Erik L. L. Sonnhammer. Hieranoid: hierarchical orthology inference. 425(11):2072–2081.
21. Jaina Mistry, Robert D. Finn, Sean R. Eddy, Alex Bateman, and Marco Punta. Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. 41(12):e121–e121.
22. Alexandros Stamatakis. RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. 30(9):1312–1313.
23. Aaron E. Darling, Guillaume Jospin, Eric Lowe, Frederick A. Matsen, Holly M. Bik, and Jonathan A. Eisen. PhyloSift: phylogenetic analysis of genomes and metagenomes. 2:e243.




Fig. 1: The species tree containing the 13 strains under study and *Escherichia coli* K-12 MG1655 studied in Keio collection [7]. We have generated the tree by running RAxML [22] on Phylosift [23] amino acid markers.




Fig. 2: The plots show the position of the genes within the genome (normalised by the lengths of the genomes) versus the insertion indices of the genes




Fig. 3: We have generated the logos from 10 nucleotides flanking the 100 top most frequent insertion sites.




Fig. 4: The plots show the G-C contents of the genes (normalised by the lengths of the genes) against their insertion indices




Fig. 5: We have divided our genes into 3 segments: 5% of the genes on the 5 end, 20% of the genes on the 3 end, and the rest in the middle.




Fig. 6: The first figure shows the number of core genes between each group of species and the second figure shows the number of core essential genes.