

Exercise 2.8

Wang Yue from CS Elite Class

October 24, 2020

Find the derivative of the function using the definition of derivative. State the domain of the function and the domain of its derivative.

26. $f(x) = x + \sqrt{x}$

Solution:

The domain of $f(x)$ is $[0, \infty)$.

$$\begin{aligned} f'(x_0) &= \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \\ &= \lim_{x \rightarrow x_0} \frac{x - x_0 + \sqrt{x} - \sqrt{x_0}}{x - x_0} \\ &= 1 + \lim_{x \rightarrow x_0} \frac{\sqrt{x} - \sqrt{x_0}}{x - x_0} \\ &= 1 + \lim_{x \rightarrow x_0} \frac{1}{\sqrt{x} + \sqrt{x_0}} \\ &= \frac{1}{2\sqrt{x_0}} + 1 \end{aligned}$$

$\therefore f'(x) = \frac{1}{2\sqrt{x}} + 1$, and the domain of $f'(x)$ is $(0, \infty)$.

27. $g(x) = \sqrt{9 - x}$

Solution:

The domain of $g(x)$ is the solution set of $9 - x \geq 0$, which is $(-\infty, 9]$.

$$\begin{aligned}
g'(x_0) &= \lim_{x \rightarrow x_0} \frac{\sqrt{9-x} - \sqrt{9-x_0}}{x - x_0} \\
&= \lim_{x \rightarrow x_0} \frac{9-x - (9-x_0)}{(x-x_0)(\sqrt{9-x} + \sqrt{9-x_0})} \\
&= \lim_{x \rightarrow x_0} \frac{-1}{\sqrt{9-x} + \sqrt{9-x_0}} \\
&= \frac{-1}{2\sqrt{9-x_0}}
\end{aligned}$$

$\therefore g'(x) = \frac{-1}{2\sqrt{9-x}}$, and the domain of $g'(x)$ is $(-\infty, 9)$.

29. $G(t) = \frac{1-2t}{3+t}$

Solution:

The domain of $G(t)$ is $\{t \in R | t \neq -3\}$.

$$G(t) = \frac{-6-2t+7}{3+t} = \frac{7}{t+3} - 2$$

$$\begin{aligned}
G'(t_0) &= \lim_{t \rightarrow t_0} \frac{\frac{7}{t+3} - \frac{7}{t_0+3}}{t - t_0} \\
&= 7 \lim_{t \rightarrow t_0} \frac{\frac{t_0-t}{(t+3)(t_0+3)}}{t - t_0} \\
&= -7 \lim_{t \rightarrow t_0} \frac{1}{(t+3)(t_0+3)} \\
&= \frac{-7}{(t_0+3)^2}
\end{aligned}$$

$\therefore G'(t) = \frac{-7}{(t+3)^2}$, the domain of which is also $(-\infty, -3) \cup (-3, \infty)$

51. Let $f(x) = \sqrt[3]{x}$.

(a) If $a \neq 0$, use Equation 2.7.5 to find $f'(a)$.

Solution:

$$\begin{aligned}
f'(a) &= \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \\
&= \lim_{x \rightarrow a} \frac{\sqrt[3]{x} - \sqrt[3]{a}}{x - a} \\
&= \lim_{x \rightarrow a} \frac{(x^{\frac{1}{3}} - a^{\frac{1}{3}})(x^{\frac{2}{3}} + (ax)^{\frac{1}{3}} + a^{\frac{2}{3}})}{(x - a)(x^{\frac{2}{3}} + (ax)^{\frac{1}{3}} + a^{\frac{2}{3}})} \\
&= \lim_{x \rightarrow a} \frac{1}{x^{\frac{2}{3}} + (ax)^{\frac{1}{3}} + a^{\frac{2}{3}}} \\
&= \frac{1}{3(a)^{\frac{2}{3}}} \\
&= \frac{1}{3}a^{-\frac{2}{3}}
\end{aligned}$$

(b) Show that $f'(0)$ does not exist.

Solution:

\because by problem (a), we can know the domain of $f'(a)$ is $(-\infty, 0) \cup (0, \infty)$

$\therefore f'(0)$ has no definition, which means $f'(0)$ does not exist.

(c) Show that $y = \sqrt[3]{x}$ has a vertical tangent line at $(0, 0)$.

Solution:

$\because \lim_{x \rightarrow 0} f'(x) = \lim_{x \rightarrow 0} \frac{1}{3x^{\frac{2}{3}}} = \infty$

\therefore the tangent line of $f(x)$ as $x \rightarrow 0$ is vertical

$\because \sqrt[3]{0} = 0$


$\therefore y = \sqrt[3]{x}$ has a vertical tangent line at $(0, 0)$

55.

(a) Sketch the graph of the function $f(x) = x|x|$.

Solution:

The graph of $f(x) = x|x|$ is below.



(b) For what values of x is f differentiable?

$$\therefore f(x) = \begin{cases} x^2 & \text{if } x \geq 0 \\ -x^2 & \text{if } x < 0 \end{cases}$$

$$\therefore f'(x) = \begin{cases} 2x & \text{if } x \geq 0 \\ -2x & \text{if } x < 0 \end{cases}$$

So obviously, for any value of x , $f(x)$ is differentiable.

(c) Find a formula for f' .

By the problem (b), a formula for f' can be

$$f'(x) = 2|x|$$

57. Prove each of the following.

(a) The derivative of an even function is an odd function.

Proof. $\because f(x) = f(-x)$

\therefore taking derivative on the both side, we can get

$$f'(x) = -f'(-x)$$

which means $f'(x)$ is an odd function. □

(b) The derivative of an odd function is an even function.

Proof. $\because f(x) = -f(-x)$

\therefore taking derivative on the both side, we can get

$$f'(x) = f'(-x)$$

which means $f'(x)$ is an even function.

□