

1. The-Real-and-Complex-Number-Systems
 - 1.1. Definition of Rational Numbers
 - 1.2. Rationals are Inadequate
 - 1.3. Order
 - 1.4. Ordered Set
 - 1.5. Upper Bounds and Lower Bounds
 - 1.6. Supremum and Infimum
 - 1.7. Least-Upper-Bound Property
 - 1.8. Relation between Supremum and Infimum

Note Information

- **ID:** 202501131947
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

Rational numbers are of the form $\frac{m}{n}$, where m and n are integers and $n \neq 0$.

Explanation

Review

1. Define the rational number system.

Links to Other Notes

-

Table of Contents

- TOC

Note Information

- **ID:** 202501132004
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-System
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

The rational number system is inadequate for many purposes, both as a field and as an ordered set.

Explanation

For example, there is no rational p such that $p^2 = 2$.

Proof 0.1 *Suppose on the contrary there was a p that satisfied $p^2 = 2$. We could write $p = \frac{m}{n}$, where m and n are integers and coprime. The original expression implies*

$$\begin{aligned}\left(\frac{m}{n}\right)^2 &= 2 \\ \frac{m^2}{n^2} &= 2 \\ m^2 &= 2n^2\end{aligned}$$

From this expression, we see that m^2 is even, and thus, m is even. Plugging $2k$ in for m , it is clear that m^2 is divisible by 4. It follows that $2n^2$ is divisible by 4 as well, which implies n^2 is even, and thus, n is even. Therefore, our assumption leads to a contradiction that both m and n are even, thus violating the coprime property of m and n . Hence, it is impossible for p to be rational.

Proof 0.2 (Alternative) *Let A be the set of all positive rationals p such that $p^2 < 2$ and let B consist of all positive rationals p such that $p^2 > 2$. By showing there is no largest element in A and no smallest element in B , we effectively partition the set of rational numbers, thus implying there is no rational p that falls outside these two sets, therefore satisfying $p^2 = 2$.*

To prove that for every p in A we can find a rational q in A such that $p < q$, we associate with each rational $p > 0$ the number

$$q = p - \frac{p^2 - 2}{p + 2} = \frac{2p + 2}{p + 2}. \quad (1)$$

Then

$$q^2 - 2 = \frac{2(p^2 - 2)}{(p + 2)^2}. \quad (2)$$

If p is in A then $p^2 - 2 < 0$, (1) shows that $q > p$, and (2) shows that $q^2 < 2$. Thus q is in A . If p is in B then $p^2 - 2 > 0$, (1) shows that $0 < q < p$, and (2) shows that $q^2 > 2$. Thus q is in B .

Review

1. Prove that there is no rational p such that $p^2 = 2$ in two different ways.

Links to Other Notes

- Definition of Rational Numbers

Table of Contents

- TOC

Note Information

- **ID:** 202501141228
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

Let S be a set. An order on S is a relation, denoted by $<$, with the following two properties:

- If $x \in S$ and $y \in S$ then one and only one of the statements

$$x < y, \quad x = y, \quad y < x$$

is true.

- If $x, y, z \in S$, if $x < y$ and $y < z$, then $x < z$.

Explanation

Review

1. Define order.

Links to Other Notes

-

Table of Contents

- TOC

Note Information

- **ID:** 202501141241
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

An ordered set is a set S in which an order is defined.

Explanation

For example, \mathbb{Q} is an ordered set if $r < s$ is defined to mean that $s - r$ is a positive rational number.

Review

1. Define ordered set and give an example.

Links to Other Notes

- Order

Table of Contents

- TOC

Note Information

- **ID:** 202501141250
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

Suppose S is an ordered set, and $E \subset S$. If there exists a $\beta \in S$ such that $x \leq \beta$ for every $x \in E$, we say that E is bounded above, and call β an upper bound of E . Lower bounds are defined in the same way.

Explanation

Review

1. Define upper bound.
2. Define lower bound.

Links to Other Notes

- Order
- Ordered Set

Table of Contents

- TOC

Note Information

- **ID:** 202501141546
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

Suppose S is an ordered set, $E \subset S$, and E is bounded above. Suppose there exists an $\alpha \in S$ with the following properties:

- α is an upper bound of E .
- If $\gamma < \alpha$ then γ is not an upper bound of E .

Then α is called the least upper bound of E or the supremum of E , and we write

$$\alpha = \sup E.$$

The greatest lower bound, or infimum, of a set E which is bounded below is defined in the same manner: If $\alpha = \inf E$, then α is a lower bound of E and no β with $\beta > \alpha$ is a lower bound of E .

Explanation

For example, consider the sets A and B from the alternative proof in Rationals are Inadequate. A and B are subsets of the ordered set Q . The set A is bounded above by the members of B . Since B contains no smallest member, A has no supremum in Q . B is bounded below by the members of A . Since A has no largest member, B has no infimum in Q .

Review

1. Define supremum.
2. Define infimum.

3. If $\alpha = \sup E$ exists, then must α be a member of E ? Give an example to justify your answer.
4. Let E consist of all numbers $1/n$, where $n = 1, 2, 3, \dots$. What is $\sup E$ and $\inf E$? Are $\sup E$ and $\inf E$ members of E ?

Links to Other Notes

- Definition of Rational Numbers
- Rationals are Inadequate
- Order
- Ordered Set
- Upper Bounds and Lower Bounds

Table of Contents

- TOC

Note Information

- **ID:** 202501141632
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

An ordered set S is said to have the least-upper-bound property if the following is true: If $E \subset S$, E is not empty, and E is bounded above, then $\sup E$ exists in S .

Explanation

Review

1. Define the least-upper-bound property.
2. Define the greatest-lower bound property.
3. Does \mathbb{Q} have the least-upper-bound property? Explain.

Links to Other Notes

- Definition of Rational Numbers
- Rationals are Inadequate
- Order
- Ordered Set
- Upper Bounds and Lower Bounds
- Supremum and Infimum

Table of Contents

- TOC

Note Information

- **ID:** 202501141654
- **Timestamp:** Wednesday 15th January, 2025 06:52
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
 - Analysis I
 - Rudin W., Principles of Mathematical Analysis

Main Content

Main Idea

If an ordered set has the least-upper-bound property, then it also has the greatest-lower-bound property.

Explanation

Theorem 0.3 *Suppose S is an ordered set with the least-upper-bound property, $B \subset S$, B is not empty, and B is bounded below. Let L be the set of all lower bounds of B . Then*

$$\alpha = \sup L$$

exists in S , and $\alpha = \inf B$. In particular, $\inf B$ exists in S .

Proof 0.4 *Since B is bounded below, L is not empty. Since L consists of exactly those $y \in S$ which satisfy the inequality $y \leq x$ for every $x \in B$, we see that every $x \in B$ is an upper bound of L . Thus L is bounded above. Our hypothesis about S implies therefore that L has a supremum in S , namely α . If $\gamma < \alpha$ then γ is not an upper bound of L , hence $\gamma \notin B$. It follows that $\alpha \leq x$ for every $x \in B$. Thus $\alpha \in L$. If $\alpha < \beta$ then $\beta \notin L$, since α is an upper bound of L . We have shown that $\alpha \in L$ but $\beta \notin L$ if $\beta > \alpha$. In other words α is a lower bound of B , but β is not if $\beta > \alpha$. This means that $\alpha = \inf B$.*

Review

1. Prove that if an ordered set has the least-upper-bound property, then it also has the greatest-lower-bound property.

Links to Other Notes

- Definition of Rational Numbers
- Rationals are Inadequate
- Order
- Ordered Set
- Upper Bounds and Lower Bounds
- Supremum and Infimum
- Least-Upper-Bound Property

Table of Contents

- TOC