

1. The-Real-and-Complex-Number-Systems
  - 1.1. Definition of Rational Numbers
  - 1.2. Rationals are Inadequate
  - 1.3. Order
  - 1.4. Ordered Set
  - 1.5. Upper Bounds and Lower Bounds

## Note Information

- **ID:** 202501131947
- **Timestamp:** Tuesday 14<sup>th</sup> January, 2025 12:57
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
  - Analysis I
  - Rudin W., Principles of Mathematical Analysis

## Main Content

### Main Idea

Rational numbers are of the form  $\frac{m}{n}$ , where  $m$  and  $n$  are integers and  $n \neq 0$ .

### Explanation

## Review

1. Define the rational number system.

## Links to Other Notes

- 

## Table of Contents

- TOC

## Note Information

- **ID:** 202501132004
- **Timestamp:** Tuesday 14<sup>th</sup> January, 2025 12:57
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-System
- **References:**
  - Analysis I
  - Rudin W., Principles of Mathematical Analysis

## Main Content

### Main Idea

The rational number system is inadequate for many purposes, both as a field and as an ordered set.

### Explanation

For example, there is no rational  $p$  such that  $p^2 = 2$ .

**Proof 0.1** *Suppose on the contrary there was a  $p$  that satisfied  $p^2 = 2$ . We could write  $p = \frac{m}{n}$ , where  $m$  and  $n$  are integers and coprime. The original expression implies*

$$\begin{aligned}\left(\frac{m}{n}\right)^2 &= 2 \\ \frac{m^2}{n^2} &= 2 \\ m^2 &= 2n^2\end{aligned}$$

*From this expression, we see that  $m^2$  is even, and thus,  $m$  is even. Plugging  $2k$  in for  $m$ , it is clear that  $m^2$  is divisible by 4. It follows that  $2n^2$  is divisible by 4 as well, which implies  $n^2$  is even, and thus,  $n$  is even. Therefore, our assumption leads to a contradiction that both  $m$  and  $n$  are even, thus violating the coprime property of  $m$  and  $n$ . Hence, it is impossible for  $p$  to be rational.*

**Proof 0.2 (Alternative)** *Let  $A$  be the set of all positive rationals  $p$  such that  $p^2 < 2$  and let  $B$  consist of all positive rationals  $p$  such that  $p^2 > 2$ . By showing there is no largest element in  $A$  and no smallest element in  $B$ , we effectively partition the set of rational numbers, thus implying there is no rational  $p$  that falls outside these two sets, therefore satisfying  $p^2 = 2$ .*

*To prove that for every  $p$  in  $A$  we can find a rational  $q$  in  $A$  such that  $p < q$ , we associate with each rational  $p > 0$  the number*

$$q = p - \frac{p^2 - 2}{p + 2} = \frac{2p + 2}{p + 2}. \quad (1)$$

*Then*

$$q^2 - 2 = \frac{2(p^2 - 2)}{(p + 2)^2}. \quad (2)$$

*If  $p$  is in  $A$  then  $p^2 - 2 < 0$ , (1) shows that  $q > p$ , and (2) shows that  $q^2 < 2$ . Thus  $q$  is in  $A$ . If  $p$  is in  $B$  then  $p^2 - 2 > 0$ , (1) shows that  $0 < q < p$ , and (2) shows that  $q^2 > 2$ . Thus  $q$  is in  $B$ .*

## Review

1. Prove that there is no rational  $p$  such that  $p^2 = 2$  in two different ways.

## Links to Other Notes

- Definition of Rational Numbers

## Table of Contents

- TOC

## Note Information

- **ID:** 202501141228
- **Timestamp:** Tuesday 14<sup>th</sup> January, 2025 12:57
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
  - Analysis I
  - Rudin W., Principles of Mathematical Analysis

## Main Content

### Main Idea

Let  $S$  be a set. An order on  $S$  is a relation, denoted by  $<$ , with the following two properties:

- If  $x \in S$  and  $y \in S$  then one and only one of the statements

$$x < y, \quad x = y, \quad y < x$$

is true.

- If  $x, y, z \in S$ , if  $x < y$  and  $y < z$ , then  $x < z$ .

### Explanation

## Review

1. Define order.

## Links to Other Notes

- 

## Table of Contents

- TOC

## Note Information

- **ID:** 202501141241
- **Timestamp:** Tuesday 14<sup>th</sup> January, 2025 12:57
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
  - Analysis I
  - Rudin W., Principles of Mathematical Analysis

## Main Content

### Main Idea

An ordered set is a set  $S$  in which an order is defined.

### Explanation

For example,  $\mathbb{Q}$  is an ordered set if  $r < s$  is defined to mean that  $s - r$  is a positive rational number.

## Review

1. Define ordered set and give an example.

## Links to Other Notes

- Order

## Table of Contents

- TOC

## Note Information

- **ID:** 202501141250
- **Timestamp:** Tuesday 14<sup>th</sup> January, 2025 12:57
- **Tags:** Mathematics, Analysis-I, The-Real-and-Complex-Number-Systems
- **References:**
  - Analysis I
  - Rudin W., Principles of Mathematical Analysis

## Main Content

### Main Idea

Suppose  $S$  is an ordered set, and  $E \subset S$ . If there exists a  $\beta \in S$  such that  $x \leq \beta$  for every  $x \in E$ , we say that  $E$  is bounded above, and call  $\beta$  an upper bound of  $E$ . Lower bounds are defined in the same way.

### Explanation

## Review

1. Define upper bound.
2. Define lower bound.

## Links to Other Notes

- Order
- Ordered Set

## Table of Contents

- TOC