

# Differential Equations Week **12**

Garud Shah

May 16, 2025

**Problem 1**

Solve the following differential equation:

$$\begin{cases} P' &= P(c - r(N)) \\ Q' &= Pr(N) \\ N &= P + Q \end{cases} \quad (1.1)$$

for

$$(a) \ r(N) = b(1 + \log N), \ P(0) = 1, \ Q(0) = 0, \ N(0) = 1$$

$$(b) \ r(N) = (2N - 1)s.$$

**Solution 1.1** — Note that  $Q(t)$  is very annoying, let's turn this into an  $P, N$  differential equation system.

$$\begin{cases} P' &= P(c - r(N)) \\ N' &= cP \end{cases}. \quad (1.2)$$

Now we solve the phase plane differential equation,

$$\frac{dP}{dN} = \frac{c - r(N)}{c} \quad (1.3)$$

$$P = N - \frac{1}{c} \int_{dontcare}^N r(x) dx + \alpha. \quad (1.4)$$

Now, plug back in:

$$N' = c \left( N - \frac{1}{c} \int_{dontcare}^N r(x) dx + \alpha \right) \quad (1.5)$$

$$t = \int \frac{1}{cN - \int_{dontcare}^N r(x) dx + c\alpha} dN \quad (1.6)$$

**Solution 1.2** — Plug in (set dontcare to 1):

$$t = \int \frac{1}{cN - bN \log N} dN \quad (1.7)$$

$$= \int \frac{1}{N} \cdot \frac{1}{c - b \log N} dN \quad (1.8)$$

$$= \int \frac{1}{c - bu} du \quad (1.9)$$

$$= -\frac{1}{b} \log |c - b \log |N|| + C \quad (1.10)$$

$$c - ae^{-bt} = b \log |N| \quad (1.11)$$

$$N = e^{\frac{1}{b}(c - ae^{-bt})} \quad (1.12)$$

$$P = e^{\frac{1}{b}(c - ae^{-bt})} - \frac{1}{c} N \log N \quad (1.13)$$

$$= e^{\frac{1}{b}(c - ae^{-bt})} \left( 1 - \frac{1}{bc} (c - ae^{-bt}) \right) \quad (1.14)$$