

Differential Equations Week **7**

Garud Shah

March 2, 2025

Can you please mark the equations you have problems with in the index of comments? I'm NOT asking for that to be the number, just when you make a comment, within the *text* of the comment (not the label) mark the bad equation (for all future weeks too, this is a clarification of last's week's!)

Problem 1 (Problem 1)

Solve the differential equation

$$y'' = ay, a \geq 0. \quad (1.1)$$

Solution — If $a = 0$ trivial, $ax + b$

We have characteristic equation:

$$\lambda^2 = a \quad (1.2)$$

$$\lambda = \pm\sqrt{a}, \quad (1.3)$$

thus the answer is:

$$c_1 e^{\sqrt{a}} + c_2 e^{-\sqrt{a}}. \quad (1.4)$$

Problem 2 (Problem 2)

Solve the following IVP's:

1. $y'' + 16y = 0, y(0) = 2, y'(0) = 0$
2. $y'' + 100y' + y = 0, y(0) = 1, y'(0) = 0.$
3. $y'' - 6y' + 8y = 0, y(0) = 1, y'(0) = 0.$

Solution 2.1 (Solution 2a) — $\omega = \sqrt{16}$, the answer is $2 \cos(4t)$.

Solution 2.2 (Solution 2b) — Roots of characteristic equation are $-50 \pm 7\sqrt{51}$, which gives us solutions $c_1 e^{(-50+7\sqrt{51})t} + c_2 e^{(-50-7\sqrt{51})t}$, so $c_1 + c_2 = 1$ and $c_1 - c_2 = 0$, thus answer is $\frac{1}{2} \left(e^{(-50+7\sqrt{51})t} + e^{(-50-7\sqrt{51})t} \right)$.

Solution 2.3 (Solution 2c) — We have roots 2 and 4 which means we have $c_1 e^{2t} + c_2 e^{4t}$, or $c_1 + c_2 = 1$ and $c_1 - c_2 = 0$ thus answer is $\frac{1}{2} (e^{2t} + e^{4t})$.

Problem 3 (Problem 3)

For the differential equation:

$$I\theta'' + b\theta' + k\theta = 0, \quad (3.1)$$

when will θ oscillate?

Solution 3.1 — $b^2 < 4Ik$ by criticality criteria...

Problem 4 (Problem 4)

Show that for

$$y'' + y = 0 \quad (4.1)$$

$$y(t) = c_1 \cos t + c_2 \sin t, \quad (4.2)$$

- (a) There is a unique solution that satisfies $y(0) = 2$ and $y(\pi/2) = 0$.
- (b) There is not a solution that satisfies $y(0) = 2$ and $y(\pi) = 0$.
- (c) There are infinite solutions that satisfy $y(0) = 2$ and $y(\pi) = -2$.

Solution 4.1 (Solution 4a) — Yes as the Wronskian for those two values is non-zero.

Solution 4.2 (Solution 4b) — As $y(0) = c_1$ and $y(\pi) = -c_1$ and $-2 \neq 0$ done.

Solution 4.3 (Solution 4c) — As $y(0) = c_1$ and $y(\pi) = -c_1$ and $-2 = 2$, y_1 can be anything done.

Problem 5 (Problem 5)

Solve the 2nd-Order Euler equation:

$$ax^2y'' + bxy' + cy = 0. \quad (5.1)$$

Solution 5.1 — Note that:

$$\frac{dy}{dx} = \frac{dy}{dv} \cdot \frac{dv}{dx}, \quad (5.2)$$

thus when substituting $v = \log x$, we need:

$$ay'' + by' + cy = 0. \quad (5.3)$$

Thus, if $ax^2 + bx + c$ has distinct roots r_1 and r_2 OR has a single root r ,

$$y(v) = c_1e^{r_1v} + c_2e^{r_2v} = c_1x^{r_1} + c_2x^{r_2} \quad (5.4)$$

$$OR \quad (5.5)$$

$$y(v) = c_1e^{rv} + c_2ve^{rv} = (c_1 + c_2 \log x)x^r. \quad (5.6)$$