

引力波天文学笔记

GasinAn

2022 年 6 月 17 日

Copyright © 2022 by GasinAn

All rights reserved. No part of this book may be reproduced, in any form or by any means, without permission in writing from the publisher, except by a BNUser.

The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories, technologies and programs to determine their effectiveness. The author and publisher make no warranty of any kind, express or implied, with regard to these techniques or programs contained in this book. The author and publisher shall not be liable in any event of incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these techniques or programs.

Printed in China

目录

第一章 引力波	5
1.1 Linearized Gravity . .	5
1.2 Radiation Gauge . . .	5
1.3 quadrupole Approxima- tion	5

第一章 引力波

[1].

1.1 Linearized Gravity

流形 \mathbb{R}^4 . 任意坐标系 $\{x^\mu\}$, $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} = \eta_{\mu\nu} + \gamma_{\mu\nu}s + O(s^2)$, 得

$$R_{\mu\nu\lambda\sigma} = \partial_\sigma \partial_{[\mu} h_{\lambda]\nu} - \partial_\nu \partial_{[\mu} h_{\lambda]\sigma} + O(s^2). \quad (1.1)$$

$$\bar{h}_{\mu\nu} := h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\eta^{\lambda\sigma}h_{\lambda\sigma} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h.$$

$$-\frac{1}{2}\partial^\lambda \partial_\lambda \bar{h}_{\mu\nu} + \partial^\lambda \partial_{(\mu} \bar{h}_{\nu)\lambda} - \frac{1}{2}\eta_{\mu\nu} \partial^\lambda \partial^\sigma \bar{h}_{\lambda\sigma} + O(s^2) = 8\pi T_{\mu\nu}. \quad (1.2)$$

存在 $\{x^\mu\}$, 使得 $\partial^\nu \bar{h}_{\mu\nu} + O(s^2) = 0$ (Lorentz gauge). 令 $\{x^\mu\}$ 满足 $\partial^\nu \bar{h}_{\mu\nu} + O(s^2) = 0$, 则

$$\partial^\lambda \partial_\lambda \bar{h}_{\mu\nu} + O(s^2) = -16\pi T_{\mu\nu}. \quad (1.3)$$

略去 $O(s^2)$ 条件: $h_{\mu\nu}, \partial_\lambda h_{\mu\nu} \dots$ 小.

1.2 Radiation Gauge

存在 $\{x^\mu\}$, 使得 $h + O(s^2) = 0$ (TT gauge [2]) 且 $h_{0\mu} + O(s^2) = 0$.

1.3 quadrupole Approximation

下略 $O(s^2)$. 由(1.3)得

$$\bar{h}_{\mu\nu}(t, \vec{r}) = 4 \int \frac{T_{\mu\nu}(t - |\vec{r} - \vec{r}'|, \vec{r}')}{|\vec{r} - \vec{r}'|} dV'. \quad (1.4)$$

$$\hat{h}_{\mu\nu}(\omega, \vec{r}) := \frac{1}{\sqrt{2\pi}} \int \bar{h}_{\mu\nu}(t, \vec{r}) e^{i\omega t} dt \quad (1.5)$$

$$= 4 \int \frac{\hat{T}_{\mu\nu}(\omega, \vec{r}')}{|\vec{r} - \vec{r}'|} e^{i\omega|\vec{r} - \vec{r}'|} dV'. \quad (1.6)$$

由 $\partial^\nu \bar{h}_{\mu\nu} = 0$,

$$-i\omega \hat{h}_{0\mu} = \sum_i \frac{\partial \hat{h}_{i\mu}}{\partial x^i}. \quad (1.7)$$

$|\vec{r}| \gg |\vec{r}'|$ 且 $\omega \ll 1/|\vec{r}'|$,

$$\hat{h}_{ij}(\omega, \vec{r}) = 4 \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{T}_{ij}(\omega, \vec{r}') dV'. \quad (1.8)$$

$$\int \hat{T}_{ij} dV' = -\frac{\omega^2}{2} \int \hat{T}_{00} x^i x^j dV', \quad ? \quad (1.9)$$

$$q_{ij}(t) := \int \hat{T}_{00} x^i x^j dV' \quad (1.10)$$

$$\hat{h}_{ij}(\omega, \vec{r}) = -2\omega^2 \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \hat{q}_{ij}(\omega), \quad (1.11)$$

$$\bar{h}_{ij}(t, \vec{r}) = \frac{2}{|\vec{r}|} \frac{d^2}{dt^2} q_{ij}(t - |\vec{r}|). \quad (1.12)$$

参考文献

- [1] Robert M. Wald. *General Relativity*. University of Chicago Pr., 1984.
- [2] 王运永. 引力波探测. 科学出版社, 2020.