

引力波天文学笔记

GasinAn

2022 年 11 月 23 日

目录

第一章 引力波	5
1.1 Linearized Gravity	5
1.2 Radiation Gauge	5
1.3 Quadrupole Approximation	5
1.4 + Mode and \times Mode	7
1.5 电磁—引力对比	8
1.5.1 电偶极—引力对比	9
1.5.2 电四极—引力对比	9
1.6 Varying G	10
第二章 能量	17
第三章 双星系统	19
3.1 基本公式	19
3.2 Post-Newtonian Approximation	19
3.3 Stationary Phase Approximation	19
第四章 宇宙学效应	21
第五章 电磁引力	23
5.1 时空张量转化为空间张量	23
5.2 电磁空间矢量	24
5.3 引力空间张量	25

第一章 引力波

1.1 Linearized Gravity

[6]. 流形 \mathbb{R}^4 . 任意坐标系 $\{x^\mu\}$, $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} = \eta_{\mu\nu} + \gamma_{\mu\nu}s + O(s^2)$, 得

$$R_{\mu\nu\lambda\sigma} = \partial_\sigma \partial_{[\mu} h_{\lambda]\nu} - \partial_\nu \partial_{[\mu} h_{\lambda]\sigma} + O(s^2). \quad (1.1)$$

$$\bar{h}_{\mu\nu} := h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\eta^{\lambda\sigma}h_{\lambda\sigma} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h.$$

$$-\frac{1}{2}\partial^\lambda \partial_\lambda \bar{h}_{\mu\nu} + \partial^\lambda \partial_{(\mu} \bar{h}_{\nu)\lambda} - \frac{1}{2}\eta_{\mu\nu} \partial^\lambda \partial^\sigma \bar{h}_{\lambda\sigma} + O(s^2) = 8\pi T_{\mu\nu}. \quad (1.2)$$

存在 $\{x^\mu\}$, 使得 $\partial^\nu \bar{h}_{\mu\nu} + O(s^2) = 0$ (Lorentz gauge). 令 $\{x^\mu\}$ 满足 $\partial^\nu \bar{h}_{\mu\nu} + O(s^2) = 0$, 则

$$\partial^\lambda \partial_\lambda \bar{h}_{\mu\nu} + O(s^2) = -16\pi T_{\mu\nu}. \quad (1.3)$$

略去 $O(s^2)$ 条件: $h_{\mu\nu}, \partial_\lambda h_{\mu\nu} \dots$ 小.

1.2 Radiation Gauge

[6]. 存在 $\{x^\mu\}$, 使得 “无源处” $h + O(s^2) = 0$ (TT gauge [7]) 且 $h_{0\mu} + O(s^2) = 0$.

1.3 Quadrupole Approximation

[6]. 下略 $O(s^2)$. 由(1.3)得

$$\bar{h}_{\mu\nu}(t, \vec{r}) = 4 \int \frac{T_{\mu\nu}(t - |\vec{r} - \vec{r}'|, \vec{r}')}{|\vec{r} - \vec{r}'|} dV'. \quad (1.4)$$

$$\hat{h}_{\mu\nu}(\omega, \vec{r}) := \frac{1}{\sqrt{2\pi}} \int \bar{h}_{\mu\nu}(t, \vec{r}) e^{i\omega t} dt \quad (1.5)$$

$$= 4 \int \frac{\hat{T}_{\mu\nu}(\omega, \vec{r}')}{|\vec{r} - \vec{r}'|} e^{i\omega|\vec{r} - \vec{r}'|} dV'. \quad (1.6)$$

由 $\partial^\nu \bar{h}_{\mu\nu} = 0$,

$$-i\omega \hat{h}_{0\mu} = \sum_i \frac{\partial \hat{h}_{i\mu}}{\partial x^i}. \quad (1.7)$$

$|\vec{r}| \gg |\vec{r}'|$ 且 $\omega \ll 1/|\vec{r}'|$,

$$\hat{h}_{ij}(\omega, \vec{r}) = 4 \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{T}_{ij}(\omega, \vec{r}') dV'. \quad (1.8)$$

$$\int \hat{T}_{ij} dV' = \int \sum_k (\hat{T}_{kj} \frac{\partial x'^i}{\partial x'^k}) dV' \quad (1.9)$$

$$= \sum_k \left[\int \frac{\partial}{\partial x'^k} (\hat{T}_{kj} x'^i) dV' - \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \right] \quad (1.10)$$

$$= \sum_k \int \partial'_k (\hat{T}_{kj} x'^i) dV' - \sum_k \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \quad (1.11)$$

$$= \int \hat{T}_{kj} x'^i dS' - \sum_k \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \quad (1.12)$$

$$= - \sum_k \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \quad (1.13)$$

$$= - \int (\sum_k \partial'_k \hat{T}_{kj}) x'^i dV' \quad (1.14)$$

$$= - \int (\partial_0 \hat{T}_{0j}) x'^i dV' \quad (1.15)$$

$$= -i\omega \int \hat{T}_{0j} x'^i dV' \quad (1.16)$$

$$= \int \hat{T}_{(ij)} dV' \quad (1.17)$$

$$= -i\omega \int \hat{T}_{0(j} x'^i) dV' \quad (1.18)$$

$$= -\frac{i\omega}{2} \int (\hat{T}_{0j} x'^i + \hat{T}_{0i} x'^j) dV', \quad (1.19)$$

$$(1.20)$$

$$-\frac{i\omega}{2} \int (\hat{T}_{0j} x'^i + \hat{T}_{0i} x'^j) dV' = -\frac{i\omega}{2} \int \sum_k (\hat{T}_{0k} x'^i \frac{\partial x'^j}{\partial x'^k} + \hat{T}_{0k} \frac{\partial x'^i}{\partial x'^k} x'^j) dV' \quad (1.21)$$

$$= -\frac{i\omega}{2} \sum_k \left[\int \frac{\partial}{\partial x'^k} (\hat{T}_{0k} x'^i x'^j) dV' - \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \right] \quad (1.22)$$

$$= -\frac{i\omega}{2} \sum_k \int \partial'_k (\hat{T}_{0k} x'^i x'^j) dV' + \frac{i\omega}{2} \sum_k \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \quad (1.23)$$

$$= -\frac{i\omega}{2} \sum_k \int \hat{T}_{0k} x'^i x'^j dS' + \frac{i\omega}{2} \sum_k \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \quad (1.24)$$

$$= \frac{i\omega}{2} \sum_k \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \quad (1.25)$$

$$= \frac{i\omega}{2} \int (\sum_k \partial'_k \hat{T}_{0k}) x'^i x'^j dV' \quad (1.26)$$

$$= \frac{i\omega}{2} \int (\partial_0 \hat{T}_{00}) x'^i x'^j dV' \quad (1.27)$$

$$= -\frac{\omega^2}{2} \int \hat{T}_{00} x'^i x'^j dV'. \quad (1.28)$$

$$q_{ij}(t) := \int T_{00} x'^i x'^j dV', \quad (1.29)$$

$$\hat{h}_{ij}(\omega, \vec{r}) = -2\omega^2 \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \hat{q}_{ij}(\omega), \quad (1.30)$$

$$\bar{h}_{ij}(t, \vec{r}) = \frac{2}{|\vec{r}|} \frac{d^2}{dt^2} q_{ij}(t - |\vec{r}|). \quad (1.31)$$

1.4 + Mode and \times Mode

寻新标架 $(e'^1)_a = (e^+)_a$, $(e'^2)_a = (e^\times)_a$, $(e'^3)_a = (e^r)_a$, $\bar{h}_{ij}(e^i)_a (e^j)_b = \bar{h}'_{ij}(e'^i)_a (e'^j)_b$, 取 x, y 分量后去迹, $h_+ = \frac{1}{2}(\bar{h}'_{11} - \bar{h}'_{22})$, $h_\times = \bar{h}'_{12} = \bar{h}'_{21}$? [5]
[2], $\vec{n} := \frac{\vec{r}}{|\vec{r}|}$,

$$h_{ij}^{\text{TT}} = \frac{2}{|\vec{r}|} \mathcal{P}_{ijkm} \frac{d^2}{dt^2} Q^{km}(t - |\vec{r}|), \quad (1.32)$$

$$\mathcal{P}_{ijkm} := (\delta_{ik} - \vec{n}_i \vec{n}_k) (\delta_{jm} - \vec{n}_j \vec{n}_m) - \frac{1}{2} (\delta_{ij} - \vec{n}_i \vec{n}_j) (\delta_{km} - \vec{n}_k \vec{n}_m), \quad (1.33)$$

$$Q^{km}(t) := \int T_{00} \left(x'^k x'^m - \frac{1}{3} \delta^{km} \sum_n x'^n x'^n \right) dV' \quad (1.34)$$

1.5 电磁—引力对比

$$A_\mu(t, \vec{r}) = \frac{\mu_0}{4\pi} \int \frac{J_\mu(t - |\vec{r} - \vec{r}'|, \vec{r}')}{|\vec{r} - \vec{r}'|} dV' \quad (1.35)$$

$$\bar{h}_{\mu\nu}(t, \vec{r}) = 4G \int \frac{T_{\mu\nu}(t - |\vec{r} - \vec{r}'|, \vec{r}')}{|\vec{r} - \vec{r}'|} dV' \quad (1.36)$$

$$A_\mu(t, \vec{r}) = \frac{1}{\sqrt{2\pi}} \int \hat{A}_\mu(\omega, \vec{r}) e^{-i\omega t} dt \quad (1.37)$$

$$\bar{h}_{\mu\nu}(t, \vec{r}) = \frac{1}{\sqrt{2\pi}} \int \hat{\bar{h}}_{\mu\nu}(\omega, \vec{r}) e^{-i\omega t} dt \quad (1.38)$$

$$\hat{A}_\mu(\omega, \vec{r}) = \frac{\mu_0}{4\pi} \int \frac{\hat{J}_\mu(\omega, \vec{r}')}{|\vec{r} - \vec{r}'|} e^{i\omega|\vec{r} - \vec{r}'|} dV' \quad (1.39)$$

$$\hat{\bar{h}}_{\mu\nu}(\omega, \vec{r}) = 4G \int \frac{\hat{T}_{\mu\nu}(\omega, \vec{r}')}{|\vec{r} - \vec{r}'|} e^{i\omega|\vec{r} - \vec{r}'|} dV' \quad (1.40)$$

$$\hat{A}_\mu(\omega, \vec{r}) = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{J}_\mu(\omega, \vec{r}') e^{-i\omega(\frac{\vec{r}}{|\vec{r}|} \cdot \vec{r}')} dV' \quad (1.41)$$

$$\hat{\bar{h}}_{\mu\nu}(\omega, \vec{r}) = 4G \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{T}_{\mu\nu}(\omega, \vec{r}') e^{-i\omega(\frac{\vec{r}}{|\vec{r}|} \cdot \vec{r}')} dV' \quad (1.42)$$

$$\hat{A}_\mu(\omega, \vec{r}) = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{J}_\mu(\omega, \vec{r}') \left[1 - i\omega \left(\frac{\vec{r}}{|\vec{r}|} \cdot \vec{r}' \right) - \dots \right] dV' \quad (1.43)$$

$$\hat{\bar{h}}_{\mu\nu}(\omega, \vec{r}) = 4G \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{T}_{\mu\nu}(\omega, \vec{r}') \left[1 - i\omega \left(\frac{\vec{r}}{|\vec{r}|} \cdot \vec{r}' \right) - \dots \right] dV' \quad (1.44)$$

1.5.1 电偶极—引力对比

$$\hat{A}_i = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{J}_i dV' \quad (1.45)$$

$$\hat{\bar{h}}_{ij} = 4G \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{T}_{ij} dV' \quad (1.46)$$

$$\int \hat{J}_i dV' = -i\omega \int \hat{J}_0 x'^i dV' \quad (1.47)$$

$$\int \hat{T}_{ij} dV' = -\frac{\omega^2}{2} \int \hat{T}_{00} x'^i x'^j dV' \quad (1.48)$$

$$\hat{p}_i = \int \hat{J}_0 x'^i dV' \quad (1.49)$$

$$\hat{q}_{ij} = \int \hat{T}_{00} x'^i x'^j dV' \quad (1.50)$$

$$\hat{A}_i = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} (-i\omega \hat{p}_i) \quad (1.51)$$

$$\hat{\bar{h}}_{ij} = 4G \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \left(-\frac{\omega^2}{2} \hat{q}_{ij}\right) \quad (1.52)$$

$$A_i = \frac{\mu_0}{4\pi} \frac{1}{|\vec{r}|} \frac{d}{dt} p_i(t - |\vec{r}|) \quad (1.53)$$

$$\bar{h}_{ij} = 4G \frac{1}{|\vec{r}|} \frac{1}{2} \frac{d^2}{dt^2} q_{ij}(t - |\vec{r}|) \quad (1.54)$$

1.5.2 电四极—引力对比

$$\hat{A}_i(\omega, \vec{r}) = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} (-i\omega) \int \hat{J}_i(\omega, \vec{r}') \left(\frac{\vec{r}}{|\vec{r}|} \cdot \vec{r}'\right) dV' \quad (1.55)$$

$$\hat{A}_i = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} (-i\omega) \int \hat{J}'_i n^j x'_j dV' \quad (1.56)$$

$$\hat{A}_i = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} (-i\omega) \int n^j x'_j \hat{J}'_i dV' \quad (1.57)$$

$$\hat{A}_i = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} (-i\omega) n^j \left[\int x'_{(j} \hat{J}'_{i)} dV' \right] \quad (1.58)$$

$$\int x'_{(j} \hat{J}'_{i)} dV' = \frac{1}{2} \int (\hat{J}'_j x'_i + \hat{J}'_i x'_j) dV' \quad (1.59)$$

$$= \frac{1}{2} \int \sum_k (\hat{J}'_k x'^i \frac{\partial x'^j}{\partial x'^k} + \hat{J}'_k \frac{\partial x'^i}{\partial x'^k} x'^j) dV' \quad (1.60)$$

$$= \frac{1}{2} \sum_k \left[\int \frac{\partial}{\partial x'^k} (\hat{J}'_k x'^i x'^j) dV' - \int \frac{\partial \hat{J}'_k}{\partial x'^k} x'^i x'^j dV' \right] \quad (1.61)$$

$$= \frac{1}{2} \sum_k \int \partial'_k (\hat{J}'_k x'^i x'^j) dV' - \frac{1}{2} \sum_k \int \frac{\partial \hat{J}'_k}{\partial x'^k} x'^i x'^j dV' \quad (1.62)$$

$$= \frac{1}{2} \sum_k \int \hat{J}'_k x'^i x'^j dS' - \frac{1}{2} \sum_k \int \frac{\partial \hat{J}'_k}{\partial x'^k} x'^i x'^j dV' \quad (1.63)$$

$$= -\frac{1}{2} \sum_k \int \frac{\partial \hat{J}'_k}{\partial x'^k} x'^i x'^j dV' \quad (1.64)$$

$$= -\frac{1}{2} \int (\sum_k \partial'_k \hat{J}'_k) x'^i x'^j dV' \quad (1.65)$$

$$= -\frac{1}{2} \int (\partial_0 \hat{J}'_0) x'^i x'^j dV' \quad (1.66)$$

$$= -\frac{i\omega}{2} \int \hat{J}'_0 x'^i x'^j dV' \quad (1.67)$$

$$\hat{D}_{ij} = \int \hat{J}'_0 x'^i x'^j dV' \quad (1.68)$$

$$\hat{A}_i = \frac{\mu_0}{4\pi} \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \left(-\frac{\omega^2}{2} n^j \hat{D}_{ij} \right) \quad (1.69)$$

$$A_i = \frac{\mu_0}{4\pi} \frac{1}{|\vec{r}|} n^j \frac{1}{2} \frac{d^2}{dt^2} D_{ij}(t - |\vec{r}|) \quad (1.70)$$

1.6 Varying G

$$T_{ab} = 2U_{(a} J_{b)} + U^c J_c U_a U_b \quad (1.71)$$

$$J_b = -U^a T_{ab} \quad (1.72)$$

$$\partial^a \bar{h}_{ab} = 0 \quad (1.73)$$

$$\partial^c \partial_c \bar{h}_{ab} = -16\pi \frac{G_0}{c_0^3} T_{ab} \quad (1.74)$$

$$\Gamma^c_{ab} = \frac{1}{2} \eta^{cd} (2\partial_{(a} h_{b)d} - \partial_d h_{ab}) \quad (1.75)$$

$$U^a \partial_a U^c + \Gamma^c_{ab} U^a U^b = 0 \quad (1.76)$$

$$U^a \partial_a U^c = -\frac{1}{2} \eta^{cd} (2\partial_{(a} h_{b)d} - \partial_d h_{ab}) U^a U^b \quad (1.77)$$

$$A_b = -\frac{1}{4} U^a \bar{h}_{ab} \quad (1.78)$$

$$A_0 = -\frac{1}{4} c_0 \bar{h}_{00} = -\frac{1}{2} c_0 (\bar{h}_{00} - \frac{1}{2} \eta_{00} \eta^{00} \bar{h}_{00}) = -\frac{1}{2} c_0 h_{00} \quad (1.79)$$

$$A_i = -\frac{1}{4} c_0 \bar{h}_{0i} = -\frac{1}{4} c_0 h_{0i} \quad (1.80)$$

$$U^\mu \partial_\mu U^i = -\frac{1}{2} \eta^{i\sigma} (\partial_\mu h_{\nu\sigma} + \partial_\nu h_{\mu\sigma} - \partial_\sigma h_{\mu\nu}) U^\mu U^\nu \quad (1.81)$$

$$-\frac{1}{2} \eta^{i\sigma} (\partial_0 h_{0\sigma} + \partial_0 h_{0\sigma} - \partial_\sigma h_{00}) U^0 U^0 = \frac{1}{2} c_0^2 \eta^{i\sigma} \partial_\sigma h_{00} \quad (1.82)$$

$$= \frac{1}{2} c_0^2 \partial^i h_{00} \quad (1.83)$$

$$= c_0 \partial^i A_0 \quad (1.84)$$

$$= -E^i \quad (1.85)$$

$$-\frac{1}{2} \eta^{i\sigma} (\partial_0 h_{j\sigma} + \partial_j h_{0\sigma} - \partial_\sigma h_{0j}) U^0 U^j = -\frac{1}{2} c_0 \eta^{i\sigma} (\partial_j h_{0\sigma} - \partial_\sigma h_{0j}) v^j \quad (1.86)$$

$$= -\frac{1}{2} c_0 \eta^{ik} (\partial_j h_{0k} - \partial_k h_{0j}) v^j \quad (1.87)$$

$$= 2\eta^{ik} (\partial_j A_k - \partial_k A_j) v^j \quad (1.88)$$

$$= 2(\partial_j A^i - \partial^i A_j) v^j \quad (1.89)$$

$$= -2\varepsilon^i_{jk} v^j B^k \quad (1.90)$$

$$-\frac{1}{2} \eta^{i\sigma} (\partial_j h_{k\sigma} + \partial_k h_{j\sigma} - \partial_\sigma h_{jk}) U^j U^k = 0 \quad (1.91)$$

$$a^i = -E^i - 4\varepsilon^i_{jk} v^j B^k \quad (1.92)$$

$$\partial^i \left(\frac{c_0}{16\pi G_0} E_i \right) = \rho \quad (1.93)$$

$$\partial^i B_i = 0 \quad (1.94)$$

$$\varepsilon^i_{jk} \partial^j E^k = -\partial_t B^i \quad (1.95)$$

$$\varepsilon^i_{jk} \partial^j \left(\frac{c_0^3}{16\pi G_0} B^k \right) = j^i + \partial_t \left(\frac{c_0}{16\pi G_0} E^i \right) \quad (1.96)$$

$$\vec{\nabla} \times \vec{E} = -\frac{\partial}{\partial t} (\mu \vec{H}) \quad (1.97)$$

$$\vec{\nabla} \times \vec{H} = +\frac{\partial}{\partial t} (\varepsilon \vec{E}) \quad (1.98)$$

$$\vec{\nabla} \cdot (\varepsilon \vec{E}) = 0 \quad (1.99)$$

$$\vec{\nabla} \cdot (\mu \vec{H}) = 0 \quad (1.100)$$

$$\vec{\nabla}^2 \vec{E} - \vec{\nabla} (\vec{\nabla} \cdot \vec{E}) - \varepsilon \mu \frac{\partial^2}{\partial t^2} \vec{E} = 0 \quad (1.101)$$

$$\vec{\nabla}^2 \vec{H} - \vec{\nabla} (\vec{\nabla} \cdot \vec{H}) - \varepsilon \mu \frac{\partial^2}{\partial t^2} \vec{H} = 0 \quad (1.102)$$

$$\varepsilon \vec{\nabla} \cdot \vec{E} + \vec{\nabla} \varepsilon \cdot \vec{E} = 0 \quad (1.103)$$

$$\mu \vec{\nabla} \cdot \vec{H} + \vec{\nabla} \mu \cdot \vec{H} = 0 \quad (1.104)$$

$$\vec{\nabla}^2 \vec{H} - \vec{\nabla} (\vec{\nabla} \cdot \vec{H}) - \frac{\partial^2}{\partial (v_p t)^2} \vec{H} = 0 \quad (1.105)$$

$$\mu \vec{\nabla} \cdot \vec{H} + \vec{\nabla} \mu \cdot \vec{H} = 0 \quad (1.106)$$

$$\vec{\nabla} \mu \cdot \vec{H} = 0 \quad (1.107)$$

$$\vec{\nabla}^2 \vec{H} - \frac{\partial^2}{\partial (v_p t)^2} \vec{H} = 0 \quad (1.108)$$

$$\vec{\nabla} \cdot \vec{H} = 0 \quad (1.109)$$

$$4\pi r_1^2 (\vec{E} \times \vec{H})|_{t=0, r=r_1} = 4\pi r_2^2 (\vec{E} \times \vec{H})|_{t=\int_{r_1}^{r_2} \frac{dr}{v_p}, r=r_1} \quad (1.110)$$

$$4\pi r_1^2 (\vec{S})|_{t=0, r=r_1} = 4\pi r_2^2 (\vec{S})|_{t=\int_{r_1}^{r_2} \frac{dr}{v_p}, r=r_1} \quad (1.111)$$

$$S_G \propto \dot{h}^2 \propto \omega^2 h^2 \quad (1.112)$$

$$S_G \propto \frac{c^3}{G} \omega^2 h^2 \quad (1.113)$$

双星系统引力辐射本为

$$h = \frac{\mathcal{M}[\pi \mathcal{M} F(t)]^{2/3}}{r} Q(\theta, \phi, \psi, \iota) \cos\left[\int 2\pi F(t) dt\right] \quad (1.114)$$

设双星系统常量 c^* , G^* , 一观者临近双星系统且与双星系统相对静止, 其与双星系统距离为 r , 测得强度 h_r , 频率 F_r , 则¹

$$h_r = \frac{\mathcal{M}[\pi \mathcal{M} F_r(t)]^{2/3}}{r/c^*} Q(\theta, \phi, \psi, \iota) \quad (1.115)$$

与双星系统距离为 r 的观者测得的引力辐射光度 $L_r \propto 4\pi r^2 (c^{*3}/G^*) F_r^2 h_r^2$, 设地球观者与双星系统距离为 d , 双星系统红移为 z , 测得强度 h_d , 频率 F_d , 则地球观者测得的引力辐射光度正比于 $L_d \propto 4\pi d^2 (c^3/G) F_d^2 h_d^2$, 且有 $L_d = L_r/(1+z)^2$, 所以 $r^2 (c^{*3}/G^*) F_r^2 h_r^2 / (1+z)^2 = d^2 (c^3/G) F_d^2 h_d^2$, 又有 $F_d = F_r/(1+z)$, 所以 $r^2 (c^{*3}/G^*) h_r^2 = d^2 (c^3/G) h_d^2$, 则

$$h_d = \sqrt{\frac{c^{*3}/G^*}{c^3/G} \frac{r^2}{d^2}} h_r \quad (1.116)$$

$$= \sqrt{\frac{c^{*3}/G^*}{c^3/G} \frac{\mathcal{M}[\pi \mathcal{M} F_r(t)]^{2/3}}{d/c^*}} Q(\theta, \phi, \psi, \iota) \quad (1.117)$$

所以地球观者测得

$$h = \sqrt{\frac{c^{*3}/G^*}{c^3/G} \frac{\mathcal{M}[\pi \mathcal{M} F_r(t)]^{2/3}}{d/c^*}} Q(\theta, \phi, \psi, \iota) \cos\left[\int 2\pi \frac{F_r(t)}{1+z} dt\right] \quad (1.118)$$

记 $F_{\text{obs}}(t) = F_r(t)/(1+z)$, $\mathcal{M}_{\text{obs}} = \mathcal{M}(1+z)$, 光度距离 $d_L = d(1+z)$, 则

$$h = \sqrt{\frac{c^{*3}/G^*}{c^3/G} \frac{\mathcal{M}[\pi \mathcal{M} F_r(t)]^{2/3}}{d(1+z)/c^*}} Q(\theta, \phi, \psi, \iota) \cos\left[\int 2\pi F_{\text{obs}}(t) dt\right] \quad (1.119)$$

$$= \sqrt{\frac{c^{*3}/G^*}{c^3/G} \frac{\mathcal{M}_{\text{obs}}[\pi \mathcal{M}_{\text{obs}} F_{\text{obs}}(t)]^{2/3}}{d_L/c^*}} Q(\theta, \phi, \psi, \iota) \cos\left[\int 2\pi F_{\text{obs}}(t) dt\right] \quad (1.120)$$

$$= \sqrt{\frac{c^{*5}/G^*}{c^5/G} \frac{\mathcal{M}_{\text{obs}}[\pi \mathcal{M}_{\text{obs}} F_{\text{obs}}(t)]^{2/3}}{d_L/c}} Q(\theta, \phi, \psi, \iota) \cos\left[\int 2\pi F_{\text{obs}}(t) dt\right] \quad (1.121)$$

¹ \mathcal{M} 和 c^* , G^* 简并, 所以可以笼统地仍记作 \mathcal{M} .

用引力波测距测得 $d_{L,G}$, 则

$$d_{L,G} = d_L \sqrt{\frac{c^5/G}{c^{*5}/G^*}} \quad (1.122)$$

[4]

$$h(t) = \frac{\mathcal{M}[\pi \mathcal{M} F(t)]^{2/3}}{\xi d_L} Q(\text{angles}) \cos \Phi(t) \quad (1.123)$$

$$\tilde{h}(f) = \frac{\sqrt{30}}{48\pi^{2/3}} \frac{\mathcal{M}^{5/6} Q}{\xi d_L} f^{-7/6} e^{i[2\pi f t(f) - \Phi(f) - \frac{\pi}{4}]} \quad (1.124)$$

问题转化为估计 ξ

$$p(\mu) \propto p^{(0)}(\mu) \exp[-\frac{1}{2} \Gamma_{ab}(\mu^a - \hat{\mu}^a)(\mu^b - \hat{\mu}^b)] \quad (1.125)$$

$$p^{(0)}(\mu) \propto \exp[-\frac{1}{2} \Gamma_{ab}^{(0)}(\mu^a - \bar{\mu}^a)(\mu^b - \bar{\mu}^b)] \quad (1.126)$$

设待估参数为 $\mu = (\ln \xi, \ln(d_L/d_{L0}), \ln Q, \dots), \dots$ 为其他参数 (如 \mathcal{M}), 则 $\tilde{h}_{,\ln \xi} = \tilde{h}_{,\ln(d_L/d_{L0})} = -\tilde{h}_{,\ln Q} = -\tilde{h}$, \tilde{h} 对其他参数求偏导皆为纯虚数, 则由 $\Gamma_{ab} = \langle h_{,a} | h_{,b} \rangle$ 和 $\text{SNR} := \rho = \sqrt{\langle h | h \rangle}$ 得

$$\Gamma_{ab} = \begin{bmatrix} \rho^2 & \rho^2 & -\rho^2 & 0 & \dots \\ \rho^2 & \rho^2 & -\rho^2 & 0 & \dots \\ -\rho^2 & -\rho^2 & \rho^2 & 0 & \dots \\ 0 & 0 & 0 & ? & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \quad (1.127)$$

又设

$$\Gamma_{ab}^{(0)} = \begin{bmatrix} 0 & 0 & 0 & 0 & \dots \\ 0 & 1/\sigma_{\ln d_L}^2 & 0 & 0 & \dots \\ 0 & 0 & 1/\sigma_{\ln Q}^2 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \quad (1.128)$$

则由 $\Sigma_{ab} = (\Gamma_{ab}^{(0)} + \Gamma_{ab})^{-1}$ 得

$$\Sigma_{ab} = \begin{bmatrix} \left[\begin{array}{ccc} \rho^2 & \rho^2 & -\rho^2 \\ \rho^2 & \rho^2 + 1/\sigma_{\ln(d_L/d_{L0})}^2 & -\rho^2 \\ -\rho^2 & -\rho^2 & \rho^2 + 1/\sigma_{\ln Q}^2 \end{array} \right]^{-1} & 0 \\ 0 & [?]^{-1} \end{bmatrix} \quad (1.129)$$

而

$$\begin{bmatrix} \rho^2 & \rho^2 & -\rho^2 \\ \rho^2 & \rho^2 + 1/\sigma_{\ln(d_L/d_{L0})}^2 & -\rho^2 \\ -\rho^2 & -\rho^2 & \rho^2 + 1/\sigma_{\ln Q}^2 \end{bmatrix}^{-1} \quad (1.130)$$

$$= \begin{bmatrix} 1/\rho^2 + \sigma_{\ln(d_L/d_{L0})}^2 + \sigma_{\ln Q}^2 & -\sigma_{\ln(d_L/d_{L0})}^2 & \sigma_{\ln Q}^2 \\ -\sigma_{\ln(d_L/d_{L0})}^2 & \sigma_{\ln(d_L/d_{L0})}^2 & 0 \\ \sigma_{\ln Q}^2 & 0 & \sigma_{\ln Q}^2 \end{bmatrix} \quad (1.131)$$

第二章 能量

第三章 双星系统

3.1 基本公式

$$\mathcal{M} := \mu^{3/5} M^{2/5} \quad (3.1)$$

$$h_+ = \frac{4\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} \frac{1 + \cos^2 \iota}{2} \cos \Phi(t) \quad (3.2)$$

$$h_\times = \frac{4\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} \cos \iota \sin \Phi(t) \quad (3.3)$$

$$h = F_+ h_+ + F_\times h_\times \quad (3.4)$$

3.2 Post-Newtonian Approximation

3.3 Stationary Phase Approximation

[4], if $\zeta(t)$ varies slowly near $t = t_0$ where the phase has a stationary point: $\phi'(t_0) = 0$,

$$\int \zeta(t) e^{i\phi(t;f)} dt = \int \zeta(t) e^{i[\phi(t_0) + \phi'(t_0)(t-t_0) + \frac{1}{2}\phi''(t_0)(t-t_0)^2 + \dots]} dt \quad (3.5)$$

$$\simeq e^{i\phi(t_0)} \int \zeta(t) e^{i[\frac{1}{2}\phi''(t_0)(t-t_0)^2]} dt \quad (3.6)$$

$$\simeq e^{i\phi(t_0)} \int \zeta(t_0) e^{\frac{-\sqrt{-i\phi''(t_0)}^2 (t-t_0)^2}{2}} dt \quad (3.7)$$

$$= \frac{\sqrt{2\pi}}{\sqrt{-i\phi''(t_0)}} \zeta(t_0) e^{i\phi(t_0)}. \quad (3.8)$$

$$h = \frac{\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} Q \cos \Phi(t) \quad (3.9)$$

$$= \frac{\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} Q \frac{1}{2} [e^{i\Phi(t)} + e^{-i\Phi(t)}] \quad (3.10)$$

$$\tilde{h}(f) = \int h(t) e^{i2\pi f t} dt \quad (3.11)$$

$$= \int \frac{\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} Q \frac{1}{2} [e^{i\Phi(t)} + e^{-i\Phi(t)}] e^{i2\pi f t} dt \quad (3.12)$$

$$= \int \frac{\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} Q \frac{1}{2} \{e^{i[2\pi f t + \Phi(t)]} + e^{i[2\pi f t - \Phi(t)]}\} dt \quad (3.13)$$

$$\simeq \int \frac{\mathcal{M}}{D} [\pi \mathcal{M} F(t)]^{2/3} Q \frac{1}{2} e^{i[2\pi f t - \Phi(t)]} dt \quad (3.14)$$

$$= \int \frac{\mathcal{M}}{D} [\pi \mathcal{M} F]^{2/3} Q \frac{1}{2} e^{i[2\pi f t(F) - \Phi(F)]} \frac{dt}{dF} dF \quad (3.15)$$

$$\simeq \frac{\sqrt{2\pi}}{\sqrt{-i[2\pi f t(F) - \Phi(F)]''_{F=f}}} \quad (3.16)$$

$$\left[\frac{\mathcal{M}}{D} (\pi \mathcal{M} F)^{2/3} Q \frac{1}{2} \frac{dt}{dF} \right]_{F=f} e^{i[2\pi f t(f) - \Phi(f)]} \quad (3.17)$$

$$\simeq \frac{\sqrt{2\pi}}{\sqrt{-i \left\{ 2\pi f \left[-\frac{5}{256} \mathcal{M} (\pi \mathcal{M} F)^{-8/3} \right] - \left[\frac{1}{16} (\pi \mathcal{M} F)^{-5/3} \right] \right\}''_{F=f}}} \quad (3.18)$$

$$\left\{ \frac{\mathcal{M}}{D} (\pi \mathcal{M} F)^{2/3} Q \frac{1}{2} \left[\frac{5\pi \mathcal{M}^2}{96} (\pi \mathcal{M} F)^{-11/3} \right] \right\}_{F=f} e^{i[2\pi f t(f) - \Phi(f)]} \quad (3.19)$$

$$= \frac{\sqrt{30}}{48\pi^{2/3}} \frac{\mathcal{M}^{5/6} Q}{D} f^{-7/6} e^{i[2\pi f t(f) - \Phi(f) - \frac{\pi}{4}]} \quad (\text{pnspace.py}) \quad (3.20)$$

或 [1], $h(t) = 2A(t) \cos \phi(t)$, $d \ln A / dt \ll d\phi / dt$ 且 $|d^2 \phi / dt^2| \ll (d\phi / dt)^2$.

第四章 宇宙学效应

第五章 电磁引力

[3].

5.1 时空张量转化为空间张量

$$h_{ab} := g_{ab} + Z_a Z_b. \quad (5.1)$$

$$h_a{}^b = \delta_a{}^b + Z_a Z^b. \quad (5.2)$$

$$Z^a h_{ab} = 0. \quad (5.3)$$

$$V_{\langle a} := h_a{}^b V_b. \quad (5.4)$$

$$Z^a V_{\langle a} = 0. \quad (5.5)$$

$$T_{\langle ab \rangle} := h_{(a}{}^c h_{b)}{}^d T_{cd} - \frac{1}{3} h_{cd} T^{cd} h_{ab}. \quad (5.6)$$

$$Z^a (h_a{}^c h_b{}^d T_{cd}) = 0. \quad (5.7)$$

$$Z^a (h_b{}^c h_a{}^d T_{cd}) = 0. \quad (5.8)$$

$$Z^a (h_{(a}{}^c h_{b)}{}^d T_{cd}) = 0. \quad (5.9)$$

$$Z^a (h_{cd} T^{cd} h_{ab}) = 0. \quad (5.10)$$

$$Z^a T_{\langle ab \rangle} = 0. \quad (5.11)$$

$$T_{\langle (ab) \rangle} = T_{\langle ab \rangle}. \quad (5.12)$$

$$h^{ab} T_{\langle ab \rangle} = 0. \quad (5.13)$$

$$\varepsilon_{abc} := \varepsilon_{abcd} Z^d. \quad (5.14)$$

$$\varepsilon_{0123} := -\sqrt{|g|}. \quad (5.15)$$

$$T_a := \frac{1}{2} \varepsilon_{abc} T^{[bc]}. \quad (5.16)$$

$$[U, V]_a := \varepsilon_{abc} U^b V^c. \quad (5.17)$$

$$[S, T]_a := \varepsilon_{abc} g_{de} S^{bd} T^{ce}. \quad (5.18)$$

$$D_t T^{a\dots}_{b\dots} := Z^c \nabla_c T^{a\dots}_{b\dots}. \quad (5.19)$$

$${}^3\nabla_a T^{b\dots}_{c\dots} := h_a{}^p h^b{}_q \dots h_c{}^r \dots \nabla_p T^{q\dots}_{r\dots}. \quad (5.20)$$

$$(\operatorname{div} V) := {}^3\nabla^a V_a. \quad (5.21)$$

$$(\operatorname{curl} V)_a := \varepsilon_{bca} {}^3\nabla^b V^c. \quad (5.22)$$

$$(\operatorname{div} T)_a := {}^3\nabla^b T_{ab}. \quad (5.23)$$

$$(\operatorname{curl} T)_{ab} := \varepsilon_{cd(a} {}^3\nabla^c g_{b)e} T^{ed}. \quad (5.24)$$

5.2 电磁空间矢量

$${}^*F_{ab} := \frac{1}{2} \varepsilon_{abcd} F^{cd} \quad (5.25)$$

$$E_a := F_{ab} Z^b = E_{\langle a \rangle}. \quad (5.26)$$

$$B_a := {}^*F_{ab} Z^b = B_{\langle a \rangle}. \quad (5.27)$$

$$\rho = -Z^a J_a. \quad (5.28)$$

$$j_a = h_a{}^b J_b. \quad (5.29)$$

$$\nabla_{[a} F_{bc]} = 0. \quad (5.30)$$

$$\nabla^a F_{ab} = \mu J_b. \quad (5.31)$$

$$(\operatorname{div} E) = \mu\rho - \dots \quad (5.32)$$

$$(\operatorname{div} B) = + \dots \quad (5.33)$$

$$(\operatorname{curl} E)_a + \dots = -D_t B_{\langle a} - \dots \quad (5.34)$$

$$(\operatorname{curl} B)_a + \dots = \mu j_a + D_t E_{\langle a} + \dots \quad (5.35)$$

5.3 引力空间张量

$${}^*C_{abcd} := \frac{1}{2}\varepsilon_{abef}C^ef_{cd}. \quad (5.36)$$

$$E_{ab} := C_{acbd}Z^cZ^d = E_{\langle ab \rangle}. \quad (5.37)$$

$$B_{ab} := {}^*C_{acbd}Z^cZ^d = B_{\langle ab \rangle}. \quad (5.38)$$

$$(\operatorname{div} E)_a = \kappa \frac{1}{3} \nabla_a \rho - \dots \quad (5.39)$$

$$(\operatorname{div} B)_a = \kappa(\rho + p)\omega_a + \dots \quad (5.40)$$

$$(\operatorname{curl} E)_{ab} + \dots = -D_t B_{\langle ab \rangle} - \dots \quad (5.41)$$

$$(\operatorname{curl} B)_{ab} + \dots = \kappa \frac{1}{2}(\rho + p)\sigma_{ab} + D_t E_{\langle ab \rangle} + \dots \quad (5.42)$$

参考文献

- [1] K. G. Arun, B. R. Iyer, B. S. Sathyaprakash, and P. A. Sundararajan. Parameter estimation of inspiralling compact binaries using 3.5 post-newtonian gravitational wave phasing: The nonspinning case. *Physical Review D*, 71(8):084008, Apr 2005.
- [2] L. Blanchet. Gravitational radiation from relativistic sources. In J.-A. Marck and J.-P. Lasota, editors, *Relativistic Gravitation and Gravitational Radiation*, page 33, Jan. 1997.
- [3] R. Maartens. Nonlinear gravito-electromagnetism. *General Relativity and Gravitation*, 40(6):1203–1217, June 2008.
- [4] E. Poisson and C. M. Will. Gravitational waves from inspiraling compact binaries: Parameter estimation using second-post-newtonian waveforms. *Physical Review D*, 52:848–855, Jul 1995.
- [5] B. S. Sathyaprakash and B. F. Schutz. Physics, astrophysics and cosmology with gravitational waves. *Living Reviews in Relativity*, 12(1):2, Dec. 2009.
- [6] R. M. Wald. *General Relativity*. University of Chicago Pr., 1984.
- [7] 王运永. 引力波探测. 科学出版社, 2020.