

# 引力波天文学笔记

GasinAn

2022 年 8 月 23 日

Copyright © 2022 by GasinAn

All rights reserved. No part of this book may be reproduced, in any form or by any means, without permission in writing from the publisher, except by a BNUser.

The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories, technologies and programs to determine their effectiveness. The author and publisher make no warranty of any kind, express or implied, with regard to these techniques or programs contained in this book. The author and publisher shall not be liable in any event of incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these techniques or programs.

Printed in China

# 目录

第一章	引力波	5
1.1	Linearized Gravity . .	5
1.2	Radiation Gauge . . .	5
1.3	Quadrupole Approximation . . . . .	5
1.4	+ Mode and $\times$ Mode .	7
第二章	电磁引力	9
2.1	时空张量转化为空间张量 . . . . .	9
2.2	电磁空间矢量 . . . . .	10
2.3	引力空间张量 . . . . .	11



# 第一章 引力波

## 1.1 Linearized Gravity

[4]. 流形  $\mathbb{R}^4$ . 任意坐标系  $\{x^\mu\}$ ,  $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} = \eta_{\mu\nu} + \gamma_{\mu\nu}s + O(s^2)$ , 得

$$R_{\mu\nu\lambda\sigma} = \partial_\sigma \partial_{[\mu} h_{\lambda]\nu} - \partial_\nu \partial_{[\mu} h_{\lambda]\sigma} + O(s^2). \quad (1.1)$$

$$\bar{h}_{\mu\nu} := h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\eta^{\lambda\sigma}h_{\lambda\sigma} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h.$$

$$-\frac{1}{2}\partial^\lambda \partial_\lambda \bar{h}_{\mu\nu} + \partial^\lambda \partial_{(\mu} \bar{h}_{\nu)\lambda} - \frac{1}{2}\eta_{\mu\nu} \partial^\lambda \partial^\sigma \bar{h}_{\lambda\sigma} + O(s^2) = 8\pi T_{\mu\nu}. \quad (1.2)$$

存在  $\{x^\mu\}$ , 使得  $\partial^\nu \bar{h}_{\mu\nu} + O(s^2) = 0$  (Lorentz gauge). 令  $\{x^\mu\}$  满足  $\partial^\nu \bar{h}_{\mu\nu} + O(s^2) = 0$ , 则

$$\partial^\lambda \partial_\lambda \bar{h}_{\mu\nu} + O(s^2) = -16\pi T_{\mu\nu}. \quad (1.3)$$

略去  $O(s^2)$  条件:  $h_{\mu\nu}, \partial_\lambda h_{\mu\nu} \dots$  小.

## 1.2 Radiation Gauge

[4]. 存在  $\{x^\mu\}$ , 使得  $h + O(s^2) = 0$  (TT gauge [5]) 且  $h_{0\mu} + O(s^2) = 0$ .

## 1.3 Quadrupole Approximation

[4]. 下略  $O(s^2)$ . 由(1.3)得

$$\bar{h}_{\mu\nu}(t, \vec{r}) = 4 \int \frac{T_{\mu\nu}(t - |\vec{r} - \vec{r}'|, \vec{r}')}{|\vec{r} - \vec{r}'|} dV'. \quad (1.4)$$

$$\hat{h}_{\mu\nu}(\omega, \vec{r}) := \frac{1}{\sqrt{2\pi}} \int \bar{h}_{\mu\nu}(t, \vec{r}) e^{i\omega t} dt \quad (1.5)$$

$$= 4 \int \frac{\hat{T}_{\mu\nu}(\omega, \vec{r}')}{|\vec{r} - \vec{r}'|} e^{i\omega|\vec{r} - \vec{r}'|} dV'. \quad (1.6)$$

由  $\partial^\nu \bar{h}_{\mu\nu} = 0$ ,

$$-i\omega \hat{h}_{0\mu} = \sum_i \frac{\partial \hat{h}_{i\mu}}{\partial x^i}. \quad (1.7)$$

$|\vec{r}| \gg |\vec{r}'|$  且  $\omega \ll 1/|\vec{r}'|$ ,

$$\hat{h}_{ij}(\omega, \vec{r}) = 4 \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \int \hat{T}_{ij}(\omega, \vec{r}') dV'. \quad (1.8)$$

$$\int \hat{T}_{ij} dV' = \int \sum_k (\hat{T}_{kj} \frac{\partial x'^i}{\partial x'^k}) dV' \quad (1.9)$$

$$= \sum_k \left[ \int \frac{\partial}{\partial x'^k} (\hat{T}_{kj} x'^i) dV' - \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \right] \quad (1.10)$$

$$= \sum_k \int \partial'_k (\hat{T}_{kj} x'^i) dV' - \sum_k \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \quad (1.11)$$

$$= \int \hat{T}_{kj} x'^i dS' - \sum_k \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \quad (1.12)$$

$$= - \sum_k \int \frac{\partial \hat{T}_{kj}}{\partial x'^k} x'^i dV' \quad (1.13)$$

$$= - \int (\sum_k \partial'_k \hat{T}_{kj}) x'^i dV' \quad (1.14)$$

$$= - \int (\partial_0 \hat{T}_{0j}) x'^i dV' \quad (1.15)$$

$$= -i\omega \int \hat{T}_{0j} x'^i dV' \quad (1.16)$$

$$= \int \hat{T}_{(ij)} dV' \quad (1.17)$$

$$= -i\omega \int \hat{T}_{0(j} x'^i) dV' \quad (1.18)$$

$$= -\frac{i\omega}{2} \int (\hat{T}_{0j} x'^i + \hat{T}_{0i} x'^j) dV', \quad (1.19)$$

$$(1.20)$$

$$-\frac{i\omega}{2} \int (\hat{T}_{0j} x'^i + \hat{T}_{0i} x'^j) dV' = -\frac{i\omega}{2} \int \sum_k (\hat{T}_{0k} x'^i \frac{\partial x'^j}{\partial x'^k} + \hat{T}_{0k} \frac{\partial x'^i}{\partial x'^k} x'^j) dV' \quad (1.21)$$

$$= -\frac{i\omega}{2} \sum_k \left[ \int \frac{\partial}{\partial x'^k} (\hat{T}_{0k} x'^i x'^j) dV' - \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \right] \quad (1.22)$$

$$= -\frac{i\omega}{2} \sum_k \int \partial'_k (\hat{T}_{0k} x'^i x'^j) dV' + \frac{i\omega}{2} \sum_k \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \quad (1.23)$$

$$= -\frac{i\omega}{2} \sum_k \int \hat{T}_{0k} x'^i x'^j dS' + \frac{i\omega}{2} \sum_k \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \quad (1.24)$$

$$= \frac{i\omega}{2} \sum_k \int \frac{\partial \hat{T}_{0k}}{\partial x'^k} x'^i x'^j dV' \quad (1.25)$$

$$= \frac{i\omega}{2} \int (\sum_k \partial'_k \hat{T}_{0k}) x'^i x'^j dV' \quad (1.26)$$

$$= \frac{i\omega}{2} \int (\partial_0 \hat{T}_{00}) x'^i x'^j dV' \quad (1.27)$$

$$= -\frac{\omega^2}{2} \int \hat{T}_{00} x'^i x'^j dV'. \quad (1.28)$$

$$q_{ij}(t) := \int T_{00} x'^i x'^j dV', \quad (1.29)$$

$$\hat{h}_{ij}(\omega, \vec{r}) = -2\omega^2 \frac{e^{i\omega|\vec{r}|}}{|\vec{r}|} \hat{q}_{ij}(\omega), \quad (1.30)$$

$$\bar{h}_{ij}(t, \vec{r}) = \frac{2}{|\vec{r}|} \frac{d^2}{dt^2} q_{ij}(t - |\vec{r}|). \quad (1.31)$$

## 1.4 + Mode and $\times$ Mode

寻新标架  $(e'^1)_a = (e^+)_a$ ,  $(e'^2)_a = (e^\times)_a$ ,  $(e'^3)_a = (e^r)_a$ ,  $\bar{h}_{ij}(e^i)_a (e^j)_b = \bar{h}'_{ij}(e'^i)_a (e'^j)_b$ , 取  $x, y$  分量后去迹,  $h_+ = \frac{1}{2}(\bar{h}'_{11} - \bar{h}'_{22})$ ,  $h_\times = \bar{h}'_{12} = \bar{h}'_{21}$ ? [3]  
[1],  $\vec{n} := \frac{\vec{r}}{|\vec{r}|}$ ,

$$h_{ij}^{\text{TT}} = \frac{2}{|\vec{r}|} \mathcal{P}_{ijkm} \frac{d^2}{dt^2} Q^{km}(t - |\vec{r}|), \quad (1.32)$$

$$\mathcal{P}_{ijkm} := (\delta_{ik} - \vec{n}_i \vec{n}_k) (\delta_{jm} - \vec{n}_j \vec{n}_m) - \frac{1}{2} (\delta_{ij} - \vec{n}_i \vec{n}_j) (\delta_{km} - \vec{n}_k \vec{n}_m), \quad (1.33)$$

$$Q^{km}(t) := \int T_{00} \left( x'^k x'^m - \frac{1}{3} \delta^{km} \sum_n x'^n x'^n \right) dV' \quad (1.34)$$



## 第二章 电磁引力

[2].

### 2.1 时空张量转化为空间张量

$$h_{ab} := g_{ab} + Z_a Z_b. \quad (2.1)$$

$$h_a{}^b = \delta_a{}^b + Z_a Z^b. \quad (2.2)$$

$$Z^a h_{ab} = 0. \quad (2.3)$$

$$V_{\langle a} := h_a{}^b V_b. \quad (2.4)$$

$$Z^a V_{\langle a} = 0. \quad (2.5)$$

$$T_{\langle ab \rangle} := h_{(a}{}^c h_{b)}{}^d T_{cd} - \frac{1}{3} h_{cd} T^{cd} h_{ab}. \quad (2.6)$$

$$Z^a (h_a{}^c h_b{}^d T_{cd}) = 0. \quad (2.7)$$

$$Z^a (h_b{}^c h_a{}^d T_{cd}) = 0. \quad (2.8)$$

$$Z^a (h_{(a}{}^c h_{b)}{}^d T_{cd}) = 0. \quad (2.9)$$

$$Z^a (h_{cd} T^{cd} h_{ab}) = 0. \quad (2.10)$$

$$Z^a T_{\langle ab \rangle} = 0. \quad (2.11)$$

$$T_{\langle (ab) \rangle} = T_{\langle ab \rangle}. \quad (2.12)$$

$$h^{ab} T_{\langle ab \rangle} = 0. \quad (2.13)$$

$$\varepsilon_{abc} := \varepsilon_{abcd} Z^d. \quad (2.14)$$

$$\varepsilon_{0123} := -\sqrt{|g|}. \quad (2.15)$$

$$T_a := \frac{1}{2} \varepsilon_{abc} T^{[bc]}. \quad (2.16)$$

$$[U, V]_a := \varepsilon_{abc} U^b V^c. \quad (2.17)$$

$$[S, T]_a := \varepsilon_{abc} g_{de} S^{bd} T^{ce}. \quad (2.18)$$

$$D_t T^{a\dots}_{b\dots} := Z^c \nabla_c T^{a\dots}_{b\dots}. \quad (2.19)$$

$${}^3\nabla_a T^{b\dots}_{c\dots} := h_a{}^p h^b{}_q \dots h_c{}^r \dots \nabla_p T^{q\dots}_{r\dots}. \quad (2.20)$$

$$(\operatorname{div} V) := {}^3\nabla^a V_a. \quad (2.21)$$

$$(\operatorname{curl} V)_a := \varepsilon_{bca} {}^3\nabla^b V^c. \quad (2.22)$$

$$(\operatorname{div} T)_a := {}^3\nabla^b T_{ab}. \quad (2.23)$$

$$(\operatorname{curl} T)_{ab} := \varepsilon_{cd(a} {}^3\nabla^c g_{b)e} T^{ed}. \quad (2.24)$$

## 2.2 电磁空间矢量

$${}^*F_{ab} := \frac{1}{2} \varepsilon_{abcd} F^{cd} \quad (2.25)$$

$$E_a := F_{ab} Z^b = E_{\langle a \rangle}. \quad (2.26)$$

$$B_a := {}^*F_{ab} Z^b = B_{\langle a \rangle}. \quad (2.27)$$

$$\rho = -Z^a J_a. \quad (2.28)$$

$$j_a = h_a{}^b J_b. \quad (2.29)$$

$$\nabla_{[a} F_{bc]} = 0. \quad (2.30)$$

$$\nabla^a F_{ab} = \mu J_b. \quad (2.31)$$

$$(\operatorname{div} E) = \mu\rho - \dots \quad (2.32)$$

$$(\operatorname{div} B) = + \dots \quad (2.33)$$

$$(\operatorname{curl} E)_a + \dots = -D_t B_{\langle a} - \dots \quad (2.34)$$

$$(\operatorname{curl} B)_a + \dots = \mu j_a + D_t E_{\langle a} + \dots \quad (2.35)$$

## 2.3 引力空间张量

$${}^*C_{abcd} := \frac{1}{2}\varepsilon_{abef}C^ef_{cd}. \quad (2.36)$$

$$E_{ab} := C_{acbd}Z^cZ^d = E_{\langle ab \rangle}. \quad (2.37)$$

$$B_{ab} := {}^*C_{acbd}Z^cZ^d = B_{\langle ab \rangle}. \quad (2.38)$$

$$(\operatorname{div} E)_a = \kappa \frac{1}{3} \nabla_a \rho - \dots \quad (2.39)$$

$$(\operatorname{div} B)_a = \kappa(\rho + p)\omega_a + \dots \quad (2.40)$$

$$(\operatorname{curl} E)_{ab} + \dots = -D_t B_{\langle ab \rangle} - \dots \quad (2.41)$$

$$(\operatorname{curl} B)_{ab} + \dots = \kappa \frac{1}{2}(\rho + p)\sigma_{ab} + D_t E_{\langle ab \rangle} + \dots \quad (2.42)$$



## 参考文献

- [1] L. Blanchet. Gravitational radiation from relativistic sources. In J.-A. Marck and J.-P. Lasota, editors, *Relativistic Gravitation and Gravitational Radiation*, page 33, Jan. 1997.
- [2] R. Maartens. Nonlinear gravito-electromagnetism. *General Relativity and Gravitation*, 40(6):1203–1217, June 2008.
- [3] B. S. Sathyaprakash and B. F. Schutz. Physics, astrophysics and cosmology with gravitational waves. *Living Reviews in Relativity*, 12(1):2, Dec. 2009.
- [4] R. M. Wald. *General Relativity*. University of Chicago Pr., 1984.
- [5] 王运永. 引力波探测. 科学出版社, 2020.