

Probe Varying Gravitational Constant via GW

Bing Sun¹ and Jiachen An^{2,3}

¹*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

²*Institute for Frontiers in Astronomy and Astrophysics,
Beijing Normal University, Beijing 102206, China*

³*Department of Astronomy, Beijing Normal University, Beijing 102206, China*

(Dated: October 30, 2023)

Abstract

In this note, we give the effect of varying gravitational constant G on gravitational waves (GWs) during propagations. We separately make G in Fierz-Pauli action and Einstein-Hilbert action vary, and then we maintain two modified GW equations with the same form. We then solve the equations for the situation in our interest, and our solution can be easily applied to gravitational constant in Yukawa form when it has a minor correction and a slow variation.

I. ACTION METHOD FOR FIERZ-PAULI ACTION

The Fierz-Pauli action for the field variable h_{ab} is given by [1]

$$S = \frac{1}{64\pi G} \int d^4x \left(-\partial_a h_{bc} \partial^a h^{bc} + \partial_a h_b^b \partial^a h_c^c - 2\partial_a h_c^a \partial^c h_b^b + 2\partial_a h_c^a \partial_b h^{bc} \right). \quad (1)$$

We set gravitational constant G as a function in a form ϕ^{-1} so that the action becomes

$$S = \frac{1}{64\pi} \int d^4x \phi \left(-\partial_a h_{bc} \partial^a h^{bc} + \partial_a h_b^b \partial^a h_c^c - 2\partial_a h_c^a \partial^c h_b^b + 2\partial_a h_c^a \partial_b h^{bc} \right). \quad (2)$$

In the Lorentz gauge of transverse trace variable $\bar{h}_{ab} = h_{ab} - \frac{1}{2}\eta_{ab}h$, $\partial^b \bar{h}_{ab} = 0$, the action (2) has a simple form

$$S = \frac{1}{64\pi} \int d^4x \phi \left(-\partial_a \bar{h}_{bc} \partial^a \bar{h}^{bc} + \frac{1}{2} \partial_c \bar{h} \partial^c \bar{h} \right). \quad (3)$$

The equation of motion (EOM) for this action becomes

$$\square h_{ab} + \partial_c h_{ab} \partial^c \ln \phi = 0. \quad (4)$$

where $h_{ab} = \bar{h}_{ab} - \frac{1}{2}\eta_{ab}\bar{h}$. We separate the variable as $h_{ab} = H_{ab}e^{-i\omega t}$, and then

$$\nabla^2 H_{ab} + \omega^2 H_{ab} + i\omega \partial_t \ln \phi H_{ab} + \delta^{ij} \partial_j \ln \phi \partial_i H_{ab} = 0. \quad (5)$$

If the GW travels along z -axis in a transverse-traceless (TT) gauge, $H_{ij} = H_{ij}(z)$ and the equation becomes

$$\partial_z^2 H_{ij} + \omega^2 H_{ij} + i\omega \partial_t \ln \phi H_{ij} + \partial_z \ln \phi \partial_z H_{ij} = 0. \quad (6)$$

Moreover, if $\phi = \phi(\vec{r})$, the expression eq.(6) is solvable

$$\partial_z^2 H_{ij} + \partial_z \ln \phi \partial_z H_{ij} + \omega^2 H_{ij} = 0. \quad (7)$$

In addition, if $\phi \neq \phi(\vec{r})$ but $\phi = \phi(t)$, the new variable $h_{ij} = C_{ij}e^{ikz}H(t)$ obeys the relation from eq.(4), and then

$$\partial_t^2 H + \partial_t \ln \phi \partial_t H + k^2 H = 0, \quad (8)$$

which has a similar form like eq.(7).

Obviously, eq.(7) has the form like eq.(A1) in Appendix A. Now we use definitions and methods in Appendix A. For eq.(7),

$$p = \frac{1}{2} \partial_z \ln \phi, \quad q = 0, \quad (9)$$

therefore $\Gamma \propto \phi^{1/2} \propto G^{-1/2}$ and $\Xi = -\frac{1}{4}\phi^{-2}(\partial_z \phi)^2 + \frac{1}{2}\phi^{-1}\partial_z^2 \phi = \frac{3}{4}G^{-2}(\partial_z G)^2 - \frac{1}{2}G^{-1}\partial_z^2 G$. If G takes the Yukawa form [2]

$$G = G_\infty[1 + \alpha(1 + z/\lambda)e^{-z/\lambda}], \quad (10)$$

there will be

$$\Xi = \frac{1}{\lambda^2} \frac{[\alpha(z/\lambda)e^{-z/\lambda}]^2 + 2[\alpha e^{-z/\lambda}]^2 - 2[\alpha(z/\lambda)e^{-z/\lambda}] + 2[\alpha e^{-z/\lambda}]}{4[\alpha(z/\lambda + 1)e^{-z/\lambda} + 1]^2}. \quad (11)$$

Moreover, if $|\frac{\Xi}{\omega^2}| \ll 1$ and $1/\lambda \ll \omega/c$, which means that G has minor correction and slow variation, there will be $K \approx 1 + \frac{1}{4}\frac{\Xi}{\omega^2} \approx 1$ and $\omega \int K^{-2} dz \approx \omega \int (1 + \frac{1}{4}\frac{\Xi}{\omega^2})^{-2} dz \approx \omega \int (1 - \frac{1}{2}\frac{\Xi}{\omega^2}) dz$. Consequently, the amplitude of GW, which is proportional to $\Gamma^{-1}K$, will be approximately proportional to $G^{1/2}$, and the phase of GW, $\omega \int K^{-2} dz$, will have an approximate modification $-\omega \int \frac{1}{2}\frac{\Xi}{\omega^2} dz$.

II. ACTION METHOD FOR EINSTEIN-HILBERT ACTION

Using the *xAct* module in Mathematica, we expand the Lagrangian of Einstein-Hilbert action up to second order

$$\begin{aligned} \mathcal{L}_{\text{GR}} = \sqrt{-g}R = & -\frac{1}{2}\partial_b h \partial^b h - 2\partial_a h^{ab} \partial_c h_b^c + 2\partial_b h \partial_c h^{bc} + h \partial_b \partial_c h^{bc} \\ & + 2h^{ab}(\partial_a \partial_b h - \partial_b \partial_c h_a^c - \partial_c \partial_b h_a^c + \square h_{ab}) \\ & - h \square h - \partial_b h_{ac} \partial^c h^{ab} + \frac{3}{2} \partial_c h_{ab} \partial^c h^{ab}. \end{aligned} \quad (12)$$

Using the transverse trace variable

$$\bar{h}_{ab} = h_{ab} - \frac{1}{2}\eta_{ab}h, \quad h_{ab} = \bar{h}_{ab} - \frac{1}{2}\eta_{ab}\bar{h}, \quad (13)$$

and the gauge condition

$$\partial_a \bar{h}^{ab} = 0, \quad \partial_a h^{ab} = \frac{1}{2} \partial^b h = -\frac{1}{2} \partial^b \bar{h}, \quad (14)$$

it becomes

$$\begin{aligned} \sqrt{-g}R &= -\frac{1}{2} \partial_b h \partial^b h - 2 \partial_a h^{ab} \partial_c h_b^c + 2 \partial_b h \partial_c h^{bc} + h \partial_b \partial_c h^{bc} \\ &\quad + 2 h^{ab} (\partial_a \partial_b h - \partial_b \partial_c h_a^c - \partial_c \partial_b h_a^c + \square h_{ab}) \\ &\quad - h \square h - \partial_b h_{ac} \partial^c h^{ab} + \frac{3}{2} \partial_c h_{ab} \partial^c h^{ab} \\ &= -\frac{1}{2} \partial_b h \partial^b h - \frac{1}{2} \partial^b h \partial_b h + \partial_b h \partial^b h + \frac{1}{2} h \square h \\ &\quad + 2 h^{ab} \partial_a \partial_b h - 2 h^{ab} \partial_b \partial_a h + h^{ab} \square h_{ab} - h \square h \\ &\quad - \partial_b h_{ac} \partial^c h^{ab} + \frac{3}{2} \partial_c h_{ab} \partial^c h^{ab} \\ &= -\frac{1}{2} h \square h + h^{ab} \square h_{ab} - \partial_b h_{ac} \partial^c h^{ab} + \frac{3}{2} \partial_c h_{ab} \partial^c h^{ab} \\ &= -\frac{1}{2} \bar{h} \square \bar{h} + \bar{h}^{ab} \square \bar{h}_{ab} - \partial_b \bar{h}_{ac} \partial^c \bar{h}^{ab} - \frac{1}{4} \partial_b \bar{h} \partial^b \bar{h} + \frac{3}{2} \partial_c \bar{h}_{ab} \partial^c \bar{h}^{ab}, \end{aligned} \quad (15)$$

where we used the relations

$$\begin{aligned} -h^{ab} \square h_{ab} &= 2(\bar{h}^{ab} - \frac{1}{2} \eta^{ab} \bar{h}) \square (\bar{h}_{ab} - \frac{1}{2} \eta_{ab} \bar{h}) \\ &= 2 \bar{h}^{ab} \square \bar{h}_{ab} - 2 \bar{h} \square \bar{h} + 2 \bar{h} \square \bar{h} \\ &= \bar{h}^{ab} \square \bar{h}_{ab}, \end{aligned} \quad (16)$$

$$h \square h = -\bar{h} \square \bar{h}, \quad (17)$$

$$\begin{aligned} -\partial_b h_{ac} \partial^c h^{ab} &= -\partial_b (\bar{h}_{ac} - \frac{1}{2} \eta_{ac} \bar{h}) \partial^c (\bar{h}^{ab} - \frac{1}{2} \eta^{ab} \bar{h}) \\ &= -\partial_b \bar{h}_{ac} \partial^c \bar{h}^{ab} + \partial_b \bar{h} \partial_a \bar{h}^{ab} - \frac{1}{4} \partial_b \bar{h} \partial^b \bar{h} \\ &= -\partial_b \bar{h}_{ac} \partial^c \bar{h}^{ab} - \frac{1}{4} \partial_b \bar{h} \partial^b \bar{h}, \end{aligned} \quad (18)$$

$$\begin{aligned} \frac{3}{2} \partial_c h_{ab} \partial^c h^{ab} &= \frac{3}{2} \partial_c (\bar{h}_{ab} - \frac{1}{2} \eta_{ab} \bar{h}) \partial^c (\bar{h}^{ab} - \frac{1}{2} \eta^{ab} \bar{h}) \\ &= \frac{3}{2} \partial_c \bar{h}_{ab} \partial^c \bar{h}^{ab} - \frac{3}{2} \partial_c \bar{h} \partial^c \bar{h} + \frac{3}{2} \partial_c h \partial^c \bar{h} \\ &= \frac{3}{2} \partial_c \bar{h}_{ab} \partial^c \bar{h}^{ab}. \end{aligned} \quad (19)$$

Similarly the action for the GW is

$$\begin{aligned} S &= \frac{1}{16\pi G} \int d^4x \left[-\frac{1}{2} \bar{h} \square \bar{h} + \bar{h}^{ab} \square \bar{h}_{ab} - \partial_b \bar{h}_{ac} \partial^c \bar{h}^{ab} - \frac{1}{4} \partial_b \bar{h} \partial^b \bar{h} + \frac{3}{2} \partial_c \bar{h}_{ab} \partial^c \bar{h}^{ab} \right] \\ &= \frac{1}{16\pi} \int d^4x \phi \left[-\frac{1}{2} \bar{h} \square \bar{h} + \bar{h}^{ab} \square \bar{h}_{ab} - \partial_b \bar{h}_{ac} \partial^c \bar{h}^{ab} - \frac{1}{4} \partial_b \bar{h} \partial^b \bar{h} + \frac{3}{2} \partial_c \bar{h}_{ab} \partial^c \bar{h}^{ab} \right]. \end{aligned} \quad (20)$$

We work in TT gauge,

$$S = \frac{1}{16\pi} \int d^4x \phi \left[\bar{h}^{ij} \square \bar{h}_{ij} - \partial_j \bar{h}_{ik} \partial^k \bar{h}^{ij} + \frac{3}{2} \partial_c \bar{h}_{ij} \partial^c \bar{h}^{ij} \right], \quad (21)$$

and EOM becomes

$$\phi \square \bar{h}_{ij} + \square(\phi \bar{h}_{ij}) + 2\partial^k (\phi \partial_j \bar{h}_{ik}) - \frac{3}{2} \partial^a (\phi \partial_a \bar{h}_{ij}) = 0, \quad (22)$$

or

$$\square \bar{h}_{ij} + (2\phi^{-1} \square \phi) \bar{h}_{ij} + \partial^a \ln \phi \partial_a \bar{h}_{ij} + 4\partial^k \ln \phi \partial_j \bar{h}_{ik} = 0. \quad (23)$$

Assuming $\phi = \phi(\vec{r})$, we obtain the differential equations of \bar{h}_{ij}

$$\square \bar{h}_{ij} + \left(\frac{2}{\phi} \nabla^2 \phi \right) \bar{h}_{ij} + \partial^k \ln \phi \partial_k \bar{h}_{ij} + 4\partial^k \ln \phi \partial_j \bar{h}_{ik} = 0. \quad (24)$$

We then separate the variables as $\bar{h}_{ij} = H_{ij} e^{-i\omega t}$ so the spatial differential equations become

$$\nabla^2 H_{ij} + \left[\omega^2 + \frac{2}{\phi} \nabla^2 \phi \right] H_{ij} + \partial^k \ln \phi \partial_k H_{ij} + 4\partial^k \ln \phi \partial_j H_{ik} = 0. \quad (25)$$

Because the GW travels along z -axis, the H_{ij} is the function of z only,

$$\partial_z^2 H_{ij} + \left[\omega^2 + \frac{2}{\phi} \partial_z^2 \phi \right] H_{ij} + \partial_z \ln \phi \partial_z H_{ij} + 4\delta_j^z \partial^k \ln \phi \partial_z H_{ik} = 0. \quad (26)$$

In TT gauge, H_{ij} furthermore only has x, y components so the equations become

$$\partial_z^2 h + \partial_z \ln \phi \partial_z h + (\omega^2 + 2\phi^{-1} \partial_z^2 \phi) h = 0, \quad (27)$$

where h represents h_+ or h_\times , and approximately this system has a frequency shift $\frac{\phi^{-1} \partial_z^2 \phi}{\omega^2} \omega$.

In addition, if $\phi \neq \phi(\vec{r})$ but $\phi = \phi(t)$, the EOM becomes

$$\frac{1}{2} \square \bar{h}_{ij} + \frac{\bar{h}_{ij}}{\phi} \square \phi + \frac{1}{2} \partial^a \ln \phi \partial_a \bar{h}_{ij} + 2\partial^k \ln \phi \partial_j \bar{h}_{ik} = 0, \quad (28)$$

or after variable separation $\bar{h}_{ij} = C_{ij} e^{ikz} H(t)$,

$$\partial_t^2 H + \partial_t \ln \phi \partial_t H + (k^2 + 2\phi^{-1} \partial_t^2 \phi) H = 0, \quad (29)$$

which has a similar form like eq.(27).

Again, obviously eq.(27) has the form like eq.(A1) and now we use definitions and methods in Appendix A. For eq.(27),

$$p = \frac{1}{2} \partial_z \ln \phi, \quad q = 2\phi^{-1} \partial_z^2 \phi, \quad (30)$$

therefore $\Gamma \propto \phi^{1/2} \propto G^{-1/2}$ and $\Xi = -\frac{1}{4}\phi^{-2}(\partial_z \phi)^2 - \frac{3}{2}\phi^{-1}\partial_z^2 \phi = -\frac{13}{4}G^{-2}(\partial_z G)^2 + \frac{3}{2}G^{-1}\partial_z^2 G$.

If G takes the Yukawa form eq.(10), there will be

$$\Xi = \frac{1}{\lambda^2} \frac{-7[\alpha(z/\lambda)e^{-z/\lambda}]^2 - 6[\alpha e^{-z/\lambda}]^2 + 6[\alpha(z/\lambda)e^{-z/\lambda}] - 6[\alpha e^{-z/\lambda}]}{4[\alpha(z/\lambda + 1)e^{-z/\lambda} + 1]^2}. \quad (31)$$

Again, if $|\frac{\Xi}{\omega^2}| \ll 1$ and $1/\lambda \ll \omega/c$, which means that G has minor correction and slow variation, the amplitude of GW will be approximately proportional to $G^{1/2}$ and the phase of GW will have an approximate modification $-\omega \int \frac{1}{2} \frac{\Xi}{\omega^2} dz$.

Appendix A: Solving Wave Equation

In this appendix, we solve the wave equation. By expressing h_{ij} as a Fourier transform, the wave equation for each Fourier amplitude becomes the following general linear form,

$$\frac{d^2}{dz^2} H(z) + 2p(z) \frac{d}{dz} H(z) + [\omega^2 + q(z)] H(z) = 0. \quad (A1)$$

For any $H(z)$, there are real function $A(z)$ and $\Phi(z)$ which satisfy

$$H = A e^{i\Phi}. \quad (A2)$$

We plug eq.(A2) into eq.(A1) and then divide both sides of the equation by $e^{i\Phi}$. From the real and imaginary part of the result, we achieve two equations,

$$\frac{d^2 A}{dz^2} + 2p \frac{dA}{dz} + \left[\omega^2 \left(1 - \frac{k^2}{\omega^2} \right) + q \right] A = 0, \quad (A3)$$

$$2 \frac{dA}{dz} k + A \frac{dk}{dz} + 2p A k = 0, \quad (A4)$$

where $k = \frac{d\Phi}{dz}$. It is no matter for us to suppose that $A > 0$ and $k > 0$, and then

$$2 \frac{1}{A} \frac{dA}{dz} + \frac{1}{k} \frac{dk}{dz} + 2p = 0, \quad (A5)$$

therefore

$$A \propto e^{-\int p dz} k^{-1/2}. \quad (A6)$$

We plug eq.(A6) into eq.(A3) and then achieve

$$\frac{d^2 K}{dz^2} - \left(\frac{1}{\Gamma} \frac{d^2 \Gamma}{dz^2} - q \right) K + \omega^2 K (1 - K^{-4}) = 0, \quad (A7)$$

where $\Gamma = e^{\int p dz}$ and $K = (k/\omega)^{-1/2}$. If $A = \check{A}$ and $k = +\check{k}$ satisfy eq.(A3) and eq.(A4), $A = \check{A}$ and $k = -\check{k}$ will satisfy eq.(A3) and eq.(A4) as well. This means that since eq.(A1) is linear, the general solution of eq.(A1) can be expressed as

$$H = C_+ \Gamma^{-1} \check{K} e^{+i\omega \int \check{K}^{-2} dz} + C_- \Gamma^{-1} \check{K} e^{-i\omega \int \check{K}^{-2} dz}, \quad (\text{A8})$$

where \check{K} is a particular solution of eq.(A7).

Now we need to find a particular solution of eq.(A7). We let $\Xi = \frac{1}{\Gamma} \frac{d^2 \Gamma}{dz^2} - q$ and make $\omega = 1$, which means that we are now using natural units where $c = 1$ and $\omega = 1$, then

$$\frac{d^2 K}{dz^2} + K[(1 - \Xi) - K^{-4}] = 0. \quad (\text{A9})$$

If $\Xi = \text{const}$, it is easy to find a particular solution

$$K = (1 - \Xi)^{-1/4} = 1 + \frac{1}{4}\Xi + \frac{5}{32}\Xi^2 + O(\Xi^3), \quad (\text{A10})$$

which means

$$k = (1 - \Xi)^{1/2} = 1 - \frac{1}{2}\Xi - \frac{1}{8}\Xi^2 + O(\Xi^3), \quad (\text{A11})$$

and this is similar to which of damped oscillations. If $\Xi \neq \text{const}$, we are interested in a special situation that $\Xi(z) = \kappa^2 \tilde{\Xi}(\tilde{z})$, where $\tilde{z} = \kappa z$ and κ is a small amount. The reason which leads to our interest is that if gravitational constant G takes the Yukawa form, we can make parameter λ in Yukawa form satisfies $\kappa = 1/\lambda$ when $1/\lambda \ll \omega/c$. In this situation,

$$K^3 \frac{d^2 K}{d\tilde{z}^2} \kappa^2 - K^4 \tilde{\Xi}(\tilde{z}) \kappa^2 + K^4 - 1 = 0. \quad (\text{A12})$$

We suppose that

$$K = \sum_{n=0}^{\infty} K_n(\tilde{z}) \kappa^{2n}, \quad (\text{A13})$$

and if we plug eq.(A13) into eq.(A12), we will maintain a power series of κ^2 which is equal to 0. We make each of the coefficients of this power series equal to 0, for example,

$$K_0^4 - 1 = 0, \quad (\text{A14})$$

$$K_0^3 K_0'' - K_0^4 \tilde{\Xi} + 4K_0^3 K_1 = 0, \quad (\text{A15})$$

$$(K_0^3 K_1'' + 3K_0^2 K_1 K_0'') - 4K_0^3 K_1 \tilde{\Xi} + (4K_0^3 K_2 + 6K_0^2 K_1^2) = 0. \quad (\text{A16})$$

Finally, by solving all of these equations in order, we will find each of the coefficients in eq.(A13), for example,

$$K_0 = 1, \tag{A17}$$

$$K_1 = \frac{1}{4}\tilde{\Xi}, \tag{A18}$$

$$K_2 = \frac{5}{32}\tilde{\Xi}^2 - \frac{1}{16}\frac{d^2\tilde{\Xi}}{d\tilde{z}^2}, \tag{A19}$$

and these coefficients will make our particular solution of K similar to eq.(A10).

-
- [1] S. K. Roy, R. Koley, and P. Majumdar, Probing the post-minkowskian approximation using recursive addition of self-interactions, *Physical Review D* **102**, 084045 (2020).
 - [2] E. Adelberger, B. Heckel, and A. Nelson, Tests of the gravitational inverse-square law, *Annual Review of Nuclear and Particle Science* **53**, 77 (2003).