

Basic Concepts: Frequent Itemsets (Patterns)


- ☐ Itemset: A set of one or more items
- \square k-itemset: $X = \{x_1, ..., x_k\}$
- (relative) support, s: The fraction of transactions that contains X (i.e., the probability that a transaction
 contains X)
- An itemset X is frequent if the support of X is no less than a minsup threshold (denoted as σ)

Tid	Items bought
10	Beer, Nuts, Diaper
20	Beer, Coffee, Diaper
30	Beer, Diaper, Eggs
40	Nuts, Eggs, Milk
50	Nuts, Coffee, Diaper, Eggs, Milk

- Let minsup = 50%
- □ Freq. 1-itemsets:
 - ☐ Beer: 3 (60%); Nuts: 3 (60%)
 - □ Diaper: 4 (80%); Eggs: 3 (60%)
- ☐ Freq. 2-itemsets:
 - {Beer, Diaper}: 3 (60%)

From Frequent Itemsets to Association Rules

Tid	Items bought
10	Beer, Nuts, Diaper
20	Beer, Coffee, Diaper
30	Beer, Diaper, Eggs
40	Nuts, Eggs, Milk
50	Nuts, Coffee, Diaper, Eggs, Milk



Note: Itemset: $X \cup Y$, a subtle notation!

- \square Association rules: $X \rightarrow Y$ (s, c)
 - □ Support, s: The probability that a transaction contains X ∪ Y
 - □ Confidence, c: The conditional probability that a transaction containing X also contains Y
 - \Box c = sup(X \cup Y) / sup(X)
- □ Association rule mining: Find all of the rules, $X \rightarrow Y$, with minimum support and confidence
- ☐ Frequent itemsets: Let *minsup = 50%*
 - ☐ Freq. 1-itemsets: Beer: 3, Nuts: 3, Diaper: 4, Eggs: 3
 - ☐ Freq. 2-itemsets: {Beer, Diaper}: 3
- ☐ Association rules: Let *minconf* = 50%
 - Beer → Diaper (60%, 100%)
 - Diaper → Beer (60%, 75%) (Q: Are these all rules?)