

# The 3D Organization of Chromatin Explains Evolutionary Fragile Genomic Regions by

Camille Berthelot, Matthieu Muffato,  
Judith Abecassis and Hugues Roest Crollius

Speaker: Ilia Minkin

24th April 2015

# Two types of genome alterations

1. Small point mutations:

ACTTG  
AGT-G

# Two types of genome alterations


## 1. Small point mutations:

ACTTG  
AGT-G

## 2. Large rearrangements:

- ▶ Inversions
- ▶ Transpositions
- ▶ Fusions
- ▶ ...

# Genome Rearrangements



Source: *Dierssen et al, 2009*

# Motivation

Rearrangements:

- ▶ Are a major driving force in evolution
- ▶ Play large role in diseases (e.g. cancer)

Known mechanisms:

- ▶ Non-homologous end joining
- ▶ Non-allelic homologous recombination
- ▶ Replication fork stalling
- ▶ ...

# The Big Question

Are rearrangements more likely to happen in one parts of a genome than the others?

# The Big Question

Are rearrangements more likely to happen in one parts of a genome than the others?

Two hypotheses:

1. Rearrangements are distributed uniformly
2. Some regions are more likely to be disrupted

# A Short Survey

In 1984, Nadeau and Taylor presented first arguments in favor of random breakage



# A Short Survey

In 1984, Nadeau and Taylor presented first arguments in favor of random breakage

Pevzner and Tesler in 2003 showed evidence of fragile regions

# A Short Survey

In 1984, Nadeau and Taylor presented first arguments in favor of random breakage

Pevzner and Tesler in 2003 showed evidence of fragile regions

Ma et al., 2006 argued for random model with higher resolution analysis of rearrangements

# A Short Survey

In 1984, Nadeau and Taylor presented first arguments in favor of random breakage

Pevzner and Tesler in 2003 showed evidence of fragile regions

Ma et al., 2006 argued for random model with higher resolution analysis of rearrangements

Alekseyev and Pevzner, 2010 proposed that fragile regions may born and die

# A Short Survey

In 1984, Nadeau and Taylor presented first arguments in favor of random breakage

Pevzner and Tesler in 2003 showed evidence of fragile regions

Ma et al., 2006 argued for random model with higher resolution analysis of rearrangements

Alekseyev and Pevzner, 2010 proposed that fragile regions may born and die

The story is to be continued...

# The Study


Two questions:

- ▶ Do fragile regions exist?
- ▶ If they do, what is cause of fragility?

A note: fragility is not "physical", it only means higher possibility of rearrangements


# Genomes Representations

Genomes are sequences of gene markers that are unbreakable:




# Genomes Representations

Genomes are sequences of gene markers that are unbreakable:




Here red dashes are breakpoints:



# Methodology

Suppose that we have an ancestral genome and its successors



How does ancestral intergene length affects its breakage rate?



# Methodology

Null hypothesis: breakpoint density is uniform

As intergene length  $\uparrow$ , # of breakpoints  $\uparrow$  as well

It yields Poisson distribution of breakage rate

# Methodology


Boreoeutheria: the last common ancestor of primates, rodents, and laurasiatherians

Stages of the study:

- ▶ Reconstruct gene order of Boreoeutheria
- ▶ Annotate ancestral intergenes
- ▶ Identify breakpoints w.r.t. human, mouse, dog, cow and horse
- ▶ Do Poisson regression of "breakage rate"

Expect linear law if the null hypothesis is true

# Intergene Annotation



# How to Explain the Equation?

$$r = 2.410^{-3} \times L^{0.38}$$

93% of variation in breakpoint occurrence is explained by intergene length

Maybe GC content is the real cause?

# Is GC Content The Explanation?

No.

# Is GC Content The Explanation?

No.

Added GC content in regression – non-significant coefficient

# Are CNEs The Explanation?

CNEs – conservative non-coding elements.  
Located in the intergenes, may affect rearrangement rate. Do they?

# Are CNEs The Explanation?

CNEs – conservative non-coding elements.

Located in the intergenes, may affect rearrangement rate. Do they?

No.



# Are CNEs The Explanation?

CNEs – conservative non-coding elements.  
Located in the intergenes, may affect rearrangement rate. Do they?

No.

Added CNE rate in regression – improved explanation rate only by 3%

# Inversions within Intergenes

OK, maybe some breakpoints are more likely than the others.

We work with gene markers — see only rearrangements disrupting their order.

What if there are many missing rearrangements within intergenes?

We can try to simulate rearrangements and see what happens

# Inversions within Intergenes

Rearrangements have been shown to occur between regions in close 3D proximity in the nucleus

Contact probability is a good proxy for rearrangement probability

Simulate and sample breakpoint pairs, choose detectable ones

Even if we restrict to detectable breakpoints only, simulation confirms the random breakpoint hypothesis

# Open Chromatin is the Culprit

Stick with the simulation – restrict rearrangements to only **open chromatin** regions

Voilà – simulation coincides with the model!  
It implies that chromatin state and proximity of genes may explain fragility of some genomic regions

# Conclusion

It seems that chromatin state and proximity of genes may explain fragility of some genomic regions

Thank you!