

SUBJECT ID = RR

112

ENDOMETRIOSIS PELVIC MRI ASSESSMENT –

BR PROFORMA REPORT BLIND REVIEW

Uterus

- 1: Absent
- 2: Present

Uterine anatomy

- 1. Conventional
- 2. Arcuate
- 3. Septate
 - a. Full septum
 - b. Subseptate
- 4. Bicornuate unicollis
- 5. Bicornuate bicollis
- 6. Didelphys
- 7. Other (free text enabled).

Uterine Lie (can be more than one selection)

- 1. Anteverted
- 2. Anteverted
- 3. Retroverted
- 4. Retroflexed
- 5. Axial
- 6. Others (please specify) (Free text enabled)

Uterine Size (body + cervix – 3 planes in mm)

- 1. (Free text) 83x52x62

Endometrial thickness (sag plane in mm to nearest mm)

- 1. (Free text) 7mm

Endometrial lesions

- 1. Not identified.
- 2. Present, Polyp.
 - 2b-1: No. of polyps (free text)
 - 2b-2: Size of each polyp. (free text)

Adenomyosis

- 1. No MRI supportive features
- 2. Supportive MRI features as described:
 - 1. Submucosal cysts.
 - 2. Abnormal junctional zone thickening and measurement
 - i. Anterior (mm)
 - ii. Fundal (mm)
 - iii. Posterior (mm)

Presence of an adenomyoma

- 1: No
- 2: Yes

Fibroids

- 1: No
- 2: Yes
 - 2a: Number of fibroids:
 - 2b: Largest fibroids (location and size mm all fibroids >10mm and/or impact on the cavity) – (Free text)
 - 2b: Submucosal fibroids
 - 2b-0: No

REVIEWER INITIALS

2b-1: Yes

2b-1-1: (description: free text)

Left ovary

- 1: Absent (Branching logic – move to “Right ovary”)
- 2: Present

Left ovary size (3 planes and volume)

- 1. NN x NN x NN mm
- 2. Volume (above x 0.52).

33x19x21
6.8

Left ovary follicle measurements and count

- 1. N follicles between 2 and 9 mm in diameter
 - a. (Free text) >25
- 2. N follicles > 9 mm
 - a. (Free text) 0
- 3. Dominant follicle
 - a. Y
 - b. N.

Left ovary position

- 1: Lateral adnexa. Unremarkable.
- 2: High positioning in iliac fossa.
- 3: Tethered/ distorted appearances – (may be multiple options)
 - 3a: Intimate relationship to the lateral uterus
 - 3b Intimate relationship/ tethering to adjacent bowel.
 - 3c. Tethering to pelvic sidewall
 - 3d: Tethering to the uterosacral ligament

SUBJECT ID = RR

112

3e: Other: (free text)

Left ovarian endometrioma

- 1: No
- 2: Yes
- 1a: Size: NN in millimetres (mm)
- 1b: T2 shading (intermediate/low homogeneous T2 signal).
- 1b-0: No
- 1b-1: Yes
- 1c: T1 fat sat hyperintense
- 1c-0: Absent
- 1c-1: Present
- 1d: Internal nodularity, septation or other complexity.
- 1d-1: No
- 1d-2: Yes
- 1-d-2-1: (Free text)

Left ovarian lesions/cysts other than suspected endometriomas

- 1: Not classifiable
- 2: Lesion features
- 2a: Unilocular cyst
- 2b: Unilocular-solid cyst
- 2c: Multilocular cyst
- 2d: Multilocular-solid cyst
- 2e: Solid
- 3: Vascularity (post gadolinium enhancement).
- 3a: Present

3b: Absent

4 Fat component (T1/ T2 hyperintense. Low signal on fat suppression).

4a: Present.

4b: Absent.

Right ovary

- 1: Absent (Branching logic – move to "Adnexa")
- 2: Present

Right ovary size (3 planes and volume)

1. NN x NN x NN mm

29x21x34

2. Volume (above x 0.52).

10.8

Right ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter
- a. (Free text) 25
2. N follicles > 9 mm
- a. (Free text) 1
3. Dominant follicle

a. Y

b. N.

16mm.

Right ovary position

- 1: Lateral adnexa. Unremarkable.
- 2: High positioning in iliac fossa.
- 3: Tethered/ distorted appearances – may be multiple options.
- 3a: Intimate relationship to the lateral uterus
- 3b: Intimate relationship/ tethering to

REVIEWER INITIALS

SM

adjacent bowel.

3c. Tethering to pelvic sidewall

3d: Tethering to the uterosacral ligament

Right ovarian endometrioma

- 1: No
- 2: Yes
- 2a: Size: NN in millimetres (mm)
- 2b: T2 shading (intermediate/low homogeneous T2 signal).
- 2b-0: No
- 2b-1: Yes
- 2c: T1 fat sat hyperintense
- 2c-0: Absent
- 2c-1: Present
- 2d: Internal nodularity, septation, complex.
- 2d-1: No
- 2d-2: Yes

Right ovarian lesions/cysts other than suspected endometriomas

- 1: Not classifiable
- 2: Lesion features
- 2a: Unilocular cyst
- 2b: Unilocular-solid cyst
- 2c: Multilocular cyst
- 2d: Multilocular-solid cyst
- 2e: Solid
- 3: Vascularity (post gadolinium enhancement).
- 3a: Present

SUBJECT ID = RR

112

3b: Absent

4 Fat component (T1/ T2 hyperintense. Low signal on fat suppression).

4a: Present.

4b: Absent.

Adnexa

1: Hydrosalpinx

1a: No

1b: Yes

2: Hematosalpinx

2a: No

2b: Yes

3: Other (free text).

Are both ovaries immediately approximated "kissing"?

1: No

2: Yes

Urinary bladder nodule

Definition: Is there presence of a nodule in the bladder.

1: Absent

2: Present

2a: Size: NN in millimetres (mm)

Uterovesical region

Definition: Assessment of whether there is a visible preserved fat plane +/- physiologic fluid +/- absent distortion between the anterior uterine serosa and bladder.

1: Normal.

2: Abnormal.

2a: (free text if required)

smooth
sw

Ureteric nodule(s)?

1: Absent

2: Present

2a: Location (free text + distance to ureteric orifice/ VUJ)

2b: Size (mm)

Pouch of Douglas obliteration

Definition: Assessment for abnormal loss of fat plane +/- physiologic fluid and immediate approximation of rectosigmoid and/or small bowel to the posterior uterine serosa, cervix +/- vaginal wall.

Discrete linear bands may be visible (↓ T1, ↓ T2)

1: Negative

2: Partial

2a: Left

2b: Right

3: Complete

3a: Positive = obliteration.

3b: Positive = band adhesions.

Nodules present on the posterior vaginal fornix?

Definition: Thickening of superior 1/3 of posterior vaginal wall +/- nodularity. Nodules: ↓ T2 ↑ T1 (if active haemorrhagic deposits)

1: No

2: Yes

REVIEWER INITIALS

SM

2a: Dimension of nodule to be measured in millimetres (mm).

2b1: Inactive.

2b2: Active

Vaginal forniceal elevation?

Definition: Upper level of fornix on sagittal view is superior to the angle of the uterine isthmus with stretching of vaginal wall, and/or acute angulation of the fornix.

1: No

2: Yes

2a: Left.

2b: Right

2c: Left and Right.

Rectovaginal nodules present?

Definition: Presence of deep infiltrating endometriosis in the anterior rectal wall and posterior vaginal fornix, located below the peritoneum of the Pouch of Douglas. Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑ T1, ↑ to intermediate- T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes

2a: Size (mm)

2b1: Inactive.

2b2: Active

SUBJECT ID = RR

112

Uterosacral ligament nodules or thickening?

Definition: Inactive/ fibrotic disease nodules characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑ T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes nodules

2a: Left

2a-1: Size (mm)

2b: Right

2b-1: Size (mm)

2c1: Inactive.

2c2: Active

3: Yes thickening.

3a: Left.

3b: Right

3c: Both.

Retrocervical nodule present?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑ T1, ↑ to intermediate- T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes

2a: Size (mm)

2b1: Inactive.

2b2: Active

Rectum and colon:

Is there bowel deep infiltrating endometriosis seen?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑ T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

"Mushroom cap sign" is specific to severe invasive bowel endometriosis and is characterized as a plaque with ↓ T2 at its 'base' and ↑ T2 at its 'cap'.

1: No

2: Yes

2a: Distance from the anal verge

2a-1: Length (mm)

2b: Lesion type

2b-1: Isolated lesion

2b-2: Multiple lesions

2b-3: Curved lesion

2b-4: Straight lesion

2c: Maximal depth layer of invasion each lesion (muscularis, submucosa, mucosa).

2c-1: Lesion 1: (free text)

(2c-2: Lesion 2 (free text) - delete if not relevant

(2c-3 etc.)

2c: Is it stuck to any structures or free lying?

2d-1: Vagina

2d-2: Uterus

2d-3: Uterosacral ligaments

REVIEWER INITIALS

SM

2d-4: Ovary

2d: Plaque thickness

2a: 1-5mm.

2b: 6-10mm.

2c: >11mm.

2e: Activity

2f1: Inactive.

2f2: Active.

2f: "Mushroom cap" appearance:

2g1: Present.

2g2: Absent.

Is there evidence of tethering of the bowel?

1: No

2: Yes, tethered to

2a: Uterus

2b: L. ovary

2c: R. ovary

2d: L. uterosacral ligament nodule

2e: R. uterosacral ligament nodule

2f: L pelvic side wall.

2g: R pelvic side wall.

2h: Other.

Any other salient findings on the study:

1. No

2. Yes

a. (Free text).

Scan/ Photo/ Email: kate.cook@benzonradiology.com.au