

SUBJECT ID = RR

032

ENDOMETRIOSIS PELVIC MRI ASSESSMENT –

BR PROFORMA REPORT BLIND REVIEW

Uterus

- 1: Absent
2: Present

Uterine anatomy

1. Conventional
2. Arcuate
3. Septate
 a. Full septum
 b. Subseptate
4. Bicornuate unicollis
5. Bicornuate bicollis
6. Didelphys
7. Other (free text enabled).

Uterine Lie (can be more than one selection)

1. Anteverted
2. Anteflexed
3. Retroverted
4. Retroflexed
5. Axial
6. Others (please specify) (Free text enabled)

Uterine Size (body + cervix – 3 planes in mm)

1. (Free text). 66 x 87 x 101

Endometrial thickness (sag plane in mm to nearest mm)

1. (Free text) 2mm

Endometrial lesions

1. Not identified.
2. Present. Polyp.
2b-1: No. of polyps (free text)
2b-2: Size of each polyp. (free text)

Adenomyosis

1. No MRI supportive features
2. Supportive MRI features as described:
 1. Submucosal cysts.
 2. Abnormal junctional zone thickening and measurement
 i. Anterior (mm)
 ii. Fundal (mm)
 iii. Posterior (mm)

Presence of an adenomyoma

- 1: No
2: Yes

Fibroids

- 1: No
2: Yes
2a: Number of fibroids: 4
2b: Largest fibroids (location and size mm all fibroids >10mm and/or impact on the cavity) – (Free text)
2b-0: Submucosal fibroids
2b-0: No

REVIEWER INITIALS

2b-1: Yes

2b-1-1: (description: free text)

③ + ④ about endometrium with minimal submucosal extension

Left ovary

- 1: Absent (Branching logic – move to “Right ovary”)
2: Present

Left ovary size (3 planes and volume)

1. NN x NN x NN mm 39 x 26 x 23
2. Volume (above x 0.52). 12.2 cc

Left ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter
 a. (Free text) 7
2. N follicles > 9 mm
 a. (Free text) 4
3. Dominant follicle
 a. Y
 b. N.

Left ovary position

- 1: Lateral adnexa. Unremarkable.
2: High positioning in iliac fossa.
3: Tethered/ distorted appearances – (may be multiple options)
3a: Intimate relationship to the lateral uterus
3b Intimate relationship/ tethering to adjacent bowel.
3c. Tethering to pelvic sidewall
3d: Tethering to the uterosacral ligament

SUBJECT ID = RR 032
3e: Other: (free text)

Left ovarian endometrioma

- 1: No
2: Yes
1a: Size: NN in millimetres (mm)
1b: T2 shading (intermediate/low homogeneous T2 signal).
1b-0: No
1b-1: Yes
1c: T1 fat sat hyperintense
1c-0: Absent
1c-1: Present
1d: Internal nodularity, septation or other complexity.
1d-1: No
1d-2: Yes
1-d-2-1: (Free text)

Left ovarian lesions/cysts other than suspected endometriomas

- 1: Not classifiable
2: Lesion features
2a: Unilocular cyst
2b: Unilocular-solid cyst
2c: Multilocular cyst
2d: Multilocular-solid cyst
2e: Solid
3: Vascularity (post gadolinium enhancement).
3a: Present

- 3b: Absent
4 Fat component (T1/ T2 hyperintense. Low signal on fat suppression).
4a: Present.
4b: Absent.

Right ovary

- 1: Absent (Branching logic – move to “Adnexa”)
2: Present

Right ovary size (3 planes and volume)

1. NN x NN x NN mm 20 x 17 x 27
2. Volume (above x 0.52). 4.8cc

Right ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter
a. (Free text) 2
2. N follicles > 9 mm
a. (Free text)
3. Dominant follicle
a. Y
b. N.

Right ovary position

- 1: Lateral adnexa. Unremarkable.
2: High positioning in iliac fossa.
3: Tethered/ distorted appearances – may be multiple options.
3a: Intimate relationship to the lateral uterus
3b Intimate relationship/ tethering to

REVIEWER INITIALS HJ
adjacent bowel.

- 3c. Tethering to pelvic sidewall
3d: Tethering to the uterosacral ligament

Right ovarian endometrioma

- 1: No
2: Yes
2a: Size: NN in millimetres (mm)
2b: T2 shading (intermediate/low homogeneous T2 signal).
2b-0: No
2b-1: Yes
2c: T1 fat sat hyperintense
2c-0: Absent
2c-1: Present
2d: Internal nodularity, septation, complex.
2d-1: No
2d-2: Yes

Right ovarian lesions/cysts other than suspected endometriomas

- 1: Not classifiable
2: Lesion features
2a: Unilocular cyst
2b: Unilocular-solid cyst
2c: Multilocular cyst
2d: Multilocular-solid cyst
2e: Solid
3: Vascularity (post gadolinium enhancement).
3a: Present

SUBJECT ID = RR

082

3b: Absent

4 Fat component (T1/ T2 hyperintense. Low signal on fat suppression).

4a: Present.

4b: Absent.

Adnexa

1: Hydrosalpinx

1a: No

1b: Yes

2: Hematosalpinx

2a: No

2b: Yes

3: Other (free text).

Are both ovaries immediately approximated "kissing"?

1: No

2: Yes

Urinary bladder nodule

Definition: Is there presence of a nodule in the bladder.

1: Absent

2: Present

2a: Size: NN in millimetres (mm)

Uterovesical region

Definition: Assessment of whether there is a visible preserved fat plane +/- physiologic fluid +/- absent distortion between the anterior uterine serosa and bladder.

1: Normal.

2: Abnormal.

2a: (free text if required)

Ureteric nodule(s)?

1: Absent

2: Present

2a: Location (free text + distance to ureteric orifice/ VUJ)

2b: Size (mm)

Pouch of Douglas obliteration

Definition: Assessment for abnormal loss of fat plane +/- physiologic fluid and immediate approximation of rectosigmoid and/or small bowel to the posterior uterine serosa, cervix +/- vaginal wall.

Discrete linear bands may be visible (↓ T1, ↓ T2)

1: Negative

2: Partial

2a: Left

2b: Right

3: Complete

3a: Positive = obliteration.

3b: Positive = band adhesions.

Nodules present on the posterior vaginal fornix?

Definition: Thickening of superior 1/3 of posterior vaginal wall +/- nodularity. Nodules: ↓ T2 ↑ T1 (if active haemorrhagic deposits)

1: No

2: Yes

REVIEWER INITIALS

mJ

2a: Dimension of nodule to be measured in millimetres (mm).

2b1: Inactive.

2b2: Active

Vaginal forniceal elevation?

Definition: Upper level of fornix on sagittal view is superior to the angle of the uterine isthmus with stretching of vaginal wall, and/or acute angulation of the fornix.

1: No

2: Yes

2a: Left.

2b: Right

2c: Left and Right.

Rectovaginal nodules present?

Definition: Presence of deep infiltrating endometriosis in the anterior rectal wall and posterior vaginal fornix, located below the peritoneum of the Pouch of Douglas. Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal. Active disease as ↑ T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes

2a: Size (mm)

2b1: Inactive.

2b2: Active

SUBJECT ID = RR 032

Uterosacral ligament nodules or thickening?

Definition: Inactive/ fibrotic disease nodules characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

- 1: No
- 2: Yes nodules
- 2a: Left
- 2a-1: Size (mm)
- 2b: Right
- 2b-1: Size (mm)
- 2c1: Inactive.
- 2c2: Active
- 3: Yes thickening.
- 3a: Left.
- 3b: Right
- 3c: Both.

Retrocervical nodule present?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

- 1: No
- 2: Yes
- 2a: Size (mm)
- 2b1: Inactive.
- 2b2: Active

Rectum and colon:

Is there bowel deep infiltrating endometriosis seen?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).
"Mushroom cap sign" is specific to severe invasive bowel endometriosis and is characterized as a plaque with ↓ T2 at its 'base' and ↑ T2 at its 'cap'.

- 1: No
- 2: Yes
- 2a: Distance from the anal verge
- 2a-1: Length (mm)
- 2b: Lesion type
- 2b-1: Isolated lesion
- 2b-2: Multiple lesions
- 2b-3: Curved lesion
- 2b-4: Straight lesion
- 2c: Maximal depth layer of invasion each lesion (muscularis, submucosa, mucosa).
- 2c-1: Lesion 1: (free text)
- (2c-2: Lesion 2 (free text) - delete if not relevant)
- (2c-3 etc.)
- 2c: Is it stuck to any structures or free lying?
- 2d-1: Vagina
- 2d-2: Uterus
- 2d-3: Uterosacral ligaments

REVIEWER INITIALS MJ

- 2d-4: Ovary
- 2d: Plaque thickness
- 2a: 1-5mm.
- 2b: 6-10mm.
- 2c: >11mm.
- 2e: Activity
- 2f1: Inactive.
- 2f2: Active.
- 2f: "Mushroom cap" appearance:
- 2g1: Present.
- 2g2: Absent.

Is there evidence of tethering of the bowel?

- 1: No
- 2: Yes, tethered to
- 2a: Uterus
- 2b: L. ovary
- 2c: R. ovary
- 2d: L. uterosacral ligament nodule
- 2e: R. uterosacral ligament nodule
- 2f: L pelvic side wall.
- 2g: R pelvic side wall.
- 2h: Other.

Any other salient findings on the study:

1. No
2. Yes
- a. (Free text).

Scan/ Photo/ Email: kate.cook@bensonradiology.com.au