

SUBJECT ID = RR

117

ENDOMETRIOSIS PELVIC MRI ASSESSMENT –

BR PROFORMA REPORT BLIND REVIEW

Uterus

- 1: Absent
2: Present

Uterine anatomy

1. Conventional
2. Arcuate
3. Septate
 a. Full septum
 b. Subseptate
4. Bicornuate unicollis
5. Bicornuate bicollis
6. Didelphys
7. Other (free text enabled).

Uterine Lie (can be more than one selection)

1. Anteverted
2. Anteflexed
3. Retroverted
4. Retroflexed
5. Axial
6. Others (please specify) (Free text enabled)

Uterine Size (body + cervix – 3 planes in mm)

1. (Free text). 57x31x39

Endometrial thickness (sag plane in mm to nearest mm)

1. (Free text) 6mm

Endometrial lesions

1. Not identified.
2. Present. Polyp.

2b-1: No. of polyps (free text)

2b-2: Size of each polyp. (free text)

Adenomyosis

1. No MRI supportive features
2. Supportive MRI features as described:
 1. Submucosal cysts.
 2. Abnormal junctional zone thickening and measurement
 i. Anterior (mm)
 ii. Fundal (mm)
 iii. Posterior (mm)

Presence of an adenomyoma

- 1: No
2: Yes

Fibroids

- 1: No
2: Yes

2a: Number of fibroids:

2b: Largest fibroids (location and size mm all

fibroids >10mm and/or impact on the cavity) – (Free text)

2b: Submucosal fibroids

2b-0: No

REVIEWER INITIALS

SM

2b-1: Yes

2b-1-1: (description: free text)

Left ovary

- 1: Absent (Branching logic – move to “Right ovary”)
2: Present

Left ovary size (3 planes and volume)

1. NN x NN x NN mm
2. Volume (above x 0.52).

19x12x13mm
1.5ml

Left ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter
 a. (Free text) 2
2. N follicles > 9 mm
 a. (Free text) 0
3. Dominant follicle

- a. Y
b. N.

Left ovary position

- 1: Lateral adnexa. Unremarkable.
2: High positioning in iliac fossa.
3: Tethered/ distorted appearances – (may be multiple options)
 3a: Intimate relationship to the lateral uterus
 3b: Intimate relationship/ tethering to adjacent bowel.
 3c: Tethering to pelvic sidewall
 3d: Tethering to the uterosacral ligament

SUBJECT ID = RR

3e: Other: (free text)

Left ovarian endometrioma

- 1: No
- 2: Yes
- 1a: Size: NN in millimetres (mm)
- 1b: T2 shading (intermediate/low homogeneous T2 signal).
- 1b-0: No
- 1b-1: Yes
- 1c: T1 fat sat hyperintense
- 1c-0: Absent
- 1c-1: Present
- 1d: Internal nodularity, septation or other complexity.
- 1d-1: No
- 1d-2: Yes
- 1-d-2-1: (Free text)

Left ovarian lesions/cysts other than suspected endometriomas

- 1: Not classifiable
- 2: Lesion features
- 2a: Unilocular cyst
- 2b: Unilocular-solid cyst
- 2c: Multilocular cyst
- 2d: Multilocular-solid cyst
- 2e: Solid
- 3: Vascularity (post gadolinium enhancement).
- 3a: Present

3b: Absent

4 Fat component (T1/ T2 hyperintense. Low signal on fat suppression).

4a: Present.

4b: Absent.

Right ovary

- 1: Absent (Branching logic – move to "Adnexa")
- 2: Present

Right ovary size (3 planes and volume)

1. NN x NN x NN mm
2. Volume (above x 0.52).

Right ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter
- a. (Free text)
2. N follicles > 9 mm
- a. (Free text)
3. Dominant follicle

a. Y

b. N.

Right ovary position

- 1: Lateral adnexa. Unremarkable.
- 2: High positioning in iliac fossa.
- 3: Tethered/ distorted appearances – may be multiple options.
- 3a: Intimate relationship to the lateral uterus
- 3b Intimate relationship/ tethering to

REVIEWER INITIALS

adjacent bowel.

3c. Tethering to pelvic sidewall

3d: Tethering to the uterosacral ligament

Right ovarian endometrioma

- 1: No
- 2: Yes
- 2a: Size: NN in millimetres (mm)
- 2b: T2 shading (intermediate/low homogeneous T2 signal).
- 2b-0: No
- 2b-1: Yes
- 2c: T1 fat sat hyperintense
- 2c-0: Absent
- 2c-1: Present
- 2d: Internal nodularity, septation, complex.
- 2d-1: No
- 2d-2: Yes

Right ovarian lesions/cysts other than suspected endometriomas

- 1: Not classifiable
- 2: Lesion features
- 2a: Unilocular cyst
- 2b: Unilocular-solid cyst
- 2c: Multilocular cyst
- 2d: Multilocular-solid cyst
- 2e: Solid
- 3: Vascularity (post gadolinium enhancement).
- 3a: Present

SUBJECT ID = RR

117

3b: Absent

4 Fat component (T1/ T2 hyperintense. Low signal on fat suppression).

4a: Present.

4b: Absent.

Adnexa

1: Hydrosalpinx

1a: No

1b: Yes

2: Hematosalpinx

2a: No

2b: Yes

3: Other (free text).

Are both ovaries immediately approximated "kissing"?

1: No

2: Yes

Urinary bladder nodule

Definition: Is there presence of a nodule in the bladder.

1: Absent

2: Present

2a: Size: NN in millimetres (mm)

Uterovesical region

Definition: Assessment of whether there is a visible preserved fat plane +/- physiologic fluid +/- absent

distortion between the anterior uterine serosa and bladder.

1: Normal.

2: Abnormal.

2a: (free text if required)

Ureteric nodule(s)?

1: Absent

2: Present

2a: Location (free text + distance to ureteric orifice/ VUJ)

2b: Size (mm)

Pouch of Douglas obliteration

Definition: Assessment for abnormal loss of fat plane +/- physiologic fluid and immediate approximation of rectosigmoid and/or small bowel to the posterior uterine serosa, cervix +/- vaginal wall.

Discrete linear bands may be visible (↓ T1, ↓ T2)

1: Negative

2: Partial

2a: Left

2b: Right

3: Complete

3a: Positive = obliteration.

3b: Positive = band adhesions.

Nodules present on the posterior vaginal fornix?

Definition: Thickening of superior 1/3 of posterior vaginal wall +/- nodularity. Nodules: ↓ T2 ↑T1 (if active haemorrhagic deposits)

1: No

2: Yes

REVIEWER INITIALS

SH

2a: Dimension of nodule to be measured in millimetres (mm).

2b1: Inactive.

2b2: Active

Vaginal forniceal elevation?

Definition: Upper level of fornix on sagittal view is superior to the angle of the uterine isthmus with stretching of vaginal wall, and/or acute angulation of the fornix.

1: No

2: Yes

2a: Left.

2b: Right

2c: Left and Right.

Rectovaginal nodules present?

Definition: Presence of deep infiltrating endometriosis in the anterior rectal wall and posterior vaginal fornix, located below the peritoneum of the Pouch of Douglas. Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes

2a: Size (mm)

2b1: Inactive.

2b2: Active

SUBJECT ID = RR

17

Uterosacral ligament nodules or thickening?

Definition: Inactive/ fibrotic disease nodules characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

- 1: No
2: Yes nodules
2a: Left
2a-1: Size (mm)
2b: Right
2b-1: Size (mm)
2c1: Inactive.
2c2: Active
3: Yes thickening.
3a: Left.
3b: Right
3c: Both.

Retrocervical nodule present?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

- 1: No
Yes
2a: Size (mm)
2b1: Inactive.
2b2: Active

Rectum and colon:

Is there bowel deep infiltrating endometriosis seen?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

"Mushroom cap sign" is specific to severe invasive bowel endometriosis and is characterized as a plaque with ↓ T2 at its 'base' and ↑ T2 at its 'cap'.

- 1: No
2: Yes
2a: Distance from the anal verge
2a-1: Length (mm)
2b: Lesion type
2b-1: Isolated lesion
2b-2: Multiple lesions
2b-3: Curved lesion
2b-4: Straight lesion
2c: Maximal depth layer of invasion each lesion (muscularis, submucosa, mucosa).
2c-1: Lesion 1: (free text)
(2c-2: Lesion 2 (free text) - delete if not relevant
(2c-3 etc.)
2c: Is it stuck to any structures or free lying?
2d-1: Vagina
2d-2: Uterus
2d-3: Uterosacral ligaments

REVIEWER INITIALS

SK

- 2d-4: Ovary
2d: Plaque thickness
2a: 1-5mm.
2b: 6-10mm.
2c: >11mm.
2e: Activity
2f1: Inactive.
2f2: Active.
2f: "Mushroom cap" appearance:
2g1: Present.
2g2: Absent.

Is there evidence of tethering of the bowel?

- 1: No
2: Yes, tethered to
2a: Uterus
2b: L. ovary
2c: R. ovary
2d: L. uterosacral ligament nodule
2e: R. uterosacral ligament nodule
2f: L pelvic side wall.
2g: R pelvic side wall.
2h: Other.

Any other salient findings on the study:

1. No
2. Yes
a. (Free text)
Anterior uterine plaque ~ 26mm
myometrial invasion

Scan/ Photo/ Email: kate.cook@bensonradiology.com.au