SUBJECT ID = RR

ENDOMETRIOSIS PELVIC MRI ASSESSMENT -BR PROFORMA REPORT BLIND REVIEW

## Uterus

Absent

Present

Uterine anatomy

Conventional

- Arcuate
- Septate
  - Full septum
  - Subseptate
- Bicornuate unicollis
- Bicornuate bicollis
- Didelphys
- Other (free text enabled).

Uterine Lie (can be more than one selection)

- Anteverted
- Anteflexed
- Retroverted
- Retroflexed
- 5. Axial
- Others (please specify) (Free text enabled)

Uterine Size (body + cervix - 3 planes in mm )

Endometrial thickness (sag plane in mm to nearest mm)

1. (Free text)

**Endometrial lesions** 

- Not identified.
- 2. Present. Polyp.

2b-1: No. of polyps (free text)

Size of each polyp. (free text)

Adenomyosis

No MRI supportive features

- Supportive MRI features as described:
  - Submucosal cysts.
  - Abnormal junctional zone thickening and measurement
    - Anterior (mm)
    - Fundal (mm)
    - Posterior (mm)

Presence of an adenomyoma

Yes

Fibroids

Yes

No

2a: Number of fibroids:

Largest fibroids (location and size mm all 2b: fibroids >10mm and/or iimpact on the cavity) - (Free text)

> Submucosal fibroids 2b:

> > 2b-0: No

REVIEWER INITIALS

2b-1:

2b-1-1: (description: free text)

Left ovary

Absent (Branching logic - move to "Right ovary") 1:

Present

1. NN x NN x NN mm 22 x 3 x 28 2. Volume (above x 0.52). 4.2 ml.

Left ovary follicle measurements and count

N follicles between 2 and 9 mm in diameter

a. (Free text)

2. N follicles > 9 mm

a. (Free text)

Dominant follicle

Left ovary position

Lateral adnexa. Unremarkable.

High positioning in iliac fossa.

3: Tethered/ distorted appearances - (may be multiple options)

> 3a: Intimate relationship to the lateral uterus 3b Intimate relationship/ tethering to adjacent bowel.

3c. Tethering to pelvic sidewall 3d:Tethering to the uterosacral ligament Left ovarian endometrioma

1: / No

2: Yes

1a: Size: NN in millimetres (mm)

 T2 shading (intermediate/low homogeneous T2 signal).

1b-0: No

1b-1: Yes

1c: T1 fat sat hyperintense

1c-0: Absent

1c-1: Present

 Internal nodularity, septation or other complexity.

1d-1: No

1d-2: Yes

1-d-2-1: (Free text)

Left ovarian lesions/cysts other than suspected

endometriomas

2:

Not classifiable

Lesion features

2a: Unilocular eyst

2b: Unilocylar-solid cyst

2c: Multilocular cyst

2d: Multilocular-solid cyst

2e: Solid

Yascularity (post gadolinium enhancement).

3a: Present

3b: Absent

4 Fat component (T1/T2 hyperintense. Low signal on fat suppression).

4a: Present.

1b: Absent.

Right ovary

1: Absent (Branching logic - move to "Adnexa")

72: Present

Right ovary size (3 planes and volume)

1. NN x NN x NN mm 16 × 13 × 31

2. Volume (above x 0.52). 2 . 4

Right ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter

a. (Free text)

N follicles > 9 mm

a. (Free text)

3. Dominant follicle

а. Y b. N.

Right ovary position

1: Lateral adnexa. Unremarkable.

High positioning in iliac fossa.

Tethered/ distorted appearances – may be multiple options.

3a: Intimate relationship to the lateral uterus

3b Intimate relationship/ tethering to

REVIEWER INITIALS

adjacent bowel.

3c. Tethering to pelvic sidewall

3d: Tethering to the uterosacral ligament

Right ovarian endometrioma

No

Yes

2a: Size: NN in millimetres (mm)

2b: T2 shading (intermediate/low

homogeneous T2 signal).

2b-0: No

2b-1: Yes

2c: T1 fat sat hyperintense

2c-0: Absent

2c-1: Present

2d: Internal nodularity, septation, complex.

2d-1: No

2d-2: Yes

Right ovarian lesions/cysts other than suspected endometriomas

1: Not classifiable

2: Lesion features

2a: Unilocular cyst

2b: Unilocular-solid cyst

2c: Multilocular cyst

d: Multilocular-solid cyst

2e: Soli

Vascularity (post gadolinium enhancement).

3a: Present

SUBJECT ID = RR

3b: Absent

Fat component (T1/T2 hyperintense. Low signal on fat suppression).

> 4a: Present.

> > Absent.

Adnexa

2:

1: Hydrosalpinx

No

Yes

Hematosalpinx

No

Yes

3: Other (free text).

Are both ovaries immediately approximated "kissing"?

2:

Urinary bladder nodule

Definition: Is there presence of a nodule in the bladder.

Absent

2: Present

> Size: NN in millimetres (mm) 2a:

Uterovesical region

Definition: Assessment of whether there is a visible preserved fat plane +/- physiologic fluid +/- absent

distortion between the anterior uterine serosa and bladder.

Normal.

2: Abnormal.

> 2a: (free text if required)

Ureteric nodule(s)?

Absent

Present

2a: Location (free text + distance to ureteric

orifice/VUJ)

2b: Size (mm)

Pouch of Douglas obliteration

Definition: Assessment for abnormal loss of fat plane +/physiologic fluid and immediate approximation of rectosigmoid and/or small bowel to the posterior uterine serosa, cervix +/- vaginal wall.

Discrete linear bands may be visible (\$\sqrt{T1}, \$\sqrt{T2}\$)

Negative

2:

Partial

2a: Left

2b: Right

Complete

3a: Positive = obliteration.

Positive = band adhesions. 3b:

Nodules present on the posterior vaginal fornix?

Definition: Thickening of superior 1/3 of posterior vaginal wall +/- nodularity. Nodules: ↓ T2 ↑T1 (if active

haemorrhagic deposits)

No

Yes

REVIEWER INITIALS

Dimension of nodule to be measured in 2a: millimetres (mm).

> 2b1: Inactive.

2b2: Active

Vaginal forniceal elevation?

Definition: Upper level of fornix on sagittal view is superior to the angle of the uterine isthmus with stretching of vaginal wall, and/or acute angulation of the fornix.

Yes 2:

2a: Left.

2b: Right

2c: Left and Right.

Rectovaginal nodules present?

Definition: Presence of deep infiltrating endometriosis in the anterior rectal wall and posterior vaginal fornix, located below the peritoneum of the Pouch of Douglas. Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as 11, 1 to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

2: Yes

> 2a: Size (mm)

2b1: Inactive.

2b2: Active SUBJECT ID = RR

Uterosacral ligament nodules or thickening?

Definition: Inactive/fibrotic disease nodules characterised as  $\downarrow$  T1  $\downarrow$  T2 signal.

Active disease as 171, 1 to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

1: / No

2: Yes nodules

2a: Left

2a-1: Size (mm)

2b: Right

2b-1: Size (mm)

2c1: Inactive.

2c2: Active

3: Yes thickening.

3a: Left.

3b: Right

3c: Both.

Retrocervical nodule present?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to Intermediate- T2 signal (hemorrhagic/ proteinacous content + glandular deposits).

: / No

Yes

2a: Size (mm)

2b1: Inactive.

2b2: Active

Rectum and colon:

Is there bowel deep infiltrating endometriosis seen?

Definition: Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal.

Active disease as ↑T1, ↑ to intermediate-T2 signal (hemorrhagic/ proteinacous content + glandular deposits).

"Mushroom cap sign" is specific to severe invasive bowel endometriosis and is characterized as a plaque with ↓ T2 at its 'base' and ↑ T2 at its 'cap'.

1:/

No

Yes

2a: Distance from the anal verge

2a-1: Length (mm)

2b: Lesion type

2b-1: Isolated lesion

2b-2: Multiple lesions

2b-3: Curved lesion

2b-4: Straight lesion

 Maximal depth layer of invasion each leasion (muscularis, submucosa, mucosa).

2c-1: Lesion 1: (free text)

(2c-2: Lesion 2 (free text) - delete if not relevant

(2c-3 etc.)

2c: Is it stuck to any structures or free lying?

2d-1: Vagina

2d-2: Uterus

2d-3: Uterosacral ligaments

REVIEWER INITIALS

2d-4: Ovary

2d: Plaque thickness

2a: 1-5mm.

2b: 6-10mm.

2c: >11mm.

2e: Activity

2f1: Inactive.

2f2: Active.

2f: "Mushroom cap" appearance:

2g1: Present.

2g2: Absent.

Is there evidence of tethering of the bowel?

/ No

Yes, tethered to

2a: Uterus

2b: L. ovary

2c: R. ovary

2d: L. uterosacral ligament nodule

2e: R. uterosacral ligament nodule

2f: L pelvic side wall.

2g: R pelvic side wall.

2h: Other.

Any other salient findings on the study:

1. N6

Z. Yes

. (Free text).

Scan/ Photo/ Emaii: kate.cook@bensonradiology.com.au