

SUBJECT ID = RR

429

ENDOMETRIOSIS PELVIC MRI ASSESSMENT -

BR PROFORMA REPORT BLIND REVIEW

Uterus

1. Absent
2. Present

Uterine anatomy

1. Conventional
2. Arcuate
3. Septate
a. Full septum
b. Subseptate
4. Bicornuate unicollis
5. Bicornuate bicollis
6. Didelphys
7. Other (free text enabled)

Uterine Lie (can be more than one selection)

1. Anteverted
2. Anteflexed
3. Retroverted
4. Retroflexed
5. Axial
6. Others (please specify) (Free text enabled)

Uterine Size (body + cervix - 3 planes in mm)

1. (Free text) 42x57x18x39mm

Endometrial thickness (sag plane in mm to nearest mm)

1. (Free text) 4mm

Endometrial lesions

1. Not identified
2. Present Polyp
2b-1. No. of polyps (free text)
2b-2. Size of each polyp. (free text)

Adenomyosis

1. No MRI supportive features
2. Supportive MRI features as described
1. Submucosal cysts.
2. Abnormal junctional zone thickening and measurement
i. Anterior (mm)
ii. Fundal (mm)
iii. Posterior (mm)

Presence of an adenomyoma

1. No
2. Yes

Fibroids

1. No
2. Yes
2a. Number of fibroids
2b. Largest fibroids (location and size mm all fibroids >10mm and/or impact on the cavity) - (Free text)
2b. Submucosal fibroids
2b-0. No

REVIEWER INITIALS

SR

2b-1. Yes

2b-1-1. (description, free text)

Left ovary

1. Absent (Branching logic - move to "Right ovary")
2. Present

Left ovary size (3 planes and volume)

1. NN x NN x NN mm 22x16x25
2. Volume (above x 0.52) 46ml.

Left ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter
a. (Free text) ~20
2. N follicles > 9 mm
a. (Free text) 0
3. Dominant follicle
a. Y
b. N

Left ovary position

1. Lateral adnexa Unremarkable
2. High positioning in iliac fossa
3. Tethered/ distorted appearances - (may be multiple options)
3a. Intimate relationship to the lateral uterus
3b. Intimate relationship/ tethering to adjacent bowel.
3c. Tethering to pelvic sidewall
3d. Tethering to the uterosacral ligament

SUBJECT ID = RR

3e: Other: (free text)

429

REVIEWER INITIALS

adjacent bowel

3c: Tethering to pelvic sidewall

3d: Tethering to the uterosacral ligament

Left ovarian endometrioma

1: No
2: Yes

1a: Size: NN in millimetres (mm)

1b: T2 shading (intermediate/low homogeneous T2 signal)

1b-0: No

1b-1: Yes

1c: T1 fat sat hyperintense

1c-0: Absent

1c-1: Present

1d: Internal nodularity, septation or other complexity

1d-1: No

1d-2: Yes

1-d-2-1 (Free text)

Left ovarian lesions/cysts other than suspected endometriomas

1: Not classifiable

2: Lesion features

2a: Unilocular cyst

2b: Unilocular-solid cyst

2c: Multilocular cyst

2d: Multilocular-solid cyst

2e: Solid

3: Vascularity (post gadolinium enhancement)

3a: Present

3b: Absent

4: Fat component (T1/ T2 hyperintense Low signal on fat suppression).

4a: Present.

4b: Absent.

Right ovary

1: Absent (Branching logic - move to "Adnexa")

2: Present ?? looks like a mass.

Right ovary size (3 planes and volume)

1: NN x NN x NN mm 84x59x60mm.

2: Volume (above x 0.52) 154ml.

Right ovary follicle measurements and count

1: N follicles between 2 and 9 mm in diameter

a: (Free text) 0

2: N follicles > 9 mm

a: (Free text) 0

3: Dominant follicle

a: Y
b: N

Right ovary position

1: Lateral adnexa Unremarkable

2: High positioning in iliac fossa

3: Tethered/ distorted appearances - may be multiple options.

3a: Intimate relationship to the lateral uterus

3b: Intimate relationship/ tethering to

Right ovarian endometrioma

1: No
2: Yes

2a: Size: NN in millimetres (mm)

2b: T2 shading (intermediate/low homogeneous T2 signal)

2b-0: No

2b-1: Yes

2c: T1 fat sat hyperintense

2c-0: Absent

2c-1: Present

2d: Internal nodularity, septation, complex

2d-1: No

2d-2: Yes

Right ovarian lesions/cysts other than suspected endometriomas

1: Not classifiable

2: Lesion features

2a: Unilocular cyst

2b: Unilocular-solid cyst

2c: Multilocular cyst

2d: Multilocular-solid cyst

2e: Solid

3: Vascularity (post gadolinium enhancement).

3a: Present

N/A

SUBJECT ID = RR

429

3b Absent

4 Fat component (T1/ T2 hyperintense Low signal on fat suppression).

4a Present

4b Absent

N/A (we don't have those sequences)

Adnexa

1 Hydrosalpinx

1a No

1b Yes

2 Hematosalpinx

2a No

2b Yes

3 Other (free text)

Are both ovaries immediately approximated "kissing"?

1 No

2 Yes

Uterine bladder nodule

Definition: Is there presence of a nodule in the bladder.

1 Absent

2 Present

2a Size: N/A in millimetres (mm)

Uterovesical region

Definition: Assessment of whether there is a visible preserved fat plane +/- physiologic fluid +/- absent

distortion between the anterior uterine serosa and bladder

1 Normal

2 Abnormal.

2a (free text if required)

Uteric nodule(s)?

1 Absent

2 Present

2a Location (free text + distance to uterine orifice/ VUJ)

2b Size (mm)

Pouch of Douglas obliteration

Definition: Assessment for abnormal loss of fat plane +/- physiologic fluid and immediate approximation of rectosigmoid and/or small bowel to the posterior uterine serosa, cervix +/- vaginal wall.

Discrete linear bands may be visible (↓ T1, ↓ T2)

1 Negative

2 Partial

2a Left

2b Right

3 Complete

3a Positive = obliteration

3b Positive = band adhesions

Nodules present on the posterior vaginal fornix?

Definition: Thickening of superior 1/3 of posterior vaginal wall +/- nodularity. Nodules ↓ T2 ↑ T1 (if active haemorrhagic deposits)

1 No

2 Yes

REVIEWER INITIALS

CM

2a Dimension of nodule to be measured in millimetres (mm)

2b1 Inactive

2b2 Active

Vaginal forniceal elevation?

Definition: Upper level of fornix on sagittal view is superior to the angle of the uterine isthmus with stretching of vaginal wall, and/or acute angulation of the fornix.

1 No

2 Yes

2a Left

2b Right

2c Left and Right

Rectovaginal nodules present?

Definition: Presence of deep infiltrating endometriosis in the anterior rectal wall and posterior vaginal fornix located below the peritoneum of the Pouch of Douglas. Inactive/ fibrotic disease characterised as ↓ T1 ↓ T2 signal

Active disease as ↑ T1, ↑ to intermediate- T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits)

1 No

2 Yes

2a Size (mm)

2b1 Inactive

2b2 Active

SUBJECT ID = RR

429

Uterosacral ligament nodules or thickening?

Definition: Inactive/ fibrotic disease nodules characterised as \downarrow T1 \downarrow T2 signal

Active disease as \uparrow T1, \uparrow to intermediate T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits)

1 No

2 Yes nodules

2a Left

2a-1 Size (mm)

2b Right

2b-1 Size (mm)

2c1 Inactive

2c2 Active

3 Yes thickening

3a Left

3b Right

3c Both

Retrocervical nodule present?

Definition: Inactive/ fibrotic disease characterised as \downarrow T1 \downarrow T2 signal

Active disease as \uparrow T1, \uparrow to intermediate T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits)

1 No

2 Yes

2a Size (mm)

2b1 Inactive

2b2 Active

Rectum and colon

Is there bowel deep infiltrating endometriosis seen?

Definition: Inactive/ fibrotic disease characterised as \downarrow T1 \downarrow T2 signal.

Active disease as \uparrow T1, \uparrow to intermediate T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

"Mushroom cap sign" is specific to severe invasive bowel endometriosis and is characterized as a plaque with \downarrow T2 at its 'base' and \uparrow T2 at its 'cap'

1 No

2 Yes

2a Distance from the anal verge

2a-1 Length (mm)

2b Lesion type

2b-1 Isolated lesion

2b-2 Multiple lesions

2b-3 Curved lesion

2b-4 Straight lesion

2c Maximal depth layer of invasion each lesion (muscularis, submucosa, mucosa)

2c-1 Lesion 1 (free text)

(2c-2 Lesion 2 (free text) - delete if not relevant

(2c-3 etc.)

2c Is it stuck to any structures or free lying?

2d-1 Vagina

2d-2 Uterus

2d-3 Uterosacral ligaments

REVIEWER INITIALS

2d-4 Ovary

2d Plaque thickness

2a 1-5mm

2b 6-10mm

2c >11mm

2e Activity

2f1 Inactive

2f2 Active

2f "Mushroom cap" appearance

2g1 Present

2g2 Absent

Is there evidence of tethering of the bowel?

1 No

2 Yes, tethered to

2a Uterus

2b L ovary

2c R ovary

2d L uterosacral ligament nodule

2e R uterosacral ligament nodule

2f L pelvic side wall

2g R pelvic side wall

2h Other

Any other salient findings on the study:

1 No

2 Yes

a (Free text)

Scan/Photo/ Email: kate.cook@bensonradiology.com.au

Page 4 of 4

suspected ovarian mass; difficult to determine if degenerating fibroid & the nature of the mass as ~~self~~ limited sequences for the study.