SUBJECT ID = RR

**ENDOMETRIOSIS PELVIC MRI ASSESSMENT -BR PROFORMA REPORT BLIND REVIEW** 

#### Uterus

1:

Absent

Present

Uterine anatomy

Conventional

Arcuate

3. Septate

Full septum

Subseptate

Bicornuate unicollis

Bicornuate bicollis

Didelphys

Other (free text enabled).

Uterine Lie (can be more than one selection)

Anteverted

Anteflexed

Retroverted

Retroflexed

Axia

Others (please specify) (Free text enabled)

Uterine Size (body + cervix - 3 planes in mm)

1. (Free text).



Endometrial thickness (sag plane in mm to nearest mm)

(Free text)

**Endometrial lesions** 

Not identified.

Present. Polyp.

2b-1: No. of polyps (free text)

Size of each polyp. (free text) 2b-2:

Adenomyosis

No MRI supportive features

Supportive MRI features as described:

Submucosal cysts.

Abnormal junctional zone thickening and

measurement

Anterior (mm)

Fundal (mm)

Posterior (mm)

Presence of an adenomyoma

1: No

2: Yes

**Fibroids** 

No

2: Yes

> Number of fibroids: 2a:

2b: Largest fibroids (location and size mm all

fibroids >10mm and/or iimpact on the cavity) - (Free text)

2b: Submucosal fibroids

> 2b-0: No

**REVIEWER INITIALS** 

Yes 2b-1:

2b-1-1: (description: free text)

Left ovary

1: Absent (Branching logic - move to "Right ovary")

Present 2:

Left ovary size (3 planes and volume)

1. NN x NN x NN mm 57 X 45 X 45

Volume (above x 0.52).

Left ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter

a. (Free text)

2. N follicles > 9 mm

a. (Free text)

Dominant follicle

Left ovary position

1: Lateral adnexa. Unremarkable.

2: High positioning in iliac fossa.

Tethered/ distorted appearances - (may be

multiple options)

3a: Intimate relationship to the lateral uterus

3b Intimate relationship/ tethering to adjacent

bowel.

3c. Tethering to pelvic sidewall

3d:Tethering to the uterosacral ligament

SUBJECT ID = RR 3e: Other: (free text)

#### Left ovarian endometrioma

No

Yes

Size: NN in millimetres (mm) 1a:

> T2 shading (intermediate/low homogeneous T2 signal).

No 1b-0:

1b-1: Yes

T1 fat sat hyperintense 1c:

> 1c-0: Absent 1c-1: Present

Internal nodularity, septation or other 1d:

complexity.

1d-1: No 1d-2:

Yes

1-d-2-1: (Free text)

# Left ovarian lesions/cysts other than suspected endometriomas

1: Not classifiable

2: Lesion features

> Unilocular cyst 2a:

Unilocular-solid cyst 2b:

Multilocular cyst 2c/

Multilocular-solid cyst 2d:

Solid 2e:

Vascularity (post gadolinium enhancement).

3a: Present 3b: Absent

Fat component (T1/T2 hyperintense. Low signal on fat suppression).

> 4a: Present.

4b: Absent.

#### Right ovary

Absent (Branching logic - move to "Adnexa") Present

Right ovary size (3 planes and volume)

1. NN x NN x NN mm

2. Volume (above x 0.52).

#### Right ovary follicle measurements and count

1. N follicles between 2 and 9 mm in diameter

a. (Free text)

2. N follicles > 9 mm

a. (Free text)

Dominant follicle

# Right ovary position

Lateral adnexa. Unremarkable.

2: High positioning in iliac fossa.

3: Tethered/distorted appearances - may be multiple options.

3a: Intimate relationship to the lateral uterus

3b Intimate relationship/ tethering to

# **REVIEWER INITIALS**

adjacent bowel.

3c. Tethering to pelvic sidewall

3d: Tethering to the uterosacral ligament

## Right ovarian endometrioma

No

2: Yes

> 2a: Size: NN in millimetres (mm)

2b: T2 shading (intermediate/low homogeneous T2 signal).

> 2b-0: No

2b-1: Yes

T1 fat sat hyperintense 2c:

> 2c-0: Absent

2c-1: Present

2d: Internal nodularity, septation, complex.

> 2d-1: No

Yes 2d-2:

# Right ovarian lesions/cysts other than suspected

# endometriomas

Not classifiable

Lesion features

Unilocular cyst 2a:

2b: Unilocular-solid cyst

2€: Multilocular cyst

2d: Multilocular-solid cyst

2e: Solid

Vascularity (post gadolinium enhancement).

3a: Present

#### SUBJECT ID = RR

3b: Absent

Fat component (T1/T2 hyperintense. Low signal on fat suppression).

> 4a: Present.

4b: Absent.

#### Adnexa

Hydrosalpinx 1:

Yes

2: Hematosalpinx No

> 2b: Yes

3: Other (free text).

# Are both ovaries immediately approximated "kissing"?

No

2:

### **Urinary bladder nodule**

Definition: Is there presence of a nodule in the bladder.

Absent

2: Present

> 2a: Size: NN in millimetres (mm)

## **Uterovesical region**

Definition: Assessment of whether there is a visible preserved fat plane +/- physiologic fluid +/- absent

distortion between the anterior uterine serosa and bladder.

Normal.

2: Abnormal.

> 2a: (free text if required)

# Ureteric nodule(s)?

Absent

2: Present

> 2a: Location (free text + distance to ureteric

> > orifice/VUJ)

2b: Size (mm)

#### **Pouch of Douglas obliteration**

Definition: Assessment for abnormal loss of fat plane +/physiologic fluid and immediate approximation of rectosigmoid and/or small bowel to the posterior uterine serosa, cervix +/- vaginal wall.

Discrete linear bands may be visible ( $\sqrt{T1}$ ,  $\sqrt{T2}$ )

Negative

**Partial** 

2a: Left 2b: Right

Complete

3:

3a: Positive = obliteration.

3b: Positive = band adhesions.

## Nodules present on the posterior vaginal fornix?

Definition: Thickening of superior 1/3 of posterior vaginal wall +/- nodularity. Nodules: ↓ T2 ↑T1 (if active haemorrhagic deposits)

No 1:

Yes

## **REVIEWER INITIALS**

Dimension of nodule to be measured in 2a: millimetres (mm).

> 2b1: Inactive.

2b2: Active

#### Vaginal forniceal elevation?

Definition: Upper level of fornix on sagittal view is superior to the angle of the uterine isthmus with stretching of vaginal wall, and/or acute angulation of the fornix.

No

2: Yes

2a: Left.

2b: Right

2c: Left and Right.

#### Rectovaginal nodules present?

Definition: Presence of deep infiltrating endometriosis in the anterior rectal wall and posterior vaginal fornix, located below the peritoneum of the Pouch of Douglas. Inactive/ fibrotic disease characterised as  $\downarrow$  T1  $\downarrow$  T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal (hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes

> 2a: Size (mm)

2b1: Inactive.

2b2: Active

### SUBJECT ID = RR

# Uterosacral ligament nodules or thickening?

Definition: Inactive/ fibrotic disease nodules characterised as  $\downarrow$  T1  $\downarrow$  T2 signal.

Active disease as T1, T to intermediate- T2 signal

(hemorrhagic/ proteinaceous content + glandular deposits).

1: No

2: Yes nodules

2a: Left

2a-1: Size (mm)

2b: Right

2b-1: Size (mm)

2c1: Inactive.

2c2: Active

3: Yes thickening.

3a:

3b: Right

Left.

3c: Both.

### Retrocervical nodule present?

Definition: Inactive/ fibrotic disease characterised as  $\downarrow$  T1  $\downarrow$  T2 signal.

Active disease as ↑T1, ↑ to intermediate- T2 signal

themorrhagic/ proteinacous content + glandular deposits).

.: / No

Yes

2a: Size (mm)

2b1: Inactive.

2b2: Active

#### Rectum and colon:

#### Is there bowel deep infiltrating endometriosis seen?

Definition: Inactive/ fibrotic disease characterised as  $\downarrow$  T1  $\downarrow$  T2 signal.

Active disease as  $\uparrow$ T1,  $\uparrow$  to intermediate- T2 signal (hemorrhagic/ proteinacous content + glandular deposits). "Mushroom cap sign" is specific to severe invasive bowel endometriosis and is characterized as a plaque with  $\downarrow$  T2 at its 'base' and  $\uparrow$  T2 at its 'cap'.

1:

Yes

No

2a: Distance from the anal verge

2a-1: Length (mm)

2b: Lesion type

2b-1: Isolated lesion

2b-2: Multiple lesions

2b-3: Curved lesion

2b-4: Straight lesion

2c: Maximal depth layer of invasion each

leasion (muscularis, submucosa,

2c-1: Lesion 1: (free text)

(2c-2: Lesion 2 (free text) - delete if not relevant

(2c-3 etc.)

mucosa).

2c: Is it stuck to any structures or free lying?

2d-1: Vagina

2d-2: Uterus

2d-3: Uterosacral ligaments

## **REVIEWER INITIALS**

2d-4: Ovary

2d: Plaque thickness

2a: 1-5mm.

2b: 6-10mm.

2c: >11mm.

2e: Activity

2f1: Inactive.

2f2: Active.

2f: "Mushroom cap" appearance:

2g1: Present.

2g2: Absent.

## Is there evidence of tethering of the bowel?

/ No

2: Yes, tethered to

2a: Uterus

2b: L. ovary

2c: R. ovary

2d: L. uterosacral ligament nodule

2e: R. uterosacral ligament nodule

2f: L pelvic side wall.

2g: R pelvic side wall.

2h: Other.

# Any other salient findings on the study:

1. No

2. Yes

a. (Free text).

Scan/ Photo/ Emaii: kate.cook@bensonradiology.com.au