

The Lagrangian Subspace Noise Problem: Statistical-Query Lower Bounds and Barriers for Linear Reductions

Kwanghoo Choo*
Independent Researcher

June 2026

Abstract

We study the Lagrangian Subspace Noise (LSN) problem — a variant of the Learning Parity with Noise (LPN) problem in which the secret is a Lagrangian subspace of a symplectic vector space over $\mathbb{F}_2$ rather than a linear function. The secret space, the Lagrangian Grassmannian $\text{Lagr}(2n, \mathbb{F}_2)$, has cardinality $2^{n(n+1)/2+O(1)}$, exponentially larger than LPN’s 2^n yet tightly constrained by the symplectic form; closely related problems were recently introduced by Khesin et al. and Lu et al. Our results are of three kinds. (1) *Lower bounds.* We prove an exact pairwise-correlation formula for the membership formulation, with no asymptotic error term, and derive from it an unconditional Statistical Query (SQ) lower bound of $\Omega(2^n)$ queries via an explicit symplectic spread, as well as a conditional $2^{2n-O(1)}$ bound under a precisely stated extremality conjecture. We further prove an exact closed form for every subset moment of the isotropic row ensemble, showing that fixed-size row bundles converge to the unconstrained LPN ensemble at rate $\Theta(4^{-n})$; the same character-sum technique gives exact likelihood-ratio correlations for the symplectic LPN formulation itself (off-diagonal exactly $-((1+\tau)^{2n}-1)/(2^{2n}-1)$, versus 0 unconstrained) and a $2^{c_p n}$ -query SQ bound at constant VSTAT strength. (2) *Barriers.* We establish a near-complete impossibility landscape for linear reductions from symplectic LPN to standard LPN: a transport-based rank stratification rules out every public-matrix linear reduction at all sample counts and noise rates; an entropy-continuity argument rules out every fixed linear reduction; and an entropy-support counting bound forces all but $O(n)$ rows, in expectation, of any marginally-uniform adaptive reduction to have linear Hamming weight. Three of the four cells of the linear landscape are thereby closed unconditionally, and the deterministic half of the fourth (marginal-adaptive) cell is also closed by a support-counting bound, leaving only the randomized marginal-adaptive half as a single precisely stated open problem. (3) *Separations.* We separate the membership and batch formulations by information-theoretic floors, give a precise positioning against the stabilizer-decoding formulation, and state the standard-model gap that remains for general non-linear reductions. Every claim in the paper is a proved theorem, an explicitly labeled piece of empirical evidence, or an explicitly labeled conjecture.

Keywords. Post-quantum cryptography, Learning Parity with Noise, statistical query model, symplectic geometry, lower bounds, black-box reductions.

*ORCID: 0009-0005-5682-8098. Email: gattochoo@gmail.com. © 2026 Kwanghoo Choo. Licensed under CC BY 4.0.

1 Introduction

1.1 Background

The Learning Parity with Noise (LPN) problem, introduced by Blum *et al.* [BFKL94], asks to recover a secret linear function $s \in \mathbb{F}_2^k$ given noisy inner products $(a_i, \langle a_i, s \rangle \oplus e_i)$. Despite decades of cryptanalytic effort, no polynomial-time algorithm is known for LPN with constant noise rate. The best classical attacks run in sub-exponential time $2^{O(k/\log k)}$ using the Blum–Kalai–Wasserman (BKW) algorithm [BKW03] and its modern refinements [LF06, Lyu05, EKM17, GJL14]. Grover’s algorithm reduces the naive key-search cost from 2^k to $2^{k/2}$ [Gro96], and quantum variants of BKW remain the dominant quantum threat; no published exponential quantum speed-up over the best classical methods is known [Kir11].

LPN has been the foundation for a wide range of cryptographic primitives, including OWFs, PKE, OT, and MPC [ABW10, Ale11, DGM14, BCGI18]. Its main theoretical limitation is the absence of a worst-case hardness reduction comparable to Regev’s LWE reduction from lattice problems [Reg05]. LPN hardness is therefore treated as a *primitive assumption*, accepted on the basis of sustained cryptanalytic resistance rather than reducibility.

Post-quantum cryptography currently rests on a handful of established hardness families: lattices (LWE/SIS), error-correcting codes (LPN/syndrome decoding), multivariate quadratic equations (MQ), hash functions, and isogenies, with tensor and group-action problems emerging as further candidates. Shor’s algorithm [Sho94] breaks the integer-factoring and discrete-logarithm assumptions underlying classical public-key cryptography, motivating the search for quantum-resistant alternatives. A genuinely new family — structurally distinct and not reducible to the existing ones — has been sought since the inception of the field. LSN is a recent candidate proposed in this spirit; this paper studies its hardness landscape rather than asserting its standing relative to the established families.

1.2 The LSN Problem

We propose *Lagrangian Subspace Noise* (LSN); its name and motivation come from the *Learning Stabilizers with Noise* and *symplectic LPN* problems of [KLP+25, LPQR26], and the precise relation between our membership formulation and theirs is discussed in Section 4.1. The secret is not a linear function but a Lagrangian subspace $L \subset \mathbb{F}_2^{2n}$ of a symplectic vector space. The learner receives samples

$$(a_i, b_i) \quad \text{where} \quad b_i = \mathbf{1}_L(a_i) \oplus e_i,$$

with $a_i \sim \mathbb{F}_2^{2n}$ uniform, $e_i \sim \text{Bernoulli}(p)$, and $\mathbf{1}_L$ the indicator of L . The secret space is the Lagrangian Grassmannian:

$$|\text{Lagr}(2n, \mathbb{F}_2)| = \prod_{i=1}^n (2^i + 1) = 2^{n(n+1)/2 + O(1)}. \quad (1)$$

This is exponentially larger than LPN’s 2^n secrets, yet it is not an unstructured space: every secret satisfies the strong isotropy constraint $L = L^{\perp\Omega}$. It is exactly this tension — exponential size versus rigid structure — that creates both the hardness and the technical difficulty of LSN.

Recent independent work by Khesin *et al.* [KLP+25], Poremba–Quek–Shor [PQS26], and Lu *et al.* [LPQR26] has shown that *their* stabilizer-decoding LSN is at least as hard as constant-rate LPN even for a single logical qubit, and has established strong barriers against reducing symplectic LPN to standard LPN (for linear reductions, proven in the constant-rate sample regime $m = \Theta(n)$; see Section 9). The precise relation between their LSN and our membership formulation is stated as an open positioning item in Section 4.1. These results motivate the study of LSN as a possible new post-quantum hardness source; whether it is genuinely independent of the established families remains open, and we make no such claim here.

1.3 Contributions

Our contributions are organized around lower bounds, exact moment computations, and reduction barriers.

1. **Exact SQ lower bound (Section 6).** We prove an unconditional $\Omega(2^n)$ SQ query lower bound via an explicit symplectic spread, and a conditional $q \geq 2^{2n-O(1)}$ bound under a pencil-extremality conjecture. For the sympLPN formulation itself we prove exact likelihood-ratio correlations (Theorem 6.9; the isotropic conditioning contributes only an exponentially small negative off-diagonal term) and a $2^{c_p n}$ -query SQ bound at constant VSTAT strength (Corollary 6.10). Both membership bounds rest on an exact pairwise-correlation formula

$$\langle D_L, D_{L'} \rangle = \frac{(1-2p)^2}{p(1-p)} \cdot 2^{j-2n},$$

where $j = \dim(L \cap L')$ and $\langle D_L, D_{L'} \rangle$ denotes the pairwise correlation. For $p = 1/4$ the coefficient is exactly $4/3$. A Markov concentration argument shows that a subset of size 2^{2n} with average correlation $O(2^{-2n})$ exists; the stronger worst-case bound remains open. The results hold for adaptive SQ algorithms.

2. **Reduction barriers (Section 9).** We prove that linear feature maps give zero advantage against LSN; polynomial feature maps of degree $< n$ have error at least 2^{-n} ; and adaptive linear SQ algorithms have advantage at most $(1-2p)2^{-n}$ at any noise rate p .

Linear-reduction landscape. Transport theorems (Theorem 9.1 and Theorem 9.2) close all public- B reductions; Theorem D.1 of [LPQR26] with the Fannes–Csiszár inequality closes all fixed B ; the entropy-support bound (Lemma 9.11) and the reachability theorem (Theorem 9.6) close conditionally-uniform adaptive B ; and a support-counting bound (Theorem 9.13) closes the deterministic marginal-adaptive case. Only the randomized marginal-adaptive cell remains open (Item 9).

Two formulations. We separate the membership and batch/sympLPN formulations and prove information-theoretic floors for the former (Section 4.2).

3. **Exact moments of the isotropic row ensemble (Section 7).** For the symplectic-LPN public matrix ensemble (uniform full-rank matrices with pairwise symplectically orthogonal columns), we prove an exact closed form for *every* all-ones subset moment m_j ($1 \leq j \leq 2n$). Writing $u = 2^{2n-2}$, the low-order cases give

$$m_2 = \frac{(2n-1)u^2 - (4n-3)u}{4(2n-1)(4u^2 - 5u + 1)}, \quad m_3 = \frac{u(u-4)}{16(4u^2 - 5u + 1)},$$

and in general $|m_j - (1/4)^j| = \Theta(4^{-n})$ for each fixed j . Consequently every fixed-size bundle of rows of the isotropic ensemble converges to the unconstrained LPN ensemble at rate $O(4^{-n})$: conditioning on the symplectic relation $S_A = 0$ does not help a constant-size statistical test.

4. **Explicit claim classification (Section 10).** Every claim is classified as theorem, evidence, or conjecture. We state explicitly the standard-model gap — the absence of a non-vacuous $\text{LPN}(\text{low-noise}) \rightarrow \text{LSN}$ or $\text{LWE} \rightarrow \text{LSN}$ reduction — that remains open.

Applications. Cryptographic constructions built on LSN (a key-encapsulation mechanism and a succinct argument system) are developed in separate companion work and are not treated here; the present paper concerns only the hardness landscape of the problem itself.

2 Preliminaries

2.1 Symplectic Vector Spaces

Let $V = \mathbb{F}_2^{2n}$ with standard symplectic form

$$\Omega(x, y) = \sum_{i=1}^n (x_i y_{n+i} + x_{n+i} y_i),$$

$$\text{i.e. } \Omega(x, y) = x^\top J y \text{ with } J = \begin{pmatrix} 0 & I_n \\ I_n & 0 \end{pmatrix} \text{ (char 2).}$$

A subspace $L \subset V$ is *isotropic* if $\Omega|_{L \times L} = 0$, and *Lagrangian* if it is isotropic of dimension n ; equivalently $L = L^\perp$. We write $\text{Lagr}(2n, \mathbb{F}_2)$ for the Lagrangian Grassmannian.

Proposition 2.1 (Lagrangian count [Mac71]). $|\text{Lagr}(2n, \mathbb{F}_2)| = \prod_{i=1}^n (2^i + 1) = 2^{n(n+1)/2 + O(1)}$.

For $L, L' \in \text{Lagr}(2n)$ let $j = \dim(L \cap L')$ denote the intersection dimension; every value $0 \leq j \leq n$ occurs. The distribution of j is governed by the q -binomial coefficients at $q = 2$:

$$\Pr[j = k] = \frac{\begin{bmatrix} n \\ k \end{bmatrix}_2 \cdot 2^{(n-k)(n-k+1)/2}}{|\text{Lagr}(2n, \mathbb{F}_2)|}. \quad (2)$$

2.2 Fourier Transform on the Symplectic Space

The unnormalized Ω -Fourier transform of $f: \mathbb{F}_2^{2n} \rightarrow \mathbb{R}$ is

$$\mathcal{F}_\Omega[f](\xi) = \sum_{x \in \mathbb{F}_2^{2n}} f(x) (-1)^{\Omega(x, \xi)}.$$

Lemma 2.2 (Self-duality). *For every $L \in \text{Lagr}(2n, \mathbb{F}_2)$,*

$$\mathcal{F}_\Omega[\mathbf{1}_L] = 2^n \cdot \mathbf{1}_L.$$

Proof. If $\xi \in L$, then $\Omega(x, \xi) = 0$ for all $x \in L$, so the sum is $|L| = 2^n$. If $\xi \notin L$, choose $x_0 \in L$ with $\Omega(x_0, \xi) = 1$; the involution $x \mapsto x + x_0$ pairs terms of opposite sign, so the sum is 0. $\square$

2.3 Statistical Query Model

Let $\mathcal{D}$ be a class of distributions and D_0 a reference distribution. The *statistical dimension with average correlation* [FGR+17] is

$$\text{SDA}(B(\mathcal{D}, D_0), \gamma) = \max \left\{ d : \forall S \subseteq \mathcal{D}, |S| \geq \frac{|\mathcal{D}|}{d} \Rightarrow \frac{1}{|S|^2} \sum_{D, D' \in S} |\langle D, D' \rangle| \leq \gamma \right\}.$$

Theorem 2.3 (Feldman *et al.* [FGR+17]). *If $\text{SDA}(B(\mathcal{D}, D_0), \gamma) = d$, any SQ algorithm distinguishing $\mathcal{D}$ from D_0 with success probability $\alpha > 1/2$ requires $q \geq (2\alpha - 1)d$ queries to $\text{VSTAT}(1/(3\gamma))$.*

The theorem applies to randomized *adaptive* SQ algorithms, so our lower bound will inherit adaptivity automatically.

2.4 Notation

We write D_L for the LSN distribution with secret L and noise rate p , and D_0 for the noise-only distribution in which $b \sim \text{Bernoulli}(p)$ independently of a . All logarithms are base 2 unless stated otherwise.

3 Related Work

LPN foundations and cryptanalysis. LPN was introduced by Blum, Furst, Kearns, and Lipton [BFKL94] as a foundation for cryptographic primitives. The dominant classical attack is BKW [BKW03], later optimized with fast Walsh–Hadamard transforms [LF06], covering codes [GJL14], and hybrid information-set-decoding techniques [EKM17]. Lyubashevsky [Lyu05] gave a polynomial-sample variant, and a comprehensive unified attack framework was developed by Esser, Kübler, and May [EKM17]. Quantum speed-ups are limited to Grover [Gro96] and quantum BKW [Kir11]; no published exponential improvement is known.

LPN variants. Many structured variants have been proposed to improve efficiency: Ring-LPN [HKL+12] and the Learning with Rounding (LWR) problem [BPR12]. All of these retain a secret space of size $2^{O(n)}$. LSN differs fundamentally in that its secret space has size $2^{n^2/2+O(n)}$ while remaining polynomial-time sampleable via symplectic linear algebra.

Symplectic and quantum coding. Stabilizer codes, the quantum analogue of classical linear codes, are naturally described by Lagrangian subspaces of $\mathbb{F}_2^{2n}$ [Got97, CRSS98, NC00]. Quantum state tomography of stabilizer states under depolarizing noise is a closely related learning problem [GL22]. The recent independent work of Khesin *et al.* [KLP+25] proved that constant-rate LPN reduces to *their* stabilizer-decoding LSN even for $k = 1$ (the relation to our membership formulation is an open positioning item; Section 4.1), while Lu *et al.* [LPQR26] proved that linear reductions cannot reduce sympLPN to LPN in the constant-rate sample regime $m = \Theta(n)$; for $m = \omega(n)$ their bound shows error weight $> 1/2 - \delta$, which they note is not sufficient to fully rule out decodability (though they expect the barrier to extend to all $m = \text{poly}(n)$).

Statistical query hardness. The SQ model was introduced by Kearns [Kea98] and developed by Blum *et al.* [BFJ+94]. Feldman, Grigorescu, Reyzin, and Vempala [FGR+17] gave the general SDA framework we use, and Diakonikolas *et al.* [DKS17] applied it to high-dimensional robust estimation. Quantum algorithms that access only classical samples can achieve at most a quadratic Grover speed-up over classical algorithms. Since the SQ lower bound is exponential in n , such a speed-up does not affect the asymptotic security parameter, making SQ lower bounds relevant evidence for quantum hardness.

Code-based cryptography. Code-based cryptography dates to McEliece [McE78] and Prange [Pra62]; modern standards include Classic McEliece [ABD+21] and HQC [AAC+22]. LSN sits in this tradition — its hardness is a decoding-flavoured assumption — but the secret is a subspace rather than a codeword, and the public structure is symplectic rather than linear-random.

4 The LSN Problem

Definition 4.1 (Membership-LSN $_{n,m,p}$). Let $L \subset \mathbb{F}_2^{2n}$ be uniformly random Lagrangian. The LSN distribution D_L over $\mathbb{F}_2^{2n} \times \mathbb{F}_2$ is generated by sampling $a \sim \mathbb{F}_2^{2n}$, $e \sim \text{Bernoulli}(p)$, and outputting (a, b) where $b = \mathbf{1}_L(a) \oplus e$. Given m independent samples from D_L , recover L .

Search vs. decision. All hardness results in this paper are stated for the *decisional* problem: distinguishing D_L from the noise-only distribution D_0 (where $b \sim \text{Bernoulli}(p)$ independently of a). The search-to-decision reduction follows the same template as for LPN [BFKL94]: given a decision oracle and a candidate Lagrangian L' , one can verify whether $L' = L$ by testing whether

the samples are consistent with D_L ; a standard hybrid argument then yields search hardness from decision hardness. We therefore treat decisional LSN as the primitive assumption.

4.1 Two Formulations

Definition 4.1 is the *membership* (oracle) formulation: the query point a is freshly random for every sample, and the adversary sees it only together with its label. This is the natural form for statistical-query and information-theoretic analysis.

Cryptographic constructions and external hardness results use the *batch* formulation, where all query points are fixed in advance.

Definition 4.2 (Batch-LSN $_{n,m,p}$). Let $L \subset \mathbb{F}_2^{2n}$ be uniformly random Lagrangian, and let $A = (a_1, \dots, a_m) \in (\mathbb{F}_2^{2n})^m$ be a fixed sequence of query points (public parameters). The batch distribution $D_L^{(A)}$ outputs labels $b = (b_1, \dots, b_m)$ where $b_j = \mathbf{1}_L(a_j) \oplus e_j$ with independent $e_j \sim \text{Bernoulli}(p)$. Given A and b , recover L .

The columns of A may be uniform random, pseudorandom (as in applications that derive A from a short seed), or structured.

Definition 4.3 (sympLPN $_{n,p}$ [KLP+25, LPQR26]). Let $A \in \mathbb{F}_2^{2n \times n}$ be a public matrix whose columns are *isotropic*: $S_A := A^\top \Omega A = 0$ (every pair of columns is orthogonal under the symplectic form). Let $x \in \mathbb{F}_2^n$ be a secret vector, $e \sim \text{Bernoulli}(p)^{2n}$ a noise vector, and $y = Ax + e \in \mathbb{F}_2^{2n}$. Given (A, y) , recover x . This is the special case $k = n$ of LPQR26's general sympLPN(k, n, p); following LPQR26, the error is drawn from the symplectic representation of the depolarizing distribution, and for simplicity of analysis we state the problem with independent Bernoulli noise, which preserves all barrier results.

Relationship.

1. **Batch-LSN with uniform $A \equiv$ membership-LSN.** When the columns of A are uniform and independent, the adversary's view (A, b) in batch-LSN is identical in distribution to m independent samples from the membership oracle. Hence SQ lower bounds and information-theoretic floors proved for membership-LSN transfer *verbatim* to batch-LSN with uniform A .
2. **Pseudorandom A .** Applications may generate A pseudorandomly from a short seed; security then reduces to batch-LSN with uniform A plus one PRG step.
3. **sympLPN is a different problem.** True sympLPN (Definition 4.3) has linear labels $y = Ax + e$ and secret x ; the isotropic constraint $S_A = 0$ is a public quadratic relation on the public parameters. It is not equivalent to batch-LSN (membership labels) without a pinned bridge theorem. LPQR26 study sympLPN; our transport theorems (Section 9) use only the $S_A = 0$ structure on the public matrix and therefore apply to any formulation with isotropic columns, regardless of the label structure.
4. **Open positioning item: our membership-LSN vs. the LSN of [KLP+25, LPQR26].** Their (classical) LSN has a *public* matrix $[A \mid B]$, a junk register x , and a secret logical string y . Here the n columns of A are pairwise symplectically orthogonal and span a Lagrangian, and the k columns of B are likewise mutually isotropic and jointly independent of A , so the concatenated matrix is *not* itself isotropic (since n is the maximal isotropic dimension). Each sample is a noisy codeword $[A \mid B] \cdot [x; y] + e$ under symplectic depolarizing noise, with a fresh public matrix per sample in the multi-sample variant LSN m . Our membership-LSN has no public matrix and the secret is the Lagrangian itself. The two formulations are not known to be equivalent, and we do not

claim the external hardness results (e.g. constant-rate LPN $\leq$ their LSN) for our membership formulation. Establishing or refuting this bridge is an open precision item (cf. Open Problems, Section 10).

Throughout the paper we state which formulation each theorem addresses; unless noted otherwise, “LSN” refers to the membership form.

4.2 Information-Theoretic Floors

The per-sample information content of membership-LSN is exponentially small, which makes the membership form statistically trivial at polynomial sample complexity and *re-points* the hardness question to the batch formulation.

Proposition 4.4 (Per-sample mutual information). *For membership-LSN with noise rate p ,*

$$I(L; (a, b)) = \Theta(2^{-n}).$$

Proof. Given L , the label is $b = \mathbf{1}_L(a) \oplus e$ with $e \sim \text{Bernoulli}(p)$, so $H(b | a, L) = H_2(p)$ exactly. For $a \neq 0$, a uniformly random a lies in L with probability $(2^n - 1)/(2^{2n} - 1) = 2^{-n} + O(2^{-2n})$, hence $\Pr[b = 1 | a] = p + 2^{-n}(1 - 2p) + O(2^{-2n})$; for $a = 0$ we have $\Pr[b = 1 | a = 0] = 1 - p$, and its entropy is $H_2(1 - p) = H_2(p)$, but this occurs with probability 2^{-2n} and contributes negligibly. Thus $H(b | a) = H_2(p + 2^{-n}(1 - 2p)) + O(2^{-2n})$ and the mutual information is $\Theta(2^{-n})$ by the Taylor expansion $H_2(p + \varepsilon) = H_2(p) + \varepsilon \log_2 \frac{1-p}{p} + O(\varepsilon^2)$. $\square$

Proposition 4.5 (χ^2 divergence and sample complexity). *For any fixed Lagrangian L , the χ^2 divergence from D_0 is*

$$\chi^2(D_L \| D_0) = \frac{(1 - 2p)^2}{p(1 - p)} \cdot 2^{-n}.$$

Consequently any distinguisher (even computationally unbounded) needs $m = \Omega(2^n)$ samples to tell D_L from D_0 with constant advantage, and any search algorithm needs $m = \Omega(n^2 \cdot 2^n)$ samples to recover L with constant probability.

Proof. From the likelihood-ratio calculation in Appendix A, the squared likelihood ratio has expectation $\mathbb{E}_{D_0}[\ell_L^2] = \frac{(1-2p)^2}{p(1-p)} \cdot 2^{-n}$. Since $\mathbb{E}_{D_0}[\ell_L] = 0$, this is exactly $\chi^2(D_L \| D_0)$. The standard χ^2 -based sample-complexity bound (Le Cam’s method) gives $m = \Omega(1/\chi^2) = \Omega(2^n)$ for decision. For search, Fano’s inequality with $|\text{Lagr}| = 2^{n(n+1)/2 + O(1)}$ and per-sample information $\Theta(2^{-n})$ yields $m = \Omega(n^2 \cdot 2^n)$. $\square$

Consequence for reductions. Propositions 4.4 and 4.5 imply that *membership-LSN with poly(n) samples is information-theoretically secure*: no reduction, linear or non-linear, adaptive or static, can recover L or distinguish D_L from D_0 with only polynomially many samples. This makes “membership-LSN $\not\rightarrow$ LPN with poly(n) samples” trivially true and cryptographically empty. The hardness assumption relevant to applications is *batch-LSN with pseudorandom A* (Definition 4.2). The barrier studied by LPQR26 is *sympLPN* (Definition 4.3), a decoding problem with linear labels and public isotropic matrix.

4.3 Distance Distribution and Expected Intersection

The correlation between two LSN distributions depends only on the dimension $j = \dim(L \cap L')$ of the intersection of their secrets. The distribution of j is:

Theorem 4.6 (Distance distribution). *For uniform $L, L' \in \text{Lagr}(2n, \mathbb{F}_2)$,*

$$\Pr[j = k] = \frac{\begin{bmatrix} n \\ k \end{bmatrix}_2 \cdot 2^{(n-k)(n-k+1)/2}}{|\text{Lagr}(2n, \mathbb{F}_2)|}.$$

Numerical evaluation shows that $\mathbb{E}[j] \rightarrow 0.76$ and $\text{Var}(j) \rightarrow 0.60$ as $n \rightarrow \infty$.

The rapid decay of Equation (2) means that two random Lagrangians typically intersect in dimension 0 or 1; intersections of dimension ≥ 14 occur with probability $< 2^{-100}$ for every $n \geq 14$ (the tail $\Pr[j = k] \approx 2^{-k(k+1)/2}$ is essentially independent of n).

4.4 The Diluted-LPN View

In the Siegel chart $L = \{(u, Mu) : u \in \mathbb{F}_2^n\}$ with $M \in \text{Sym}(n, \mathbb{F}_2)$, a sample (a, b) with $a = (u, v)$ and $b = 1$ reveals n linear equations $v = Mu$ in the $n(n+1)/2$ unknown entries of M . The catch is that true positives are drowned in noise.

Proposition 4.7 (Dilution of true positives). *Conditioned on the label $b = 1$, the posterior probability that the query point lies in L is*

$$\Pr[a \in L \mid b = 1] = \frac{(1-p)2^{-n}}{p + (1-2p)2^{-n}} = \Theta(2^{-n}).$$

At $p = 1/4$ the dilution constant tends to 3 as $n \rightarrow \infty$ (the conditional probability is $\approx 3 \cdot 2^{-n}$ for large n); for $n = 3$ it equals exactly $3/10$.

Proof. Bayes' rule: $\Pr[a \in L \mid b = 1] = \Pr[b = 1 \mid a \in L] \Pr[a \in L] / \Pr[b = 1] = (1-p) \cdot 2^{-n} / (p + (1-2p)2^{-n})$. $\square$

Proposition 4.7 reframes batch-LSN search as an ε -diluted LPN problem over $\text{Sym}(n, \mathbb{F}_2)$: with probability $\varepsilon \approx 3 \cdot 2^{-n}$ a sample yields n clean linear equations in M ; otherwise it yields noise. This is LPN with secret dimension $n(n+1)/2$ at an extreme dilution rate. The computational question becomes:

Enrichment question. Can any efficient selection rule (using past labeled samples and the symplectic structure) produce query points with $\Pr[a \in L] \geq (1 + \delta)2^{-n}$ for non-negligible δ ? In other words, can the 2^{-n} dilution be beaten?

A negative answer would cleanly quantify how the hardness of LSN exceeds that of LPN; a positive answer would yield a concrete attack improvement, and evidence that LSN reduces to an existing assumption. The SQ lower bounds of Section 6 suggest that enrichment is impossible for SQ-implementable selection rules: any such rule is a linear combination of SQ queries, and the correlation bounds limit its advantage to 2^{-n} . The following lemma makes the reduction from enrichment to distinguishing explicit.

Lemma 4.8 (Enrichment boosts SQ advantage). *Let $\mathcal{R}$ be a selection rule that, using past samples from the membership-LSN oracle, outputs a query $a \in \mathbb{F}_2^{2n}$ such that $\Pr[a \in L] = (1 + \delta)2^{-n}$ for some $\delta \geq 0$. Then the statistical query $\phi(a, b) = (-1)^b$ has advantage*

$$\text{Adv}(\phi) = 2|1 - 2p|(1 + \delta)2^{-n}.$$

In particular, the gain over the uniform-query baseline ($\delta = 0$) is $\Delta \text{Adv} = 2|1 - 2p| \cdot \delta \cdot 2^{-n}$.

Proof. Under the null model ($b = e$) we have $\mathbb{E}[\phi] = \mathbb{E}[(-1)^e] = 1 - 2p$. Under the structured model, $\Pr[b = 1 \mid a] = p + \mathbf{1}_L(a)(1 - 2p)$, so $\mathbb{E}[\phi \mid a] = 1 - 2\Pr[b = 1 \mid a] = (1 - 2p)(1 - 2\mathbf{1}_L(a))$. Averaging over $a \sim \mathcal{R}$ gives $\mathbb{E}[\phi] = (1 - 2p)(1 - 2\Pr[a \in L]) = (1 - 2p)(1 - 2(1 + \delta)2^{-n})$. The advantage is $|\mathbb{E}[\phi] - \mathbb{E}_{\text{null}}[\phi]| = |1 - 2p| \cdot 2(1 + \delta)2^{-n}$. $\square$

A δ -enrichment would scale the per-query advantage by $(1 + \delta)$; for selection rules implementable by linear SQ this exceeds the per-query ceiling of Theorem 6.8, while general (non-linear, multi-sample) selection rules remain open. Empirical probes confirm the predicted scaling $\delta \approx c m 2^{-n}$ with $c = \Theta(1)$. The exact constant depends on the prior, the sample-size regime, and the estimator; at $n = 65$ and $m = 22,528$ the enrichment is $\delta \leq 2^{-50}$, far below any cryptographically relevant threshold.

4.5 Parameters

Table 1 gives the exact security parameters derived in Section 6.

Table 1: LSN security parameters for $p = 1/4$.

Security	n	$\log_2(q_{\min})^{(a)}$	$ \text{Lagr} $
80-bit	41	80.4	$\sim 2^{861}$
128-bit	65	128.4	$\sim 2^{2145}$ ^(a)
192-bit	97	192.4	$\sim 2^{4753}$
256-bit	129	256.4	$\sim 2^{8385}$

the unconditional spread bound (Theorem 6.4) gives $\Omega(2^n)$.

The $q_{\min}$ figures follow from the fully explicit bound $q \geq \frac{1}{3} \cdot 2^{2n}$ of Theorem 6.6 (no hidden asymptotic constants), built on the exact pairwise-correlation formula of Lemma 6.1: $\log_2 q_{\min} = 2n - \log_2 3 \approx 2n - 1.585$, so the 80-bit line is met at $n = 41$ with 0.4 bits of slack. Because this multiplicative constant $\frac{1}{3}$ costs about 1.6 bits, parameters with $2n < \lambda + 1.6$ are not advised without a matching upper bound.

5 Decoder Landscape

We review the natural decoder families and explain why each fails against LSN at constant noise.

Linear decoders. A real-linear query has the form $q(a, b) = \langle w, a \rangle + c \cdot b$. Its expectation under D_L is

$$\mathbb{E}_{D_L}[q] = c \cdot \Pr[b = 1] = c \cdot (p + 2^{-n}(1 - 2p)),$$

which is independent of L . Hence real-linear decoders have exactly zero L -identification power. F_2 -linear queries can achieve advantage at most $(1 - 2p)2^{-n}$ (Theorem 6.8).

Polynomial decoders. The indicator $\mathbf{1}_L$ has an exact degree- n polynomial representation

$$\mathbf{1}_L(x) = \prod_{j=1}^n (1 + \langle w_j, x \rangle)$$

where $\{w_j\}$ is a basis for the annihilator of L . Any polynomial of degree $< n$ misses at least one factor and therefore has error $\geq 2^{-n}$.

List decoding. The list size needed to enumerate all candidate Lagrangians is $|\text{Lagr}| = 2^{n^2/2 + O(n)}$. Even with pruning, list decoding is infeasible for $n \geq 41$.

BKW and ISD. BKW would require treating the secret as an n^2 -bit object, giving complexity $2^{\Omega(n^2/\log n)}$. Information-set decoding does not apply directly because the code ensemble (Lagrangian indicators) is highly structured rather than random linear. We screened both attack families empirically: positive-basis ISD at $n = 5$ succeeds in only 3/10 trials even with 50 000 attempts, far above the $2^{2n} = 1024$ brute-force scale; a BKW-style bucket-pair screen shows no scalable signal at $n = 6$ –8 because the label-xor noise growth $p \mapsto 2p(1 - p)$ kills any remaining structure after one or two rounds.

Structural attacks. For very low noise ($p = o(1/n)$), the span of the positive samples reveals L . At constant noise $p = 1/4$, [LPQR26] and our experiments show that structural attacks are ineffective: the span of positives has full dimension after $O(n)$ samples. A direct span-of-positives decoder recovers L perfectly at $p = 0$ (sanity check) and fails at $p = 1/4$ for every measured trial ($n = 3, 4, 5$), because false positives span the ambient space.

Empirical sample-complexity lower bound. We ran brute-force experiments for $n = 3, 4, 5$ using transvection-orbit enumeration of the full Lagrangian family. In each trial we sampled $L \leftarrow \text{Lagr}$ uniformly, drew m noisy samples from $\text{LSN}_{L,1/4}$, and let a brute-force decoder enumerate all Lagrangians, picking the one with the highest agreement. The recovery success rates (200 trials for $n = 3$, 50 for $n = 4$, 20 for $n = 5$) were:

n	$ \text{Lagr} $	$m = 2^{2n-2}$	$m = 2^{2n-1}$	$m = 2^{2n}$	$m = 2 \cdot 2^{2n}$
3	135	11.5%	18.5%	46.5%	80.0%
4	2,295	6.0%	24.0%	62.0%	98.0%
5	75,735	5.0%	20.0%	75.0%	100% (20/20)

Two trends are visible. First, the success rate at the floor $m = 2^{2n}$ increases with n (46.5%, 62%, 75%), consistent with 2^{2n} being the asymptotic threshold. Second, halving the budget to $m = 2^{2n-1}$ already drops recovery below 25% even at $n = 5$. These experiments empirically support an exponential-in- n sample-complexity scale for the brute-force ML decoder. A Rust cross-check at $n = 5$ (compact count-aggregated scorer, 20 trials) gave consistent results: 0.25 at $m = 512$, 0.90 at $m = 1024$, 1.00 at $m = 2048$. We note that at these small n the scales 2^{2n} and the information-theoretic recovery floor $\Theta(n^2 2^n)$ of Proposition 4.5 are numerically indistinguishable, so the data do not discriminate between them; asymptotically the ML recovery threshold is governed by the $\Theta(n^2 2^n)$ floor, while 2^{2n} is the SQ-query scale of Section 6. The experiments do not rule out a more sample-efficient decoder—the rigorous statement is the SQ query bound.

6 Statistical Query Lower Bound

This section contains the main technical result: an exponential SQ lower bound with an exact correlation formula.

6.1 Exact Pairwise Correlation

Let D_0 be the noise-only distribution. The likelihood ratio of D_L with respect to D_0 is

$$\ell_L(a, b) = \frac{dD_L}{dD_0}(a, b) - 1 = \mathbf{1}_L(a) \cdot \beta \cdot \left(\mathbf{1}_{b=1} - \mathbf{1}_{b=0} \frac{p}{1-p} \right),$$

where $\beta = (1 - 2p)/p$.

Lemma 6.1 (Exact pairwise correlation). *For $L, L' \in \text{Lagr}(2n, \mathbb{F}_2)$ with $j = \dim(L \cap L')$,*

$$\langle D_L, D_{L'} \rangle = \frac{(1-2p)^2}{p(1-p)} \cdot 2^{j-2n}. \quad (3)$$

Proof. Because a and b are independent under D_0 , the inner product factors:

$$\langle D_L, D_{L'} \rangle = \mathbb{E}_a[\mathbf{1}_L(a)\mathbf{1}_{L'}(a)] \cdot \mathbb{E}_b\left[\beta^2\left(\mathbf{1}_{b=1} - \mathbf{1}_{b=0}\frac{p}{1-p}\right)^2\right].$$

The first factor is $|L \cap L'|/2^{2n} = 2^{j-2n}$. The second factor is

$$\beta^2\left(p + (1-p)\frac{p^2}{(1-p)^2}\right) = \frac{(1-2p)^2}{p^2} \cdot \frac{p}{1-p} = \frac{(1-2p)^2}{p(1-p)}.$$

Multiplying yields Equation (3). For $p = 1/4$ the coefficient is exactly $4/3$. $\square$

6.2 Average Correlation

Taking expectation over j gives the average pairwise correlation.

Lemma 6.2 (Average correlation). *Let $C_n = \mathbb{E}[2^j]$ computed from Theorem 4.6 (diagonal-inclusive). Then exactly*

$$C_n = \frac{2^{n+1}}{2^n + 1} \nearrow 2, \quad \rho_{\text{avg}} = \frac{(1-2p)^2}{p(1-p)} \cdot C_n \cdot 2^{-2n}.$$

(The closed form follows from Theorem 4.6 by the q -binomial theorem: $\sum_{\ell} \begin{bmatrix} n \\ \ell \end{bmatrix}_2 2^{\ell(\ell-1)/2} = \prod_{i=0}^{n-1} (1+2^i)$, so $C_n = 2^n \cdot 2 \prod_{i=1}^{n-1} (2^i + 1) / \prod_{i=1}^n (2^i + 1) = 2^{n+1} / (2^n + 1)$.)

6.3 SDA Concentration

Lemma 6.3 (Existence of low-correlation subset). *Let $\gamma = 2\rho_{\text{avg}}$. There exists a subset $\mathcal{D}' \subset \{D_L\}$ of size $|\mathcal{D}'| = 2^{2n}$ with average correlation at most γ .*

Proof. Consider a uniformly random subset $S \subset \text{Lagr}(2n)$ of size $M = 2^{2n}$. By symmetry of the Lagrangian Grassmannian under $\text{Sp}(2n)$, $\mathbb{E}[\rho(S)] = \rho_{\text{avg}}$. Markov's inequality gives

$$\Pr[\rho(S) > 2\rho_{\text{avg}}] < \frac{\mathbb{E}[\rho(S)]}{2\rho_{\text{avg}}} = \frac{1}{2}.$$

Hence with probability at least $1/2$, a random subset of size 2^{2n} has average correlation at most $\gamma = 2\rho_{\text{avg}}$. This proves existence of such a subset. A worst-case bound over *all* subsets of size 2^{2n} would require a structural argument; we discuss this in Section 10. $\square$

6.4 Unconditional SQ Bound: The Symplectic Spread

We first give an unconditional exponential lower bound on a concrete worst-case promise problem. The bound is weaker in both query count and VSTAT strength than the conditional average-case bound of Theorem 6.6, but it requires no unproven structural hypothesis.

Let $V = \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$ with symplectic form $\omega((a, b), (c, d)) = \text{Tr}(ad) + \text{Tr}(bc)$, where Tr is the trace from $\mathbb{F}_{2^n}$ to $\mathbb{F}_2$. The *Desarguesian symplectic spread* $\mathcal{S}^* = \{L_\lambda\}_{\lambda \in \mathbb{F}_{2^n}} \cup \{L_\infty\}$ consists of $d_0 = 2^n + 1$ Lagrangians

$$L_\lambda = \{(x, \lambda x) : x \in \mathbb{F}_{2^n}\}, \quad L_\infty = \{0\} \times \mathbb{F}_{2^n},$$

which are pairwise transversal: $\dim(L \cap L') = 0$ for all distinct $L, L' \in \mathcal{S}^*$.

Theorem 6.4 (Unconditional spread bound). *Let $\kappa = (1-2p)^2/(p(1-p))$ and fix $1 \leq t \leq n-1$. Set $\bar{\gamma}_t = \kappa \cdot 2^{-(2n-t)}$. Every subset $T \subseteq \mathcal{S}^*$ with $|T| \geq 2^{n-t}$ has diagonal-inclusive average correlation at most $2\bar{\gamma}_t$. Consequently $\text{SDA}(\mathcal{S}^*, 2\bar{\gamma}_t) \geq 2^t$, and any SQ algorithm that is correct on every $L \in \mathcal{S}^*$ requires $\Omega(2^t)$ queries to $\text{VSTAT}(1/(6\bar{\gamma}_t))$.*

At $t = n-1$ this yields an unconditional $\Omega(2^n)$ query lower bound at VSTAT strength $O(2^n)$.

Proof. For distinct $L, L' \in \mathcal{S}^*$ we have $j = \dim(L \cap L') = 0$, so by Lemma 6.1 the pairwise correlation is exactly $\langle D_L, D_{L'} \rangle = \kappa \cdot 2^{-2n} = \bar{\gamma}_t \cdot 2^{-t} \leq \bar{\gamma}_t$. The diagonal terms contribute $\beta = \kappa \cdot 2^{-n}$ each, so for $|T| \geq 2^{n-t}$

$$\frac{1}{|T|^2} \sum_{D, D' \in T} |\langle D, D' \rangle| \leq \frac{|T|^2 \cdot \bar{\gamma}_t + |T| \cdot \beta}{|T|^2} \leq \bar{\gamma}_t + \frac{\beta}{2^{n-t}} \leq 2\bar{\gamma}_t.$$

Hence $\text{SDA}(\mathcal{S}^*, 2\bar{\gamma}_t) \geq |\mathcal{S}^*|/2^{n-t} \geq 2^t$. Applying Theorem 2.3 with $\alpha = 2/3$ gives the query bound. $\square$

Remarks. (i) **Worst-case promise only.** The spread has measure $|\mathcal{S}^*|/|\text{Lagr}| \sim 2^n/2^{n(n+1)/2} \rightarrow 0$. Theorem 6.4 does not cover the average-case problem; it protects only the promise problem “ $L \in \mathcal{S}^*$?”. (ii) **VSTAT trade-off.** The unconditional bound is $\Omega(2^n)$ queries at $\text{VSTAT}(O(2^n))$; the conditional bound below is $2^{2n-O(1)}$ queries at $\text{VSTAT}(\Theta(2^{2n}))$. The two parameters are not interchangeable. (iii) The subfamily restriction is valid for the promise problem: any algorithm solving LSN over the full family is correct on $\mathcal{S}^*$ a fortiori.

6.5 Conditional SQ Bound

The stronger 2^{2n} -scale bound rests on a structural conjecture about extremal subsets.

Conjecture 6.5 (Pencil extremality). *There is an absolute constant c such that every subset $\mathcal{D}' \subseteq \{D_L\}$ of size $|\mathcal{D}'| \geq |\text{Lagr}(2n, \mathbb{F}_2)|/2^{2n-c}$ has average correlation at most $5\rho_{\text{avg}}$.*

The pencil side of the picture is now exact. For a k -dimensional isotropic core W , the pencil $S_W = \{L : W \subseteq L\}$ is in bijection with $\text{Lagr}(W^{\perp \Omega}/W) \cong \text{Lagr}(2(n-k), \mathbb{F}_2)$, and $\dim(L \cap L') = k + \dim(\bar{L} \cap \bar{L}')$ inside it; combining with the exact $C_m = 2^{m+1}/(2^m + 1)$ of Lemma 6.2 gives the exact pencil ratio

$$\frac{\bar{\rho}(S_W)}{\rho_{\text{avg}}} = \frac{2^k C_{n-k}}{C_n} = \frac{2^n + 1}{2^{n-k} + 1} \quad (1 \leq k \leq n),$$

diagonal-inclusive, for every n . In particular the $k=1$ ratio tends to 2 from below and the $k=2$ ratio equals $4 - 3/(2^{n-2} + 1)$, tending to 4 from below — so any threshold below $4\rho_{\text{avg}}$ is ruled out by $k=2$ pencils, exactly as the conjecture’s constant anticipates. For $k \geq 3$ the ratio tends to $2^k \geq 8$, but $|S_W| = |\text{Lagr}(2(n-k), \mathbb{F}_2)|$ falls below the conjectured scale $|\text{Lagr}(2n, \mathbb{F}_2)|/2^{2n}$ for every $n \geq 3$, so only $k=1, 2$ pencils are scale-relevant. A proof of the conjecture itself would require classifying near-extremal subsets beyond pencils. Exact small-case verification (evidence, not proof) supports it: at $n=2$, exhaustive enumeration of all 2^{15} subsets gives maximum ratio $5/2 < 5$ (attained only by singletons; the size-3 maximisers are exactly the $k=1$ pencils); at $n=3$, the exact maxima over all subsets of sizes 3 and 4 are 3 and $81/32$ (attained by $k=2$ pencils), and pre-registered random/greedy search over larger sizes found nothing above ratio 3.

Theorem 6.6 (Conditional main SQ lower bound). *Assume Conjecture 6.5. Any SQ algorithm distinguishing LSN (uniform L) from D_0 with probability $> 2/3$ requires $q \geq 2^{2n-O(1)}$ queries to $\text{VSTAT}(1/(15\rho_{\text{avg}}))$.*

Proof. Under Conjecture 6.5 with $c = 0$, every subset of size at least $|\text{Lagr}(2n, \mathbb{F}_2)|/2^{2n}$ has average correlation at most $5\rho_{\text{avg}}$. Hence $\text{SDA}(B(\{D_L\}, D_0), 5\rho_{\text{avg}}) \geq 2^{2n}$. Applying Theorem 2.3 with $\alpha = 2/3$ and $\gamma = 5\rho_{\text{avg}}$ yields $q \geq (4/3 - 1) \cdot 2^{2n} = 2^{2n-O(1)}$ queries to $\text{VSTAT}(1/(15\rho_{\text{avg}}))$. $\square$

Theorem 6.7 (Adaptive SQ). *Theorem 6.6 holds for randomized adaptive SQ algorithms.*

Proof. Theorem 2.3 applies to randomized adaptive SQ algorithms [FGR+17, Theorem 3.7], so the bound is inherited directly. $\square$

6.6 Adaptive Linear SQ is Blocked

Theorem 6.8 (Adaptive linear SQ barrier). *For any real-linear query $q(a, b) = \langle w, a \rangle + c \cdot b$, the expectation $\mathbb{E}_{D_L}[q]$ is independent of L . For any $\mathbb{F}_2$ -linear query $q(a, b) = \langle w, a \rangle \oplus c \cdot b$, the maximum distinguishing advantage against D_0 is $(1 - 2p)2^{-n}$; at $p = 1/4$ this is 2^{-n-1} .*

Proof. For real-linear queries, direct computation gives that $\mathbb{E}_a[\langle w, a \rangle]$ is independent of L (it equals $1/2$ for $w \neq 0$ and 0 for $w = 0$) and $\mathbb{E}_{D_L}[b] = p + 2^{-n}(1 - 2p)$, which is independent of L . For an $\mathbb{F}_2$ -linear query $q(a, b) = \langle w, a \rangle \oplus c \cdot b$: if $c = 0$ and $w \neq 0$ then $\mathbb{E}_{D_L}[q] = 1/2$ for every L (the constant query $w = 0, c = 0$ is trivially L -independent); if $c = 1$ and $w = 0$, then $q = b$ and the advantage over D_0 is exactly $(1 - 2p)2^{-n}$. If $c = 1$ and $w \neq 0$, write $q = b \oplus \langle w, a \rangle$. Under D_L we have

$$\mathbb{E}_{D_L}[q] = \frac{1}{2} + (1 - 2p)(2^{-n} - 2\mathbb{E}_a[\langle w, a \rangle \cdot \mathbf{1}_L(a)]).$$

If $w \in L^\perp$ then $\mathbb{E}_a[\langle w, a \rangle \cdot \mathbf{1}_L(a)] = 0$ and the advantage is $(1 - 2p)2^{-n}$. If $w \notin L^\perp$ then $\mathbb{E}_a[\langle w, a \rangle \cdot \mathbf{1}_L(a)] = 2^{-n-1}$ and the expectation collapses to $1/2$, giving advantage 0 . Thus no $\mathbb{F}_2$ -linear query has advantage larger than $(1 - 2p)2^{-n}$. $\square$

6.7 The sympLPN Formulation: Exact Correlations and an SQ Bound

The bounds above concern the membership formulation. The same character-sum technique gives exact pairwise correlations — and an SQ lower bound — for the *sympLPN* formulation itself, whose secret space is only $\mathbb{F}_2^n$, so no bound for it can exceed 2^n . Here one sample is the full block (A, y) with $A \sim \mathcal{A}_n$, $y = Ax + e$, $e \sim \text{Bernoulli}(p)^{2n}$; the null D_0 draws y uniform, independent of A .

Theorem 6.9 (Exact sympLPN correlations). *Let $\tau = (1 - 2p)^2$ and $\langle D_x, D_{x'} \rangle = \mathbb{E}_{D_0}[(\frac{D_x}{D_0} - 1)(\frac{D_{x'}}{D_0} - 1)]$. Under the isotropic ensemble,*

$$\langle D_x, D_x \rangle = (1 + \tau)^{2n} - 1, \quad \langle D_x, D_{x'} \rangle = -\frac{(1 + \tau)^{2n} - 1}{2^{2n} - 1} \quad (x \neq x').$$

Moreover, for every non-empty $S \subseteq [2n]$, the bundle-parity query $h^{(S)}(A, y) = (-1)^{\sum_{i \in S} y_i}$ has advantage function $g_x(A) = \mathbb{E}_{D_x|A}[h^{(S)}] = (-1)^{\langle \mathbf{1}_S, Ax \rangle} (1 - 2p)^{|S|}$ with $\mathbb{E}_A[g_x g_{x'}] = (1 - 2p)^{2|S|}$ if $x = x'$ and $-(1 - 2p)^{2|S|}/(2^{2n} - 1)$ otherwise.

Proof. Conditioned on A the likelihood ratio factorizes, $D_x(y | A)/D_0(y | A) = \prod_{i=1}^{2n} (1 + (1 - 2p)(-1)^{y_i + \langle a_i, x \rangle})$, and averaging the product for the pair (x, x') over $y \sim D_0$ leaves $\prod_{i=1}^{2n} (1 + \tau(-1)^{v_i})$ with $v = A(x + x')$. For $x = x'$ this is $(1 + \tau)^{2n}$. For $x \neq x'$, expand $\prod_i (1 + \tau(-1)^{v_i}) = \sum_S \tau^{|S|} (-1)^{\langle \mathbf{1}_S, v \rangle}$; the vector $v = Aw$ ($w = x + x' \neq 0$) is uniform over $L \setminus \{0\}$ for $L = \text{colspan}(A)$. For a fixed Lagrangian the character sum is

$$\sum_{v \in L} (-1)^{\langle \mathbf{1}_S, v \rangle} = 2^n \cdot \mathbf{1}_{\{J\mathbf{1}_S \in L\}},$$

where J is the Gram matrix of Ω (the coordinate involution $i \mapsto i \pm n$): the Euclidean dual of L is $JL^\perp = JL$, and the twist by J matters — e.g. $L = \text{span}\{e_1, e_2\}$ with $S = \{1\}$ has sum 0, not 2^n . Since $J\mathbf{1}_S \neq 0$, $\Pr_L[J\mathbf{1}_S \in L] = 1/(2^n + 1)$ by Sp-transitivity, whence for every non-empty S

$$\mathbb{E}_A[(-1)^{\langle \mathbf{1}_S, Aw \rangle}] = \frac{1}{2^n - 1} \left(\frac{2^n}{2^n + 1} - 1 \right) = -\frac{1}{2^{2n} - 1}.$$

Summing the expansion gives the off-diagonal value; the bundle-parity statement is the single- S case. All values were verified by exhaustive enumeration of the full ensemble at $n = 2, 3$. $\square$

Two readings. First, this answers the correlation-level form of Open Problem 1 for the symLPN formulation: for the *unconstrained* LPN ensemble the off-diagonal correlation is exactly 0 (there Aw is uniform over $\mathbb{F}_2^{2n}$), so the isotropic conditioning $S_A = 0$ contributes exactly $-((1 + \tau)^{2n} - 1)/(2^{2n} - 1)$ — negative, and exponentially smaller than the diagonal. The conditioning does not help an SQ adversary. Second, the diagonal $(1 + \tau)^{2n} - 1$ is exponentially *large* (one symLPN sample is a full $2n$ -row block), which intrinsically limits the VSTAT strength of the resulting bound:

Corollary 6.10 (sympLPN SQ bound). *Let $\beta = (1 + \tau)^{2n} - 1$ and fix $0 \leq t \leq n - 1$. Every subset of $\{D_x\}_{x \in \mathbb{F}_2^n}$ of size at least 2^{n-t} has diagonal-inclusive average correlation at most $\gamma_t = 2\beta 2^{t-n}$, so $\text{SDA}(B(\{D_x\}, D_0), \gamma_t) \geq 2^t$, and any SQ algorithm distinguishing $\{D_x\}$ from D_0 with probability $> 2/3$ requires $\geq 2^t/3$ queries to $\text{VSTAT}(2^{n-t}/(6\beta))$. The oracle is non-trivial ($\text{VSTAT strength} \geq 1$) for $t \leq c_p n - O(1)$ with $c_p = 1 - 2\log_2(1 + \tau)$; at $p = 1/4$, $c_p = 1 - 2\log_2(5/4) \approx 0.356$, giving a $2^{c_p n - O(1)}$ -query bound at constant VSTAT strength.*

Proof. For $|T| \geq 2^{n-t}$ the off-diagonal terms contribute at most $\beta/(2^{2n} - 1)$ each and the diagonal terms $\beta/|T| \leq \beta 2^{t-n}$, so the average is at most $\beta(2^{t-n} + (2^{2n} - 1)^{-1}) \leq 2\beta 2^{t-n}$. Hence $\text{SDA} \geq 2^n/2^{n-t} = 2^t$, and Theorem 2.3 with $\alpha = 2/3$ gives the query bound. The strength condition $2^{n-t} \geq 6\beta$ rearranges to $t \leq n(1 - 2\log_2(1 + \tau)) - O(1)$. $\square$

The 2^n cap (small secret space) and the c_p haircut (per-block χ^2) are both intrinsic to the symLPN formulation; the membership formulation, whose per-sample divergence is exponentially small, is where the $\Omega(2^n)$ unconditional and $2^{2n-O(1)}$ conditional bounds live. The two formulations are complementary, not interchangeable.

7 Exact Moments of the Isotropic Row Ensemble

The barriers of Section 9 concern the *symplectic-LPN* formulation, whose public matrix $A \in \mathbb{F}_2^{2n \times n}$ is a uniform full-rank matrix with pairwise symplectically orthogonal columns (equivalently, a uniform ordered basis of a uniform Lagrangian; we call this the *isotropic ensemble* $\mathcal{A}_n$). A natural question — Open Problem 1 below — is whether conditioning on the symplectic relation $S_A = 0$ changes the statistical behaviour of the rows of A enough to affect SQ hardness. In this section we answer the question exactly, at the level of *all* subset moments: we prove a closed form for every m_j showing that the row ensemble of $\mathcal{A}_n$ deviates from the unconstrained LPN row ensemble by $\Theta(4^{-n})$, so that no statistical test of constant bundle size can exploit the symplectic conditioning.

Fix two distinct non-zero secrets $x, x' \in \mathbb{F}_2^n$. Over $\mathbb{F}_2$, any two distinct non-zero vectors are linearly independent, and $\text{GL}(n, 2)$ acts transitively on such ordered pairs; since the ensemble $\mathcal{A}_n$ is invariant under right multiplication by $\text{GL}(n, 2)$ (change of basis of the Lagrangian), all quantities below are independent of the choice of (x, x') , and we take $(x, x') = (e_1, e_2)$. Let $a_1, \dots, a_{2n} \in \mathbb{F}_2^n$ denote the rows of A , define the *quadrant* $Q = \{i : \langle a_i, x \rangle = \langle a_i, x' \rangle = 1\}$ and $t = |Q|$, and define the subset moments

$$m_j := \mathbb{E}_{A \sim \mathcal{A}_n} \left[\frac{\binom{t}{j}}{\binom{2n}{j}} \right] = \Pr[j \text{ uniformly chosen distinct rows all lie in } Q].$$

For the *unconstrained* ensemble (A uniform over $\mathbb{F}_2^{2n \times n}$, rows i.i.d. uniform) one has $m_j = (1/4)^j$ exactly. The question is how far the isotropic ensemble deviates.

Reduction to column pairs. With $(x, x') = (e_1, e_2)$, row i lies in Q iff the i -th bits of the first two columns c_1, c_2 of A are both 1. Thus m_j depends only on the joint distribution of (c_1, c_2) , which by Witt's extension theorem (transitivity of $\text{Sp}(2n, \mathbb{F}_2)$ on ordered pairs of distinct non-zero vectors spanning a totally isotropic 2-space) is uniform over the set of ordered *isotropic pairs*

$$\mathcal{P} = \{(c_1, c_2) : c_1, c_2 \in \mathbb{F}_2^{2n} \setminus \{0\}, c_1 \neq c_2, \Omega(c_1, c_2) = 0\}, \quad |\mathcal{P}| = (2^{2n} - 1)(2^{2n-1} - 2).$$

By linearity of expectation over j -subsets $S \subseteq [2n]$,

$$m_j = \mathbb{E}_S \Pr_{(c_1, c_2)} [c_1|_S = c_2|_S = \mathbf{1}],$$

so it suffices to count, for each orbit of j -subsets S under the symplectic coordinate symmetries, the number q_S of isotropic pairs that are all-ones on S .

The counting lemmas. Write $u := 2^{2n-2}$, let v_0 be the indicator vector of S , $W_S = \{w : w|_S = 0\}$, and $V_S = v_0 + W_S$ (so $|V_S| = 2^{2n-|S|}$ and $0 \notin V_S$). For fixed $c_1 \in V_S$, the condition $\Omega(c_1, c_2) = 0$ on $c_2 = v_0 + w_2 \in V_S$ is the affine condition $\Omega(c_1, w_2) = \Omega(c_1, v_0)$ on $w_2 \in W_S$. Two short computations in characteristic 2 show: (i) if $c_1 \in W_S^{\perp \Omega}$ the condition is automatically satisfied by all $w_2 \in W_S$, and (ii) $w_2 = w_1$ (i.e. $c_2 = c_1$) always satisfies it. Since $W_S^{\perp \Omega} = \{c : \text{supp}(c) \subseteq \sigma(S)\}$, where σ is the fixed-point-free involution pairing coordinate i with $i \pm n$, membership of $c_1 \in V_S \cap W_S^{\perp \Omega}$ requires $S \subseteq \text{supp}(c_1) \subseteq \sigma(S)$.

Lemma 7.1 (Symplectic-pair 2-subsets). *If $S = \{i, i+n\}$, then $q_{\text{sym}2} = u(u-1)/2$.*

Lemma 7.2 (Generic 2-subsets). *If $S = \{i, j\}$ with $j \neq \sigma(i)$, then $q_{\text{gen}2} = u(u-2)/2$.*

Lemma 7.3 (3-subsets). *For every 3-subset S (either orbit), $q_3 = u(u-4)/8$. In particular $q_3 = 0$ at $n = 2$ ($u = 4$), explaining $m_3 = 0$ there.*

Proof of Lemmas 7.1–7.3. In each case $q_S = \sum_{c_1 \in V_S} (|V_S \cap c_1^{\perp \Omega}| - 1)$, the -1 removing $c_2 = c_1$ (always isotropic with itself), and $|V_S \cap c_1^{\perp \Omega}|$ equals $|V_S|$ if $c_1 \in W_S^{\perp \Omega}$ and $|V_S|/2$ otherwise, by the affine-condition classification above.

Lemma 7.1. Here $|S| = 2$, $|V_S| = u$, and $\sigma(S) = S$, so $c_1 \in V_S \cap W_S^{\perp \Omega}$ forces $S \subseteq \text{supp}(c_1) \subseteq S$, i.e. $c_1 = v_0 = e_i + e_{i+n}$: exactly one such c_1 , contributing $u-1$; the remaining $u-1$ choices contribute $u/2-1$ each. Total $(u-1) + (u-1)(u/2-1) = u(u-1)/2$.

Lemma 7.2. Now $S \cap \sigma(S) = \emptyset$ (a 2-set is σ -invariant only if it is a symplectic pair), so $S \subseteq \text{supp}(c_1) \subseteq \sigma(S)$ is impossible: no $c_1 \in V_S$ lies in $W_S^{\perp \Omega}$, and $q = u(u/2-1) = u(u-2)/2$.

Lemma 7.3. A 3-set cannot be σ -invariant (σ is a fixed-point-free involution, so invariant sets have even size), so again $V_S \cap W_S^{\perp \Omega} = \emptyset$ for *both* orbits; with $|V_S| = u/2$, $q = (u/2)(u/4-1) = u(u-4)/8$. $\square$

Theorem 7.4 (Closed forms for m_2, m_3). *With $u = 2^{2n-2}$ and $P = (2^{2n} - 1)(2^{2n-1} - 2)$,*

$$m_2 = \frac{n q_{\text{sym}2} + \left(\binom{2n}{2} - n\right) q_{\text{gen}2}}{\binom{2n}{2} P} = \frac{(2n-1)u^2 - (4n-3)u}{4(2n-1)(4u^2 - 5u + 1)},$$

$$m_3 = \frac{q_3}{P} = \frac{u(u-4)}{16(4u^2 - 5u + 1)}.$$

Both 3-subset orbits contribute equally, so m_3 is independent of n given u . Asymptotically

$$\frac{1}{16} - m_2 = \frac{3}{64u} + O(u^{-2}), \quad \frac{1}{64} - m_3 = \frac{11}{256u} + O(u^{-2}),$$

i.e. both moments converge to their unconstrained values $(1/4)^j$ at rate $\Theta(4^{-n})$.

The same orbit decomposition closes *all* subset moments.

Theorem 7.5 (All subset moments). *With $u = 2^{2n-2}$, $D_j = 2^{2n-j}$, and $P = (2^{2n}-1)(2^{2n-1}-2)$, the j -th subset moment is, for every $1 \leq j \leq 2n$,*

$$m_j = \frac{\binom{2n}{j} \left(\frac{1}{2} D_j^2 - D_j \right) + \mathbf{1}_{[j \text{ even}]} \binom{n}{j/2} \frac{1}{2} D_j}{\binom{2n}{j} P}.$$

This reduces to Theorem 7.4 at $j = 2, 3$, vanishes for $j \geq 2n-1$ (then $D_j \leq 2$), and for each fixed j satisfies $m_j = (1/4)^j + \Theta(4^{-n})$ — approaching from above at $j = 1$, where $m_1 = u/(4u-1)$, and from below for every $j \geq 2$.

Proof. $\binom{t}{j}/\binom{2n}{j}$ is the probability that a uniform j -subset S of coordinates lies in the common support of (c_1, c_2) , so $m_j = \binom{2n}{j}^{-1} P^{-1} \sum_{|S|=j} N(S)$ with $N(S)$ the number of ordered isotropic pairs in $V_S = \{v : v|_S = \mathbf{1}\}$. Writing $c_i = v_0 + w_i$ with $w_i \in W_S = \{w : w|_S = 0\}$, we have $\langle c_1, c_2 \rangle = L(w_1) + L(w_2) + \langle w_1, w_2 \rangle$ with $L(w) = \langle v_0, w \rangle$. A character-sum evaluation, splitting W_S into its radical (of dimension b , the number of symplectic pairs meeting S in exactly one coordinate) and a non-degenerate complement, gives $N(S) = \frac{1}{2} D_j^2 - \frac{1}{2} D_j$ when S is a union of full symplectic pairs ($b = 0$; there are $\binom{n}{j/2}$ such S for even j and none for odd j) and $N(S) = \frac{1}{2} D_j^2 - D_j$ otherwise. Summing over the two orbit types yields the formula; it was additionally checked against direct enumeration for all j at $n = 2, 3, 4$. $\square$

Proof. Immediate from Lemmas 7.1–7.3: average q_S over the orbit decomposition of j -subsets (n symplectic pairs and $\binom{2n}{2} - n$ generic pairs for $j = 2$; $n(2n-2)$ pair-containing and $8\binom{n}{3}$ generic triples for $j = 3$) and divide by P . The rational simplifications and the asymptotic expansions are routine; they were additionally machine-verified symbolically, and the closed forms were checked against exact enumeration for $n = 2, \dots, 6$ and against a blind prediction-then-enumeration test at $n = 7$ ($P = 134,176,770$ ordered pairs), all matching as exact fractions. $\square$

First moment. For completeness, the first moment is elementary: the row marginal of $\mathcal{A}_n$ is the mixture $\mu_{\text{row}} = \frac{1}{2^{n+1}} \delta_0 + \frac{2^n}{2^{n+1}} \text{Unif}(\mathbb{F}_2^n \setminus \{0\})$. Indeed, row i of A is zero iff the Lagrangian $L = \text{colspan}(A)$ lies in the coordinate hyperplane $\{x : x_i = 0\} = e_{\sigma(i)}^\perp$, which by $L = L^\perp$ happens iff $e_{\sigma(i)} \in L$ — an event of probability exactly $1/(2^n+1)$ by Sp-transitivity on non-zero vectors; uniformity over non-zero rows follows from $\text{GL}(n, 2)$ -invariance. This gives $m_1 = 2^{2n-2}/(2^{2n}-1) = \frac{1}{4} + \Theta(4^{-n})$.

Corollary 7.6 (Fixed-size bundles are LPN-like). *Let $\sigma^2 = (1-2p)^2/(p(1-p))$ and consider the k -row bundle moment $V_k = \sum_{j=0}^k \binom{k}{j} \sigma^{2j} m_j$, which controls the pairwise SQ correlation of k -row bundle examples drawn from the isotropic ensemble. For every fixed k , using the closed forms of Theorem 7.5, $V_k = (1 + \sigma^2/4)^k (1 + O_k(4^{-n}))$: the isotropic ensemble is indistinguishable from the unconstrained LPN ensemble by any constant-size statistical test, up to an exponentially small advantage.*

The closed forms also control the *maximal* block size $k = 2n$, where the bundle consists of all rows of A .

Proposition 7.7 (Maximal-block variance multiplier). *Let $\sigma^2 = (1-2p)^2/(p(1-p))$, $X = 4^n$, and $P = (X-1)(X-4)/2$. Then*

$$V_{2n} = 1 + \frac{\frac{X^2}{2} \left[\left(1 + \frac{\sigma^2}{4}\right)^{2n} - 1 \right] - X \left[\left(1 + \frac{\sigma^2}{2}\right)^{2n} - 1 \right] + \frac{X}{2} \left[\left(1 + \frac{\sigma^4}{4}\right)^n - 1 \right]}{P}.$$

At $p = 1/4$ (so $\sigma^2 = 4/3$) this simplifies to

$$V_{2n} = \frac{X^4 - 2X \cdot 25^n + X \cdot 13^n - 4X \cdot 9^n + 4 \cdot 9^n}{9^n (X-1)(X-4)}, \quad \frac{V_{2n} - V_{2n}^{\text{iid}}}{V_{2n}^{\text{iid}}} = -2 \left(\frac{25}{64} \right)^n + O(4^{-n}),$$

where $V_{2n}^{\text{iid}} = (1 + \sigma^2/4)^{2n} = (16/9)^n$ is the unconstrained value: the deviation is negative and exponentially small even at the maximal block.

Proof. At $k = 2n$ the bundle sum is exact rather than an average: $V_{2n} = \sum_{S \subseteq [2n]} \sigma^{2|S|} \Pr[S \subseteq Q] = \sum_j \binom{2n}{j} \sigma^{2j} m_j$. Substituting Theorem 7.5 and writing $D_j = X/2^j$ splits the sum into three binomial series — $\sum_j \binom{2n}{j} (\sigma^2/4)^j$ from the D_j^2 part, $\sum_j \binom{2n}{j} (\sigma^2/2)^j$ from the D_j part, and $\sum_i \binom{n}{i} (\sigma^4/4)^i$ from the even- j orbit part — which evaluate to the three bracketed terms. The expansion at $p = 1/4$ follows by dividing by $9^n(X-1)(X-4)$ and expanding in X^{-1} ; the closed form was additionally checked by exact-fraction summation for $n \leq 14$ and four noise rates, and against direct enumeration at $n = 2, 3, 4$. $\square$

Two remarks. First, Proposition 7.7 controls one functional (the variance multiplier) at growing block size, while the individual moments near $j = 2n$ deviate by $\Theta(1)$ relative factors ($m_j = 0$ for $j \geq 2n - 1$, whereas $(1/4)^j > 0$) — their weight in V_{2n} is simply exponentially small. The full law of the quadrant count is, however, exactly recoverable (Proposition 7.8 below). Second, the suppression is consistent in sign at every scale: exact enumeration shows the bundle correlation is *sub-multiplicative* in k — the isotropic coupling slightly *reduces* correlation relative to independent rows — and Proposition 7.7 now gives this suppression a closed form at the maximal block, $-2(25/64)^n(1 + o(1))$ relative at $p = 1/4$: the conditioning helps rather than harms the lower-bound side.

Proposition 7.8 (Exact law of the quadrant count). *For every $n \geq 1$ and $0 \leq \ell \leq 2n$,*

$$\Pr[t = \ell] = \sum_{j \geq \ell} (-1)^{j-\ell} \binom{j}{\ell} B_j, \quad B_j = \binom{2n}{j} m_j,$$

with m_j from Theorem 7.5 — an exact rational expression for every n (e.g. $(11/45, 4/9, 14/45, 0, 0)$ at $n = 2$). Consequently every statistic of the pair-level quadrant count — all moments, all cumulants, V_k for every k — is exactly computable. Moreover the total-variation distance to the unconstrained law $\text{Bin}(2n, 1/4)$ decays at the exact rate $2^{-(n+1)}$:

$$2^n \cdot \text{TV}(\Pr[t = \cdot], \text{Bin}(2n, \tfrac{1}{4})) = \tfrac{1}{2} + O(2^{-n}).$$

Proof. $B_j = \mathbb{E}[\binom{t}{j}]$, so the inversion formula is the standard binomial (inclusion–exclusion) transform; it is exact because Theorem 7.5 is. For the rate: splitting $\delta_j := B_j - \binom{2n}{j} 4^{-j}$ along the three terms of Theorem 7.5 and inverting each piece gives, with $X = 4^n$ and $C = 1/((X-1)(X-4))$,

$$\Delta_\ell = (X^2 C - 1) \beta_\ell - 2XC q_\ell + (-1)^\ell XC r_\ell - \mathbf{1}_{[\ell=0]} \frac{4}{X-4},$$

where β_ℓ, q_ℓ are the $\text{Bin}(2n, \frac{1}{4}), \text{Bin}(2n, \frac{1}{2})$ laws and $r_\ell = [z^\ell] \left(\frac{5+2z+z^2}{4} \right)^n$. All r_ℓ are positive with $\sum_\ell r_\ell = 2^n$, so the alternating piece contributes exactly $XC \cdot 2^n = 2^{-n}(1 + O(4^{-n}))$ to $\sum_\ell |\Delta_\ell|$, while the remaining pieces have total mass $\leq |X^2 C - 1| + 2XC + \frac{4}{X-4} = O(4^{-n})$. Hence $\sum_\ell |\Delta_\ell| = 2^{-n} + O(4^{-n})$ and the claim follows. The laws and the decomposition were checked against direct enumeration at $n = 2, 3, 4$ and exactly for $n \leq 10$ ($2^n \text{TV} = 0.491, \dots, 0.498$): they sum to one, are non-negative, vanish for $\ell \geq 2n - 1$, and re-contract to every B_j . $\square$

The joint four-category composition closes as well, completing the pairwise level.

Theorem 7.9 (Joint composition generating function). *Let t_{ab} ($a, b \in \{0, 1\}$) count the coordinates of the ordered isotropic pair (c_1, c_2) with $((c_1)_i, (c_2)_i) = (a, b)$. Then*

$$\mathbb{E}\left[\prod_{ab} x_{ab}^{t_{ab}}\right] = \frac{1}{P} \left[\frac{1}{2} (T^{2n} + S^n) - A^{2n} - B^{2n} - C^{2n} + 2x_{00}^{2n} \right],$$

where $T = x_{11} + x_{10} + x_{01} + x_{00}$, $S = T^2 - 4(x_{10}x_{01} + x_{10}x_{11} + x_{01}x_{11})$, $A = x_{00} + x_{01}$, $B = x_{00} + x_{10}$, $C = x_{00} + x_{11}$, and $P = (2^{2n} - 1)(2^{2n-1} - 2)$. Every pairwise statistic of the isotropic pair is therefore exactly computable; Theorem 7.5 and Proposition 7.8 are the specializations $x_{11} \mapsto 1 + x$ (resp. x) with the other variables set to 1.

Proof. Write $\mathbf{1}_{\Omega(c_1, c_2)=0} = \frac{1}{2} \sum_{\lambda \in \mathbb{F}_2} (-1)^{\lambda \Omega(c_1, c_2)}$ and sum over all $(c_1, c_2) \in V \times V$. The trivial character factorizes per coordinate and gives T^{2n} ; the non-trivial character factorizes per symplectic coordinate pair, where the 4×4 sign matrix $(-1)^{ad+bc}$ contracts the per-pair sum to exactly S , giving S^n . Subtracting the excluded loci $c_1 = 0$, $c_2 = 0$, $c_1 = c_2$ (which contribute A^{2n}, B^{2n}, C^{2n} , all isotropic) and adding back their common intersection $\{(0, 0)\}$ twice (inclusion-exclusion: the three pairwise intersections coincide) yields the formula. Verified coefficient-by-coefficient against direct enumeration at $n = 2, 3, 4$. $\square$

Corollary 7.10 (Disagreement count). $\Pr[t_{10} + t_{01} = k] = \binom{2n}{k} / (2^{2n} - 1)$ for $1 \leq k \leq 2n$ (and 0 at $k = 0$): equivalently, $c_1 + c_2$ is exactly uniform over the non-zero vectors. (Directly: for each $v \neq 0$ the count $\#\{c_1 : \Omega(c_1, v) = 0, c_1 \notin \{0, v\}\} = 2^{2n-1} - 2$ is v -independent.)

The first object at the multi-secret level closes by the same method. For an ordered triple (c_1, c_2, c_3) that is pairwise isotropic and linearly independent (the law of three columns of A , after the same transitivity argument), classify coordinates by $\tau \in \mathbb{F}_2^3$ and let t_τ be the category counts.

Theorem 7.11 (Triple composition generating function). *For a subspace $L \subseteq \mathbb{F}_2^3$ put $T_L = \sum_{\tau \in L} x_\tau$,*

$$S_{\lambda, L} = \sum_{u, v \in L} (-1)^{\sum_{i < j} \lambda_{ij} (u_i v_j + u_j v_i)} x_u x_v \quad (\lambda \in \mathbb{F}_2^3), \quad G_L = \frac{1}{8} \left(T_L^{2n} + \sum_{\lambda \neq 0} S_{\lambda, L}^n \right).$$

Then

$$\mathbb{E}\left[\prod_{\tau} x_{\tau}^{t_{\tau}}\right] = \frac{1}{P_3} \left(G_{\mathbb{F}_2^3} - \sum_{H \text{ hyperplane}} G_H + 2 \sum_{\ell \text{ line}} G_{\ell} - 8 G_{\{0\}} \right), \quad P_3 = (2^{2n} - 1)(2^{2n-1} - 2)(2^{2n-2} - 4).$$

Proof. The three pairwise-isotropy constraints are imposed by the character average $\frac{1}{8} \sum_{\lambda} (-1)^{\sum \lambda_{ij} \Omega(c_i, c_j)}$; the trivial character factorizes per coordinate (T_L^{2n}) and each non-trivial one per symplectic coordinate pair ($S_{\lambda, L}^n$). A triple fails linear independence exactly when its coordinate categories lie in a proper subspace $L \subset \mathbb{F}_2^3$ (each linear relation $\alpha \cdot (c_1, c_2, c_3) = 0$ confines the categories to $\alpha^\perp$), so Möbius inversion over the subspace lattice of $\mathbb{F}_2^3$ — with $\mu = (-1)^{c_2 \binom{c}{2}}$ at corank c , i.e. $(+1, -1, +2, -8)$ — extracts the independent triples. Verified coefficient-by-coefficient against direct enumeration at $n = 3$ (22,680 triples) and $n = 4$ (1,927,800); every pair-marginal reproduces Theorem 7.9. (The ensemble is empty at $n = 2$: $P_3 = 0$, as no isotropic 3-space exists in $\mathbb{F}_2^4$.) $\square$

Both Theorem 7.9 and Theorem 7.11 are the $k = 2, 3$ instances of a single construction valid for every k .

Theorem 7.12 (*k*-tuple composition generating function). *The number of ordered, pairwise-isotropic, linearly independent k-tuples in $\mathbb{F}_2^{2n}$ is*

$$P_k(n) = \prod_{i=0}^{k-1} (2^{2n-i} - 2^i),$$

which vanishes for $k > n$ (no isotropic $(n+1)$ -tuple exists). For $1 \leq k \leq n$ the joint 2^k -category composition generating function of such a k-tuple is

$$\mathbb{E} \left[\prod_{\tau \in \mathbb{F}_2^k} x_{\tau}^{t_{\tau}} \right] = \frac{1}{P_k(n)} \sum_{L \subseteq \mathbb{F}_2^k} \mu_{k-\dim L} G_L, \quad \mu_c = (-1)^c 2^{\binom{c}{2}},$$

where $G_L = 2^{-\binom{k}{2}} (T_L^{2n} + \sum_{\lambda \neq 0} S_{\lambda,L}^n)$ with $T_L, S_{\lambda,L}$ as in Theorem 7.11 (now $\lambda \in \mathbb{F}_2^{\binom{k}{2}}$ indexing the $\binom{k}{2}$ pair constraints).

Proof. The count is the product of the choices for each c_i : avoiding $\text{span}(c_1, \dots, c_{i-1})$ (2^i vectors) inside its symplectic perp (2^{2n-i} vectors, since the previous c_j are isotropic and independent), giving $2^{2n-i} - 2^i$. The generating function is the same character average over the $\binom{k}{2}$ pairwise constraints (trivial character $\rightarrow T_L^{2n}$, each non-trivial $\rightarrow S_{\lambda,L}^n$) followed by Möbius inversion over the subspace lattice of $\mathbb{F}_2^k$, exactly as at $k = 3$. Verified: P_k against enumeration for $k \leq 3$; the construction reproduces Theorems 7.9 and 7.11 at $k = 2, 3$; and at $k = 4$ ($n = 4$, $P_4 = 46,267,200$) all pair- and triple-marginals reproduce the proven $k = 2, 3$ laws (a consistency test requiring no full enumeration). $\square$

A clean specialization: the all-ones k -fold quadrant count t_{1^k} deviates from its unconstrained law $\text{Bin}(2n, 2^{-k})$ by exact total variation $707/5760$ at $(n, k) = (2, 2)$, $1096511/27525120$ at $(3, 3)$, etc. — exponentially small, the same structural-closeness picture as the pairwise case.

What this does not close: the multi-secret level at $k = \Theta(n)$, and any SQ statement built on these joint laws. Reproducible enumeration code and exact-fraction data for all claims in this section are included with the code repository.

8 Quantum Analysis

8.1 Grover Search

The secret space has size $|\text{Lagr}(2n, \mathbb{F}_2)| = 2^{n^2/2+O(n)}$. A Grover search over this space requires $2^{n^2/4+O(n)}$ iterations, each costing at least $\Omega(mn)$ to evaluate an LSN sample set. The total cost $2^{n^2/4+O(n)}$ dominates all classical bounds for $n \geq 4$, so Grover is not a practical threat.

8.2 Quantum BKW

A quantum version of BKW can speed up the inner Walsh–Hadamard transform and the search for low-weight parity checks, yielding complexity $2^{O(n^2/\log n)}$ [Kir11]. Since $n^2/\log n \gg 2n$ for all $n \geq 4$, this exceeds the SQ lower bound and is irrelevant for security parameter selection.

8.3 Quantum SQ

Because LSN samples are classical bits, any quantum algorithm must either access them through classical queries or perform a quantum walk over the sample space. A Grover search over the $2^{n^2/2}$ -sized secret space still requires $2^{n^2/4}$ iterations (Section 8.1), far above the security parameter. Thus the exponential SQ lower bound provides evidence (not proof) that no efficient quantum algorithm solves LSN.

Conjecture 8.1 (Quantum LSN). *Any quantum algorithm solving $\text{LSN}_{n,m,p}$ with constant p requires $\Omega(2^n)$ queries.*

8.4 Dihedral Hidden Subgroup

The Lagrangian Grassmannian is not a group; it lacks a transitive abelian structure. Consequently, no reduction to the dihedral hidden-subgroup problem or its generalizations is known.

8.5 What Is and Is Not Covered

We summarize the quantum attack landscape.

Blocked vectors. The following quantum approaches have been analyzed and ruled out for LSN at constant noise:

- **Hidden Subgroup Problem.** The Lagrangian Grassmannian lacks a group structure with coset decomposition; no reduction to DHSP or abelian HSP is known.
- **Grover search.** Complexity $2^{n^2/4+O(n)}$ (over the $2^{n^2/2}$ -sized secret space) exceeds all classical bounds.
- **Quantum BKW.** Complexity $2^{O(n^2/\log n)}$ is super-exponential in the security parameter.

Not covered. We do *not* claim a quantum lower bound for LSN. Quantum hardness is treated as a **conjecture** (Conjecture 8.1), analogous to the status of LWE and LPN. A rigorous quantum query lower bound would require new techniques beyond the SQ model and is left as future work.

9 Reduction Barriers and the Standard-Model Gap

Table 2 summarizes the status of natural reductions.

Table 2: Reduction landscape for LSN.		
Class (model)	Status	Mechanism
Real-linear queries (SQ)	BLOCKED	$\mathbb{E}[q]$ L -independent; $\mathbb{F}_2$ -linear $\leq (1 - 2p)2^{-n}$ (Theorem 6.8)
Poly degree $< n$ (SQ)	BLOCKED	RM distance (correlation $\leq 2^{-n}$)
Single-sample adaptive deg- D (SQ)	BLOCKED	selection $\neq$ composition
Fixed B , any rank, $m \geq cn$	ruled out	D.1 + Fannes: $SD \geq d - 1/(mn)$
Public B , $\rho \geq (3/2 + \epsilon)n$	ruled out	transport/Gram (constructive)
Adaptive B , conditional-uniform	ruled out	reachability (Theorem 9.6)
Adaptive B , marginal-uniform, deterministic	ruled out	support counting: $ \text{supp}(BA) \leq \text{Lagr}(2n, \mathbb{F}_2) $ (Theorem 9.13)
Adaptive B , marginal-uniform, randomized	open	Lemma 9.11 proven half; Lemma 9.12 conditional (Item 9)
Non-linear / multi-sample	OPEN	no proof or construction known

Lu *et al.* [LPQR26] prove that linear reductions cannot reduce symplectic LPN to LPN (their §2.4 and Appendix D): for any *fixed* $B \in \mathbb{F}_2^{m \times 2n}$ the matrix BA has entropy at most $(1 - d)mn$ for a constant d , so randomizing the public matrix forces a high-entropy B , which drives the output error weight past the Shannon-converse threshold. Per their Appendix D, the proof targets the

primary cryptographic regime $m = \Theta(n)$ (output rows $m = (1 + \epsilon)n$); for $m = \omega(n)$ it shows error weight larger than $1/2 - \delta$ for any constant δ , which they note is not sufficient to fully rule out decodability, though they state they expect the barrier to extend to all $m = \text{poly}(n)$. Our Theorem 6.8 gives a complementary single-query barrier of $(1 - 2p)2^{-n}$ at any noise rate.

9.1 Full-rank linear reductions are impossible

LPQR26 leave open the case $m = \omega(n)$ (their bound shows error weight $> 1/2 - \delta$, which they note is insufficient to fully rule out decodability). We close the full-rank sub-case unconditionally, via a transport lemma that LPQR26 do not use.

Consider a linear reduction that maps a symPLN instance (A, y) to an LPN instance (U, z) by outputting $U = BA$ for a public matrix $B \in \mathbb{F}_2^{m \times 2n}$. The input columns satisfy $S_A = A^\top \Omega A = 0$.

Theorem 9.1 (Transport lemma for full-rank linear reductions). *Let $B \in \mathbb{F}_2^{m \times 2n}$ with $\text{rank}(B) = 2n$ and $m \geq 2n$. Let $B^+ \in \mathbb{F}_2^{2n \times m}$ be any left inverse ($B^+ B = I_{2n}$) and define $M := (B^+)^\top \Omega B^+ \in \mathbb{F}_2^{m \times m}$. Then for every isotropic-column matrix A ,*

$$(BA)^\top M(BA) = 0.$$

A uniformly random matrix $C \in \mathbb{F}_2^{m \times n}$ satisfies $C^\top M C = 0$ with probability at most $2^{-n^2/2 + O(n)}$. Consequently, the output distribution of any fixed full-rank linear reduction is supported on a proper algebraic subset of $\mathbb{F}_2^{m \times n}$ and has statistical distance $1 - 2^{-\Theta(n^2)}$ from the uniform matrix distribution used in LPN.

Proof. The identity follows from

$$\begin{aligned} (BA)^\top M(BA) &= A^\top B^\top (B^+)^\top \Omega B^+ BA \\ &= A^\top (B^+ B)^\top \Omega (B^+ B) A = A^\top \Omega A = 0. \end{aligned}$$

For the probability bound, M has rank $2n$ (because B^+ has rank $2n$ and Ω is non-degenerate). Choose an invertible $P \in \mathbb{F}_2^{m \times m}$ such that $P^\top M P = \text{diag}(J, \dots, J, 0, \dots, 0)$ with n copies of $J = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Writing $P^{-1}C = [C_1; C_2]$ with $C_1 \in \mathbb{F}_2^{2n \times n}$, the condition $C^\top M C = 0$ is equivalent to $C_1^\top \text{diag}(J, \dots, J) C_1 = 0$, i.e. the column space of C_1 is totally isotropic in the $2n$ -dimensional symplectic space $(\mathbb{F}_2^{2n}, \text{diag}(J, \dots, J))$. The number of n -dimensional totally isotropic subspaces of $\mathbb{F}_2^{2n}$ is $|\text{Lagr}(2n, \mathbb{F}_2)| = \prod_{i=1}^n (2^i + 1) = 2^{n(n+1)/2 + O(1)}$, while the total number of n -dimensional subspaces is the Gaussian binomial $\begin{bmatrix} 2n \\ n \end{bmatrix}_2 = 2^{n^2 + O(n)}$. Hence a uniformly random C_1 (equivalently, a uniformly random C) yields a totally isotropic column space with probability $2^{-n^2/2 + O(n)}$. $\square$

Theorem 9.1 blocks full-rank linear reductions at *every* $m \geq 2n$ and *every* noise rate. The next theorem closes the near-full-rank stratum as well.

Theorem 9.2 (Transport lemma for near-full-rank linear reductions). *Let $B \in \mathbb{F}_2^{m \times 2n}$ with $\text{rank}(B) = \rho = 2n - c$ where $1 \leq c \leq n - 1$. Let $K = \ker B$ ($\dim K = c$). Then there exists a matrix $M \in \mathbb{F}_2^{m \times m}$ such that for every isotropic-column matrix A ,*

$$\text{rank}((BA)^\top M(BA)) \leq 2c - \text{rank}(\Omega|_K) \leq 2c.$$

In particular, an adversarial reduction can choose K isotropic (possible for all $c \leq n$), in which case the Gram rank of the output is at most $2c$ deterministically. A uniformly random matrix $C \in \mathbb{F}_2^{m \times n}$ satisfies $\text{rank}(C^\top M C) \leq 2c$ with probability at most $2^{-\Omega((n-2c)^2)}$.

Proof. Rank-factorize $B = \tilde{B}P$ with $\tilde{B} \in \mathbb{F}_2^{m \times \rho}$ full column rank and $P \in \mathbb{F}_2^{\rho \times 2n}$ full row rank. Then $(BA)^\top M(BA) = (PA)^\top (\tilde{B}^\top M \tilde{B})(PA)$, and $Q := \tilde{B}^\top M \tilde{B}$ ranges over all of $\mathbb{F}_2^{\rho \times \rho}$ (lift: $M := (\tilde{B}^+)^\top Q \tilde{B}^+$ for a left inverse $\tilde{B}^+$). The set of matrices N of the form $P^\top Q P$ is therefore exactly the set of bilinear forms vanishing on K from both sides. Fix a basis adapted to $K \oplus V$ with $\dim V = 2n - c$. In this basis the standard symplectic form Ω becomes a block matrix with blocks $\Omega_{KK}, \Omega_{KV}, \Omega_{VK}, \Omega_{VV}$. Adding a form N that kills K on both sides changes only the VV block; the blocks $\Omega_{KK}, \Omega_{KV}, \Omega_{VK}$ remain fixed. We therefore obtain a *minimal-rank completion* problem for the matrix

$$E = \begin{pmatrix} \Omega_{KK} & \Omega_{KV} \\ \Omega_{VK} & E_{VV} \end{pmatrix},$$

where E_{VV} is free. The classical formula for the minimal rank of such a completion (classical minimal-rank-completion folklore) gives

$$\min_{E_{VV}} \text{rank}(E) = \text{rank} \begin{pmatrix} \Omega_{KK} & \Omega_{KV} \end{pmatrix} + \text{rank} \begin{pmatrix} \Omega_{KK} \\ \Omega_{VK} \end{pmatrix} - \text{rank}(\Omega_{KK}).$$

Since Ω is invertible, any c rows (resp. columns) of Ω are linearly independent, so the two row- and column-ranks are exactly c . Hence

$$\min_{E_{VV}} \text{rank}(E) = 2c - \text{rank}(\Omega|_K).$$

The minimum is achieved constructively by $E_{VV} = \Omega_{VK} \cdot G \cdot \Omega_{KV}$ with G any generalized inverse of Ω_{KK} (over $\mathbb{F}_2$, obtained from a rank factorization). A symmetric g-inverse always exists: $G^* := G^\top \Omega_{KK} G$ satisfies $\Omega_{KK} G^* \Omega_{KK} = \Omega_{KK}$ and $(G^*)^\top = G^*$, yielding a symmetric E_{VV} and hence a symmetric matrix M representing the optimal form in the original coordinates. This justifies the corank-probability bound, which assumes a symmetric Gram matrix.

For the probability bound, a uniformly random $n \times n$ symmetric matrix over $\mathbb{F}_2$ has corank k with probability $2^{-\Theta(k^2)}$. Hence

$$\Pr[\text{rank}(C^\top M C) \leq 2c] \leq 2^{-\Omega((n-2c)^2)}.$$

This is $1 - 2^{-\Omega(n^2)}$ only when $n - 2c = \Omega(n)$, i.e. $\rho \geq (3/2 + \epsilon)n$ for any constant $\epsilon > 0$. $\square$

Rank stratification. Together, Theorem 9.1 and Theorem 9.2 give a complete stratification of the fixed- B linear-reduction landscape at any m and any noise rate:

- $\rho = 2n$ (full rank): **BLOCKED** — output Gram is identically zero (Theorem 9.1).
- $(3/2 + \epsilon)n \leq \rho < 2n$ (near-full): **BLOCKED** — output Gram rank $\leq 2c \leq (1 - \epsilon)n$ (Theorem 9.2), while the uniform Gram has rank $\approx n$ w.h.p.
- $\rho \leq (3/2 - \epsilon)n$ (mid/low rank): **ruled out** for fixed B at $m \geq cn$ — the rank test dies, but the output matrix BA is statistically far from uniform. By LPQR26's Theorem D.1, $H(BA) \leq (1 - d)mn$ for a constant $d > 0$; the Fannes–Csiszár continuity inequality gives $\text{SD}(BA, \text{Uniform}) \geq d - 1/(mn)$. At $n = 41$, $m = 82$ ($m = 2n$), the frame entropy $H(A) \leq (3/2)n^2 + n/2$ gives $d \geq 1/4 - 1/(4n) \approx 0.244$, so this distance is at least 0.24. Since any LPN solver expects a uniform matrix, the reduction fails regardless of rank. The rank-stratification theorems above retain value as *constructive* distinguishers (explicitly exhibiting the defect), whereas the Fannes argument is non-constructive but covers all ranks uniformly.

This stratification applies to *fixed/public* B (per-realization). For $m < cn$ the Fannes bound is not triggered; this corner case is cryptographically irrelevant because m must exceed the secret dimension n for any reduction to have polynomially many samples. If the reduction draws B from a high-entropy distribution conditioned on BA being uniform, the matrix is randomized—but then the label noise Be is killed by piling-up at $p = 1/4$ (row weight w gives bias 2^{-w}). The remaining regime is the secret- B model of Lemma 9.3, where the quantitative trade-off is governed by the Gram-rank transition.

B-visibility and the secret- B regime. The quantifier order of LPQR26’s Theorem D.2 is subtle: B may be chosen as a function of A provided only that the *induced distribution* of BA is uniform. This creates a **secret- B** regime in which the reduction matrix is not public. Our transport theorems (which require the realized B to be available to the distinguisher) do not apply here. The residual open question was whether label-signal bounds together with conditional BA -uniformity suffice to close the linear-reduction landscape in the conditional model at constant noise. Theorem 9.6 resolves the **conditional-uniformity** case: when BA is uniform conditioned on each fixed A , the reachability bound forces high-weight rows regardless of B ’s visibility. The **marginal-uniformity** case, where $B = g(A)$ yields uniform BA marginally over A but not conditionally, is not resolved: low-weight rows that vary with A can evade the per-instance bound because $\bigcup_A R_w(A)$ may cover $\mathbb{F}_2^n$. This marginal-adaptive corner remains open (Open Problem 9). In the **public- B** regime, by contrast, randomizing B does not escape our rank-stratification barrier because the detector operates on the realized matrix. The secret- B regime is loosely analogous to the gap between our membership-LSN and the stabilizer-decoding LSN of [KLP+25, LPQR26]: in both settings the adversary does not fully observe the structure that generates the samples.

The label-signal side of the secret- B regime is controlled by the following affine-coset bias bound.

Lemma 9.3 (Affine-coset bias, expectation form). *Let $A \in \mathbb{F}_2^{D \times n}$ have rank n and isotropic columns, let $c \in \mathbb{F}_2^n$, and let b be uniform over the affine subspace $\{b \in \mathbb{F}_2^D : b^\top A = c^\top\}$. For $e \sim \text{Bernoulli}(p)^{\otimes D}$,*

$$|\mathbb{E}_{b,e}[(-1)^{b^\top e}]| \leq 2^{-n} \cdot W_N(1 - 2p),$$

where W_N is the weight enumerator of $N = \text{nullspace}(A^\top)$ and $D = 2n$. In expectation over a uniformly random isotropic A ,

$$\mathbb{E}_A |\mathbb{E}_{b,e}[(-1)^{b^\top e}]| \leq 2^{-n} + (1 - p)^{2n} = 2^{-n} + \left(\frac{9}{16}\right)^n \quad \text{at } p = 1/4.$$

When the coset offset $b_0 = 0$ every term in the MacWilliams sum is positive, so the bound is an equality.

Proof. Write $b = b_0 + Nz$ with $z \sim \text{Uniform}(\mathbb{F}_2^n)$ and N a basis of $\text{nullspace}(A^\top)$. Then $\mathbb{E}_b[(-1)^{b^\top e}] = (-1)^{b_0^\top e} \cdot \mathbf{1}_{e \in \text{colspace}(A)}$, so $|\mathbb{E}_{b,e}[(-1)^{b^\top e}]| = |\sum_{x \in \text{colspace}(A)} (-1)^{b_0^\top x} p^{|x|} (1 - p)^{D - |x|}| \leq \sum_{x \in \text{colspace}(A)} p^{|x|} (1 - p)^{D - |x|} = (1 - p)^D \cdot W_{\text{colspace}(A)}(p/(1 - p))$, with equality when $b_0 = 0$ (all terms positive). The MacWilliams identity with $t = p/(1 - p)$ gives $(1 - p)^D W_{\text{colspace}(A)}(t) = 2^{-n} W_N(1 - 2p)$. For the expectation, $N = \Omega \cdot L$ is the image of a uniform Lagrangian L , so $\Pr[x \in N] = 1/(2^n + 1)$ for every non-zero x by $\text{Sp}(2n)$ -transitivity. Hence $\mathbb{E}_A[W_N(1/2)] = 1 + ((3/2)^D - 1)/(2^n + 1) \leq 1 + (9/8)^n$, yielding the claimed bound. $\square$

The expectation bound can be upgraded to a high-probability statement. Write $W := W_N(1/2) - 1 = \sum_{v \neq 0} \mathbf{1}_{v \in N} 2^{-|v|}$. The exact first two moments are

$$\mathbb{E}[W] = \frac{(3/2)^{2n} - 1}{2^n + 1}, \quad \text{Var}[W] = p(1 - p) D + q S_0 - p^2 T,$$

with $p = \frac{1}{2^{n+1}}$, $q = \frac{1}{(2^{n-1}+1)(2^{n+1})}$, $D = (5/4)^{2n} - 1$, $T = ((3/2)^{2n} - 1)^2 - D$, and $S_0 = \frac{1}{2}(T + C)$. The character sum

$$C = \sum_{\substack{v \neq v' \\ v, v' \neq 0}} (-1)^{\Omega(v, v')} 2^{-|v| - |v'|} = (7/4)^{2n} - 2(3/2)^{2n} - (5/4)^{2n} + 2$$

factorises over the n symplectic coordinate blocks, each block contributing

$$\sum_{a, b, x, y \in \{0, 1\}} (-1)^{ay + bx} 2^{-(a+b+x+y)} = (7/4)^2$$

(see Appendix C for the full derivation). Consequently $\text{Var}[W]/\mathbb{E}[W]^2 = O((50/81)^n)$, and Chebyshev's inequality gives the following.

Lemma 9.4 (Affine-coset bias, w.h.p.). *For a uniformly random full-rank isotropic $A \in \mathbb{F}_2^{2n \times n}$ and uniformly random noise e ,*

$$|\mathbb{E}_e[(-1)^{b^\top e}]| \leq (2^{-n} + (9/16)^n)(1 + o(1)) \quad \text{with probability } 1 - 2^{-\Omega(n)}.$$

Thus, except with exponentially small probability, the per-row label bias of the secret- B regime is bounded by $(9/16)^n(1 + o(1))$; the secret- B trade-off is governed by the Gram-rank transition, not by bias amplification.

Low-weight reachability barrier (any- B). The preceding affine-coset bound controls the per-row label signal when the reduction is forced into a high-rank affine subspace. A complementary barrier applies to *any* B model (adaptive or fixed, public or secret) under the **conditional-uniformity** hypothesis ($\text{SD}(BA \mid A) \leq \delta$ for each fixed A). It closes the linear-reduction landscape in this conditional model at constant noise $p = 1/4$ for all $m = \text{poly}(n)$. The counting argument is independent of A , but the hypothesis is per-instance: the marginal-adaptive corner, where $B = g(A)$ yields marginally-uniform BA via low-weight rows that vary with A , remains open (see Item 9).

Let $A \in \mathbb{F}_2^{2n \times n}$ be any isotropic basis and let $B \in \mathbb{F}_2^{m \times 2n}$ be any matrix (possibly depending on A). For $w \geq 0$ define the *reachability set*

$$R_w(A) := \{b^\top A : b \in \mathbb{F}_2^{2n}, |b| \leq w\} \subseteq \mathbb{F}_2^n.$$

Every row of BA that is produced with a row of B of weight at most w must lie in $R_w(A)$. The cardinality of $R_w(A)$ is bounded independently of A :

Lemma 9.5 (Reachability bound). *For every $A \in \mathbb{F}_2^{2n \times n}$ and every $w \geq 0$,*

$$|R_w(A)| \leq \sum_{j=0}^w \binom{2n}{j}.$$

Proof. Each $b \in \mathbb{F}_2^{2n}$ with $|b| \leq w$ contributes at most one element $b^\top A$ to $R_w(A)$. The number of such b is exactly $\sum_{j=0}^w \binom{2n}{j}$. $\square$

For $w = \lfloor 0.19n \rfloor$ and large n , the binomial sum is at most $2^{2n \cdot H(0.095)} < 2^{0.94n}$, where $H(\cdot)$ is the binary entropy function. Hence

$$\frac{|R_w(A)|}{2^n} \leq 2^{-0.06n}. \quad (4)$$

This ratio is exponentially small, regardless of A .

Theorem 9.6 (Any- B uniformity forcing). *Let $A \in \mathbb{F}_2^{2n \times n}$ be any isotropic basis and let $B \in \mathbb{F}_2^{m \times 2n}$ be any matrix (possibly depending on A). Let $w = \lfloor 0.19n \rfloor$ and suppose that the conditional distribution satisfies $\text{SD}(BA \mid A, \text{Uniform}_{\mathbb{F}_2^n}) \leq \delta$ for every A . Then for every row b_i of B ,*

$$\Pr[|b_i| \leq w \mid A] \leq \frac{|R_w(A)|}{2^n} + \delta \leq 2^{-0.06n} + \delta.$$

In particular, if $m \leq \text{poly}(n)$ and $\delta = o(1/m)$, then with probability $1 - o(1)$ over the randomness of the reduction every row of B has weight $> w$.

The bound is monotone in the uniformity gap: if $\text{SD}(BA \mid A) \leq \delta$ then the same conclusion holds with δ replaced by the actual gap. The theorem does not apply to the marginal model where only $\text{SD}(BA, \text{Uniform}) \leq \delta$ is required (averaged over A): low-weight rows that vary with A can yield marginally-uniform output while evading the per-instance reachability bound.

Proof. If $|b_i| \leq w$ then $c_i = b_i^\top A \in R_w(A)$. Under a uniform distribution on $\mathbb{F}_2^n$ the probability of $R_w(A)$ is $|R_w(A)|/2^n \leq 2^{-0.06n}$. Since $\text{SD}(BA, \text{Uniform}) \leq \delta$, the actual probability is at most $2^{-0.06n} + \delta$. A union bound over m rows gives

$$\Pr[\exists i : |b_i| \leq w] \leq m(2^{-0.06n} + \delta) = o(1)$$

when $m = \text{poly}(n)$ and $\delta \leq 1/(4mn)$. □

Theorem 9.6 applies to adaptive B , secret B , and any distribution over B , *provided the conditional-uniformity hypothesis holds*: BA must be close to uniform conditioned on each fixed A . In that model, once BA is forced to be uniform (otherwise the output is distinguishable from standard LPN), every row of B must have weight $> 0.19n$. The theorem does *not* cover the marginal-adaptive corner (Item 9).

The row-weight lower bound immediately amplifies the noise.

Corollary 9.7 (Noise amplification). *Under the hypotheses of Theorem 9.6, every output label $a_i = \langle b_i, e \rangle + \langle c_i, x \rangle$ has effective noise parameter*

$$p' \geq \frac{1 - (1 - 2p)^{w+1}}{2} = \frac{1 - 2^{-(w+1)}}{2}$$

at $p = 1/4$. For $w = \lfloor 0.19n \rfloor$ this gives $p' \geq \frac{1}{2} - 2^{-0.19n-1}$.

Proof. The label noise is the XOR of $|b_i|$ independent Bernoulli(p) variables. The bias of XOR of w' i.i.d. Bernoulli(p) bits is $(1 - 2p)^{w'}$. Since $|b_i| > w$, the bias is at most $(1 - 2p)^{w+1} = 2^{-(w+1)}$, which translates to the stated p' . □

Finally, the amplified noise forces an information-theoretic sample lower bound.

Corollary 9.8 (Recovery barrier). *No algorithm can recover $x \in \mathbb{F}_2^n$ from m samples with per-sample noise $p' \geq \frac{1}{2} - 2^{-0.19n-1}$ with probability $> 1/3$ unless*

$$m = \Omega(n \cdot 2^{0.38n}).$$

In particular, any linear reduction from symplectic LPN to standard LPN that produces uniform queries requires super-polynomially many output samples.

Proof. Each sample carries at most $1 - h_2(p')$ bits of information about x , where h_2 is the binary entropy function. For $p' = \frac{1}{2} - \epsilon$ with $\epsilon = 2^{-0.19n-1}$, we have $1 - h_2(p') = \Theta(\epsilon^2) = \Theta(2^{-0.38n})$. To resolve n bits of entropy requires $m \geq n/\Theta(2^{-0.38n}) = \Omega(n \cdot 2^{0.38n})$. □

Together, Lemma 9.5, Theorem 9.6, Corollary 9.7, and Corollary 9.8 close the linear-reduction landscape *in the conditional-uniformity model*: any linear reduction that makes BA uniform *conditioned on each* A is forced into high-weight rows, which amplifies the noise to $p' \geq \frac{1}{2} - 2^{-\Theta(n)}$, and recovery then requires $m = 2^{\Theta(n)}$. At $m = \text{poly}(n)$ the reduction fails in this model, for every B model, every isotropic A , and every noise rate $p = \Theta(1)$. The marginal-adaptive corner, where B may depend on A in a way that yields uniform BA marginally but not conditionally, is not covered by this argument.

A sharper, unconditional barrier comes from an information-theoretic floor:

Proposition 9.9 (Entropy floor). *Any reduction from $\text{LSN}(n)$ to $\text{LPN}(k)$ that preserves the secret with recovery probability $\alpha(n)$ must satisfy*

$$k \geq \alpha(n) H(L) - O(1) = \alpha(n) \Theta(n^2).$$

In particular, constant success probability requires $k = \Omega(n^2)$, and success probability $1 - o(1)$ requires $k \geq H(L) - o(n^2)$.

Proof. The LSN secret space is $\text{Lagr}(2n, \mathbb{F}_2)$ with $|\text{Lagr}| = \prod_{i=1}^n (2^i + 1)$. Hence $H(L) = \log_2 |\text{Lagr}| = n(n+1)/2 + \Theta(1)$, where the additive constant is $\log_2 \prod_{i=1}^{\infty} (1 + 2^{-i}) \approx 1.25$. If a reduction outputs an $\text{LPN}(k)$ instance from which L is recovered with probability α , Fano's inequality gives $k \geq \alpha H(L) - H_2(1 - \alpha) = \alpha H(L) - O(1)$. $\square$

Concrete-security consequence. For near-perfect recovery of L (i.e., $\alpha = 1 - o(1)$ in Proposition 9.9), the entropy floor forces $k \geq H(L) - o(n^2)$. For the parameter sets in Table 1 this gives $k \geq 863$ (80-bit), $k \geq 2147$ (128-bit), $k \geq 4755$ (192-bit), or $k \geq 8387$ (256-bit). The best known attack on $\text{LPN}(k)$ (BKW [BKW03]) requires time $2^{\Omega(k/\log k)}$. At these dimensions this yields $2^{88.5}$, $2^{194.0}$, $2^{389.3}$, and $2^{643.5}$ respectively—all exceeding the target security levels. We stress that this is a concrete-security observation, not an asymptotic impossibility: BKW on $\text{LPN}(\Theta(n^2))$ runs in $2^{\Theta(n^2/\log n)}$, which is asymptotically smaller than brute-force on LSN ($2^{\Theta(n^2)}$). For the parameter range of interest, however, the entropy floor lands BKW well above the security budget.

Single-sample adaptivity does not increase degree. A natural question is whether *adaptive* degree- D feature-map reductions circumvent the static degree barrier. For the restricted model in which the reduction is *single-sample*—it may choose each ϕ_j adaptively, but applies it only to the raw samples (a, b) , with the choice of ϕ_j made as a function of previous answers—the answer is negative: in each round the feature map $\phi_j(a, b)$ has degree at most D , and the effective degree of the entire transcript is bounded by D . When $D < n$, the Reed–Muller obstruction of Section 5 still applies: correlation with $\mathbf{1}_L$ is at most 2^{-n} , requiring exponentially many samples. Thus single-sample adaptive degree- D feature-map reductions are blocked for every $D < n$, independent of the number of rounds.

The remaining open question is whether a *multi-sample* or *genuinely non-linear* adaptive reduction exists. Such a reduction could apply non-linear functions to *fresh* samples generated from previous outputs, or could use adaptively chosen non-linear queries in a way that raises the algebraic degree of the transcript. We do not claim impossibility for this richer model. We note a complementary point: a reduction in this richer model would likely improve random self-reductions for LPN itself, and so would be of independent interest even if it placed LSN within an existing assumption.

The symplectic structure is reduction-inert. A sharper observation comes from the structure of the LSN samples themselves. The deterministic quadratic constraint $S_A = 0$ (the Ω -Gram of the public matrix columns) is *public* and *x-free*: it depends only on the public matrix

A , not on the secret x or the noise. Consequently, $S_A = 0$ adds no x -information to an attacker and provides no usable handle for a reductionist trying to embed LPN into LSN. This mirrors the accepted status of Ring-LWE, which is regarded as a lattice-family assumption by *source* (ring structure native to the problem) rather than by reduction to plain LWE. We formalize this as:

Conjecture 9.10 (Source novelty). *The following three invariants of LSN have no analogue in standard LPN, and each is destroyed by every natural reduction class that has been proposed:*

1. **Symplectic-Fourier self-duality.** *The indicator $\mathbf{1}_L$ satisfies $\mathcal{F}_\Omega[\mathbf{1}_L] = 2^n \mathbf{1}_L$, where $\mathcal{F}_\Omega$ is the symplectic Fourier transform. A standard linear function $f(x) = \langle s, x \rangle$ has no such fixed-point property; its Fourier transform is a Dirac mass, not a function.*
2. **x -free quadratic constraint (sympLPN).** *In the sympLPN variant (Definition 4.3) the public matrix A satisfies $S_A = 0$ (the Ω -Gram of columns is identically zero). This is a deterministic, public, quadratic constraint on the public parameters alone. The batch-LSN distribution with pseudorandom columns (Definition 4.2) draws pseudorandom rather than isotropic columns, so this invariant does not apply there; it is a structural distinction of the sympLPN variant studied by LPQR26. LPN has no analogous public structure linking its samples.*
3. **Stabilizer degeneracy.** *Every LSN secret L is a stabilizer state with stabilizer weight $\Omega(n)$. The secret space is the Lagrangian Grassmannian, not a vector space. LPN secrets are bare linear functions.*

Each invariant is preserved by the symplectic group $\mathrm{Sp}(2n)$ and destroyed by any of the blocked reductions: linear maps lose the self-duality (Lu et al. [LPQR26]), degree- $D < n$ polynomial maps cannot represent $\mathbf{1}_L$ (Reed–Muller distance), and entropy-compressing maps discard the stabilizer structure (Proposition 9.9).

Conjecture 9.10 is falsifiable: a single explicit reduction (membership-LSN, batch-LSN, or sympLPN) $\rightarrow$ LPN that preserves any of the three invariants would refute the conjecture in its current form. It is not decidable by reduction analysis alone; it rests on the absence of such a reduction and the structural barriers documented above.

The scrambling barrier at constant noise. The LPQR26 linear barrier (Theorem D.2) applies at any noise rate $p = \omega(1/n)$, and at our constant noise $p = 1/4$ its error-amplification step is in fact *stronger*: a randomizing row of weight w leaves per-bit bias $(1 - 2p)^w = 2^{-w}$ (piling-up), versus $\approx 1 - 2wp$ in the low-noise regime. The residual gap is therefore not about noise: linear reductions consuming $m = \omega(n)$ samples are ruled out in the conditional-uniformity model for all $m = \mathrm{poly}(n)$ by the reachability barrier (Theorem 9.6). The marginal-adaptive corner, in which $B = g(A)$ yields marginally-uniform BA via low-weight rows that vary with A , is not covered by this argument and remains open (Open Problem 9). What remains outside both regimes is only non-linear scramblers. For (ii) an algebraic obstruction remains: $S_A = 0$ is a deterministic quadratic relation among the public vectors (our observation, distinct from the LPQR entropy argument). Direct inspection of the $n = 3$ case shows that simple scramblers fall into one of three classes: a linear scrambler *transports* the quadratic relation ($S'_{A'} = 0$ with respect to the conjugated form $M^{-\top} \Omega M^{-1}$, so it remains detectable), a non-linear scrambler destroys the linear label structure required by LPN, and a random scrambler destroys the secret entirely. None of these simple classes simultaneously destroys $S_A = 0$ and creates a fresh LPN secret-bit structure. The residual question — whether a sufficiently complex non-linear scrambler exists — is the same as Open Problem 2.

The marginal-adaptive corner: entropy-support bound. The gap between conditional and marginal uniformity is bridged by an entropy-support argument that does not require per-instance uniformity. The idea is simple: the isotropic basis A carries entropy $H(A) = \Theta(n^2)$, while each low-weight row of B can only “spend” $O(n)$ bits of entropy on its output query. If the total output entropy is nm (uniformity), the budget forces most rows to have high weight.

Lemma 9.11 (Entropy-support bound for low-weight rows). *Let $A \in \mathbb{F}_2^{2n \times n}$ be a random isotropic basis and let $B = g(A, R) \in \mathbb{F}_2^{m \times 2n}$ be any randomized function of A . Let $C = BA$ and suppose $\text{SD}(C, \text{Uniform}_{\mathbb{F}_2^{m \times n}}) \leq \delta$. Let $w = \lfloor 0.19n \rfloor$. Then the expected number of rows of B with weight $\leq w$ satisfies*

$$\mathbb{E}[\#\{i : |b_i| \leq w\}] \leq 16n + 11\delta m + \frac{11m}{n} + O(1),$$

Proof. By the Fannes–Csiszár continuity bound (Cover–Thomas §17.3.3), $\text{SD}(C, \text{Uniform}) \leq \delta$ implies $H(C) \geq nm - \delta nm - 1$. Conditioning on A and using the chain rule,

$$H(C) \leq H(A) + H(C | A) = H(A) + \sum_{i=1}^m H(c_i | A, c_{<i}) \leq H(A) + \sum_{i=1}^m H(c_i | A).$$

Let T_i be the indicator of the event $|b_i| \leq w$. Then

$$\begin{aligned} H(c_i | A) &\leq H(T_i, c_i | A) = H(T_i | A) + H(c_i | T_i, A) \\ &\leq 1 + \Pr[T_i = 1 | A] \cdot 0.906n + \Pr[T_i = 0 | A] \cdot n, \end{aligned}$$

where the first term is at most 1 because T_i is binary, the second uses Lemma 9.5 ($|R_w(A)| \leq 2^{0.906n}$), and the third uses the trivial bound $H(c_i | A) \leq n$. Simplifying,

$$H(c_i | A) \leq 1 + n - \Pr[T_i = 1 | A] \cdot 0.094n.$$

Summing over i and letting $k = \sum_i T_i$,

$$\sum_{i=1}^m H(c_i | A) \leq m + nm - \mathbb{E}[k | A] \cdot 0.094n.$$

The entropy of A is at most the log-support of the isotropic-frame distribution, $H(A) \leq (3/2)n^2 + n/2 + O(1)$. Combining with the Fannes–Csiszár bound $H(C) \geq nm - \delta nm - 1$ and the chain rule,

$$nm - \delta nm - 1 \leq (3/2)n^2 + n/2 + m + nm - \mathbb{E}[k | A] \cdot 0.094n + O(1),$$

which rearranges to $\mathbb{E}[k | A] \cdot 0.094n \leq (3/2)n^2 + n/2 + m + \delta nm + O(1)$. Since $(3/2)n^2 + n/2$ divided by $0.094n$ is at most $16n + O(1)$, taking expectations over A and R gives the claimed bound. $\square$

Lemma 9.11 bounds the *expected* number of low-weight rows in the marginal-adaptive model using only marginal uniformity and the entropy of A . When $m \geq Cn$ with $C > 16$, the lemma forces at least $m - 16n - O(1)$ rows to have weight $> 0.19n$, regardless of how B depends on A . The noise-amplification argument of Corollary 9.7 and the recovery argument of Corollary 9.8 then apply to those rows.

Lemma 9.12 (Distribution-mismatch death — **conditional**). ***Status: heuristic, not yet established.** The obstacle is that the output noise components $e'_i = \langle b_i, e \rangle$ are m linear images of the fixed $2n$ -bit symplectic-LPN noise vector e ; they therefore live in a subspace of dimension at most $2n$ and are heavily correlated when $m = \omega(n)$. Both standard repair strategies hit walls: (i) a recovery-reduction argument needs Corollary 9.8 to be robust to correlated noise, but the*

$\leq 2n$ leak-rows could over-determine $x \in \mathbb{F}_2^n$ (and the solver does not know B); (ii) a noise-entropy argument $H(Be) \leq H(e) = O(n)$ holds for fixed B but fails for random high-entropy B (since Be can be near-uniform). The $\leq 2n$ -dimensional noise structure is both why the corner is plausibly closed and why no clean argument closes it yet.

If the following statement could be proved, it would close the marginal-adaptive corner: Let $p' \in (0, 1/2)$ with $1 - 2p' \geq 1/\text{poly}(n)$. Consider any reduction outputting m samples (c_i, a_i) with $a_i = \langle c_i, x \rangle + e'_i$ where e'_i has bias $\leq 2^{-0.19n}$ on all but $\ell = o(m)$ rows. Then

$$\text{SD}((C, y), \text{LPN}_{p'}) \geq 1 - o(1)$$

provided $m \geq \frac{4n}{(1-2p')^2} \cdot \text{polylog}(n)$.

Theorem 9.13 (Deterministic marginal-adaptive barrier). *Let $A \sim \text{Uniform}(\text{Lagr}(2n, \mathbb{F}_2))$, let $B = g(A) \in \mathbb{F}_2^{m \times 2n}$ be any deterministic function of A , and set $C = BA$. Then for every noise rate p' ,*

$$\text{SD}((C, y), \text{LPN}_{p'}) \geq \text{SD}(C, \text{Uniform}_{\mathbb{F}_2^{m \times n}}) \geq 1 - \frac{|\text{Lagr}(2n, \mathbb{F}_2)|}{2^{mn}},$$

which is $1 - o(1)$ as soon as mn exceeds $\log_2 |\text{Lagr}(2n, \mathbb{F}_2)| = n(n+1)/2 + O(1)$ (e.g. for any $m \geq n+1$). The deterministic marginal-adaptive cell is therefore closed unconditionally.

Proof. $C = g(A)A$ is a deterministic function of A , so its support has size at most $|\text{Lagr}(2n, \mathbb{F}_2)|$, whereas the LPN public matrix is uniform on 2^{mn} values; the statistical distance between a distribution supported on K points and the uniform distribution on $N \geq K$ points is at least $1 - K/N$. Since (C, y) determines C , the data-processing inequality carries the bound to $\text{SD}((C, y), \text{LPN}_{p'})$. For $n = 2$ the bound is $1 - 15/4^m$, tight at $m = 3$ ($49/64$) and $m = 4$ ($241/256$) by exact enumeration. $\square$

Theorem 9.14 (Randomized marginal-adaptive linear-reduction barrier — **conditional**). **Depends on Lemma 9.12.** *Let $A \sim \text{Uniform}(\text{Lagr}(2n, \mathbb{F}_2))$ and let $B = g(A, R) \in \mathbb{F}_2^{m \times 2n}$ be any randomized adaptive matrix (using fresh private randomness R). Lemma 9.11 shows that if $\text{SD}(BA, \text{Uniform}) \leq \delta$ and $m \geq Cn$ ($C > 16$), at least $m - o(m)$ rows have weight $> 0.19n$, hence effective noise bias $\leq 2^{-0.19n}$. If Lemma 9.12 holds, this yields $\text{SD}((C, y), \text{LPN}_{p'}) = 1 - o(1)$ for every p' with $1 - 2p' \geq 1/\text{poly}(n)$, and no marginal-adaptive linear reduction maps symplectic LPN to a usable standard LPN instance when $m \geq Cn$.*

Theorem 9.14 is a **conditional** barrier: Lemma 9.11 is proven, but Lemma 9.12 remains open because of the correlated-noise obstacle described above. The deterministic marginal-adaptive case is already closed unconditionally by Theorem 9.13. The status map is therefore: fixed- B **ruled out**, public- B **ruled out**, conditionally-uniform adaptive **ruled out**, deterministic marginal-adaptive **ruled out**, randomized marginal-adaptive **open**. The remaining regime $m = \Theta(n)$ with $m < Cn$ is covered by LPQR26 Appendix D (their Theorem D.2) for fixed B (any rank, including low-rank).

10 Open Problems

We state precisely the problems whose resolution would close the standard-model gap.

1. **Statistical dimension under $S_A = 0$ for symplectic LPN [LPQR26].** The full SQ proof is blocked by the deterministic quadratic symplectic relation $S_A = 0$ (the Ω -Gram of columns) that characterises the symplectic LPN variant (Definition 4.3). Prove that conditioning on $S_A = 0$ does not reduce the statistical dimension below $2^{\Omega(n)}$. Section 7 resolves this exactly for statistical tests of bounded bundle size (the conditioned moments deviate from the unconstrained ones by $\Theta(4^{-n})$ at orders $j \leq 3$); the regime of growing bundle size, equivalently general- j moments, remains open.

2. **LPN(low-noise) $\rightarrow$ LSN.** In the cryptographically relevant regime $p = O(1/\sqrt{n})$, the reduction of [KLP+25] is vacuous. Does a non-vacuous reduction exist, or can its non-existence be proved?
3. **LWE $\rightarrow$ LSN.** A reduction from LWE (or worst-case lattice problems) to LSN would provide the gold-standard foundation currently missing.
4. **Quantum lower bound.** Prove Conjecture 8.1: any quantum algorithm solving LSN requires $\Omega(2^n)$ queries.
5. **Pencil-extremality conjecture.** Prove Conjecture 6.5: that isotropic pencils ($k \leq 2$) are extremal at scale $|\text{Lagr}(2n, \mathbb{F}_2)|/2^{2n}$, giving average correlation $\leq 5\rho_{\text{avg}}$ for every subset of that size. This would upgrade Theorem 6.6 from a conditional to an unconditional worst-case bound.
6. **Sample freshness for rerandomized LSN.** Exact secret rerandomization is possible: applying a public symplectic matrix $S \in \text{Sp}(2n)$ to the public part of each sample maps the secret L to $S \cdot L$ while preserving labels. For the natural *symplectic-orbit* family — splitting one sample into two via public matrices S_1, S_2 as $(x, b) \mapsto ((S_1x, b), (S_2x, b))$ — the question is now closed negatively at every n , by theorem: for every $n \geq 1$ and every $S_1, S_2 \in \text{Sp}(2n, \mathbb{F}_2)$, the statistical distance between the transformed pair and two independent fresh samples for the rerandomized secrets is exactly

$$1 - \frac{p^2 + (1-p)^2}{4^n} = 1 - \frac{5}{8 \cdot 4^n} \quad (p = \tfrac{1}{4}),$$

i.e. 123/128 at $n = 2$, 507/512 at $n = 3$, tending to 1. The proof has two steps: a relabeling bijection $(u, b_1, z, b_2) \mapsto (u, b_1, Tz, b_2)$ reduces every pair (S_1, S_2) to (I, I) , where the transformed distribution is supported on duplicated samples and the fresh distribution places exactly $(p^2 + (1-p)^2)/4^n$ mass on that diagonal (the shared noise bit is preserved by any symplectic map, so the orbit family cannot manufacture independence). Verified by exact enumeration at $n = 2$ (all 720 maps) and $n = 3$. The obstruction is in fact universal for *label-preserving* splits: for arbitrary public bijections f_1, f_2 of $\mathbb{F}_2^{2n}$ (not necessarily linear), comparing $(x, b) \mapsto ((f_1x, b), (f_2x, b))$ against two fresh same-secret samples gives exactly

$$\text{SD} = 1 - \frac{2p(1-p) + (1-2p)^2 A(f_1, f_2)}{4^n}, \quad A = \Pr_{L,x}[\mathbf{1}_L(f_1x) = \mathbf{1}_L(f_2x)] \leq 1,$$

so every label-preserving split has $\text{SD} \geq 1 - (p^2 + (1-p)^2)/4^n$, with equality iff $f_1 = f_2$: the duplicated noise bit cannot be removed by any public relabeling of the sample points (verified by exact enumeration over symplectic, affine, and random bijections at $n = 2, 3$). The same law extends verbatim to *label-flipping* splits $(x, b) \mapsto ((f_1x, b \oplus h_1(x)), (f_2x, b \oplus h_2(x)))$ for arbitrary public $h_i : \mathbb{F}_2^{2n} \rightarrow \mathbb{F}_2$, with A replaced by $A' = \Pr_{L,x}[\mathbf{1}_L(f_1x) \oplus \mathbf{1}_L(f_2x) = h_1(x) \oplus h_2(x)]$; since for $f_1x \neq f_2x$ the membership XOR is non-constant over L (and equals 0 when $f_1x = f_2x$), $A' = 1$ forces $f_1 = f_2$ and $h_1 = h_2$ — the universal lower bound covers the whole family, with equality only for the literal duplicate. Two refinements bound the family further. First, every public map that is *valid-output* (a bijection g of $\mathbb{F}_2^{2n} \times \mathbb{F}_2$ with $g_*(D_L) = D_{L'}$ for some Lagrangian L' , i.e. that sends LSN samples to LSN samples) is a pure secret rerandomization $g(x, b) = (Sx, b)$ with $S \in \text{Sp}(2n, \mathbb{F}_2)$. (A valid-output map preserves the label bit and sends every Lagrangian to a Lagrangian; the body map is therefore an automorphism of the symplectic polar space, which over $\mathbb{F}_2$ is exactly $\text{Sp}(2n, \mathbb{F}_2)$). At $n = 2$ this is proven by exhaustion without a linearity assumption: of all 16! permutations of $\mathbb{F}_2^4$, exactly the 720 elements

of $\text{Sp}(4, \mathbb{F}_2)$ map every Lagrangian to a Lagrangian.) Such maps cannot manufacture freshness. Second, even *b-dependent* label-preserving bijections $g_i(x, b) = (\varphi_{i,b}(x), b)$ obey $\text{SD} = 1 - (p^2 + (1-p)^2)/4^n + \frac{(1-2p)^2}{2 \cdot 4^n} (2 - A_0 - A_1) \geq 1 - (p^2 + (1-p)^2)/4^n$ with equality iff $g_0 = g_1$. What remains open is whether any transformation with non-product output structure (necessarily not valid-output) yields fresh samples; a positive answer would give tight multi-user security bounds for LSN-based constructions, while a fully general negative answer would explain why hybrid arguments lose a factor of the number of users.

7. **Membership-LSN $\leftrightarrow$ stabilizer-decoding LSN bridge.** Relate our membership formulation (Definition 4.1; secret Lagrangian, no public matrix) to the LSN of [KLP+25, LPQR26] (public stabilizer matrix $[A \mid B]$ with a Lagrangian A -block, noisy-codeword samples, secret logical string). A reduction in either direction would let the external hardness results (constant-rate LPN $\leq$ their LSN; Theorem 6.6 of [KLP+25] attaches this hardness already to the *single-sample* Search variant) and our SQ lower bounds support a *single* object; absent a bridge, the two hardness cases remain parallel rather than cumulative. A reduction $\text{THEIR} \leq \text{OUR}$ must embed the k -bit logical secret into our $\Theta(n^2)$ -bit Lagrangian secret without the junk register that [KLP+25] (Lemma 6.5) exploits to bypass the analogous dimension mismatch: from the single-sample hardness carrier it meets an information-budget obstruction shared with Item 9 (one noisy codeword carries $O(n)$ secret-correlated bits, while membership recovery needs $\Omega(n^2)$ labels); the multi-sample variant instead meets a frame-alignment obstruction (the secret arrives in a fresh public frame per sample, whereas ours is one fixed Lagrangian). The reverse direction is blocked by code visibility (a public $[A \mid B]$ would encode our secret L). None of these is an impossibility result; each blocks only the natural maps.
8. **Absence of a symplectic worst-to-average self-reduction.** Like LPN, and unlike LWE, LSN has no known worst-case-to-average-case reduction (Section 4.1 treats LSN as a primitive assumption); we record a structural obstruction to the natural symplectic route. The group $\text{Sp}(2n, \mathbb{F}_2)$ acts transitively on Lagrangians, so for $g \sim \text{Unif}(\text{Sp}(2n, \mathbb{F}_2))$ the map $L \mapsto gL$ *perfectly* randomizes the secret subspace; the obstruction is purely metric. Applying g carries a sample $y = Ax + e$ to $gy = (gA)x + ge$ (with gA spanning the uniform Lagrangian gL), but ge is no longer Bernoulli(p) 2n . In the Walsh domain $\widehat{g_* B_p}(u) = (1 - 2p)^{\text{wt}(g^\top u)}$, so correcting $g_* B_p$ back to Bernoulli(p') 2n by convolving with independent fresh noise P' would require $\widehat{P'}(u) = (1 - 2p')^{\text{wt}(u)} / (1 - 2p)^{\text{wt}(g^\top u)} \leq 1$ for every u , i.e. $1 - 2p' \leq (1 - 2p)^{W(g)}$ with $W(g) := \max_{u \neq 0} \text{wt}(g^\top u) / \text{wt}(u)$. For a generic $g \in \text{Sp}(2n, \mathbb{F}_2)$ one has $W(g) = \Theta(n)$ — immediate, since $g^\top$ maps a weight-1 vector to a column of $g^\top$, a generic non-zero vector of weight $\Theta(n)$, while always $W(g) \leq 2n$ — driving the corrected rate $p' \rightarrow 1/2$ and destroying the secret. The weight-preserving subgroup ($W = 1$, the symplectic monomial maps) is *not* transitive on Lagrangians: exhaustive enumeration of all 720 elements of $\text{Sp}(4, \mathbb{F}_2)$ shows that single-application reachability with $W \leq 2$ covers only 10 of the 15 Lagrangians, while the remaining 5 require $W = 3$ (corrected rate 0.4375 at $p = 1/4$). Leaving the noise uncorrected is circular: a public g lets the distinguisher apply g^{-1} and recover the worst-case instance verbatim. Whether an intermediate subgroup H with (monomial) $< H \leq \text{Sp}(2n, \mathbb{F}_2)$ that is transitive on Lagrangians with bounded W , or a non-symplectic randomization, escapes the barrier is open; we conjecture none does. The root cause is a Hamming-symplectic incompatibility: Hamming weight is not Sp-invariant. Since $\text{Sp}(2n, \mathbb{F}_2)$ is transitive on $\mathbb{F}_2^{2n} \setminus \{0\}$, *every* Sp-invariant law is constant on the non-zero vectors — the weight-blind one-parameter family $q \delta_0 + (1 - q) \mathcal{U}(\mathbb{F}_2^{2n} \setminus \{0\})$ — and the symplectic twirl of Bernoulli(p) 2n lands at $q = (3/4)^{2n}$, carrying no weight signal; no weight-graded Sp-invariant noise exists, so there is no discrete analogue of the Gaussian self-duality underlying Regev’s lattice reduction. This is consistent with the isotropic-constraint barrier that [LPQR26] report for adapting

LPN-style security proofs, and positions LSN with LPN among assumptions trusted by sustained cryptanalysis rather than by a self-reduction backbone. The obstruction is stated for the natural correction by an *independent* fresh-noise convolution (and rejection sampling onto $\text{Bernoulli}(p')$ has acceptance $2^{-\Theta(n)}$); whether a *correlated* or non-linear correction, or an intermediate transitive subgroup H with bounded W , escapes it is open (*evidence*: the $n = 2$ enumeration and the elementary $W(g) = \Theta(n)$; *conjecture*: that no usable intermediate H or correction exists).

9. **Randomized marginal-adaptive linear reductions (sharpened).** In the isotropic-to-LPN reduction model, the distinguisher receives the public matrix $C = BA$ and the noisy output $y = Cx + e$. The deterministic case is closed by Theorem 9.13, so the remaining question concerns reductions that use fresh private randomness R . Strong empirical evidence (the recovery experiments reported with the code repository) shows that single-sample recovery of x vanishes as $n \rightarrow \infty$. Exact computation of $I(x; y | C)$ for the marginal-uniform output — in closed form over the ordered-basis isotropic ensemble — at $n = 2, 3, 4$ further shows $I(x; y | C)/n$ *decreasing* at every comparable sample ratio m/n (e.g. 0.102, 0.077, 0.053 at $m/n = 2$ for $n = 2, 3, 4$), consistent with the $o(n)$ rate at the sizes we can check though not a proof of it. A rigorous information-theoretic proof — showing that the conditional mutual information $I(x; y | C)$ is $o(n)$ for typical random C — remains open. Previous Fisher-information and total-variation approaches targeted the wrong quantity: they bounded $I(x; y)$, but since the secret x is independent of the public C one has $I(x; y) \leq I(x; y | C)$, so a bound on $I(x; y)$ does not bound the operative quantity $I(x; y | C)$.

The status map for the linear-reduction landscape is: fixed- B **ruled out**, public- B **ruled out**, conditionally-uniform adaptive **ruled out**, marginal-adaptive **open**. Lemma 9.11 shows that any marginal-adaptive $B = g(A)$ with uniform BA must have $m - o(m)$ rows of weight $> 0.19n$ (hence effective noise bias $\leq 2^{-0.19n}$), but Lemma 9.12 is not yet established because the output noise $e' = Be$ is m linear images of a fixed $2n$ -bit symplectic-LPN noise vector and therefore lives in a subspace of dimension at most $2n$. The decisive question: *Does the $\leq 2n$ -dimensional noise structure of Be survive a high-entropy secret B , or can adaptive B fake m independent $\text{Bernoulli}(p')$ from $2n$ noise bits undetectably in C ?* A proof either way would close the marginal-adaptive corner. Two concrete strategies are provably non-reducing. The *uniform- B -per- A* instance (each B drawn uniformly given A) has output statistical distance $\rightarrow 1$ from matched-rate LPN at fixed n : writing the output mixture $q P_{\text{graph}} + (1 - q) P_{\text{full}}$, a Hellinger tensorization gives $\text{SD} \geq 1 - C(n) \rho(n)^m$ with $\rho(n) = 1 - 2^{-n}(1 - \sqrt{(1 - p_{\text{eff}})/2} - \sqrt{p_{\text{eff}}/2}) < 1$, and the optimal scalar distinguisher is the minimum syndrome weight $\min_w \text{wt}(y + Cw)$ (which captures over 99% of the exact distance numerically, while the membership test saturates at q). A row-*correlated* marginal-uniform family (rows equal with probability λ) moves the distance the wrong way — correlation collapses the support dimension of Be , making the output *more* distinguishable, not less. Both observations point to the same residual: any reducing B would have to keep Be 's joint law product-like while staying marginal-uniform, which neither strategy achieves. Moreover, the $W = 0$ spike is a *B -agnostic* provable distinguisher: for *every* marginal-uniform B , whenever $e \in \text{Col}(A)$ — i.e. $e = Aw$, an event of probability $q_{\text{graph}}(n)$ that depends only on the noise prior, not on B or its private randomness — one has $Be = BAw = Cw$, so $y = C(x + w)$ is a codeword and $\min_w \text{wt}(y + Cw) = 0$; since real LPN has $\Pr[W = 0] \leq 2^n(1 - p')^m$, the output leaks at rate $\geq q_{\text{graph}}(n) - \text{negl}$ regardless of B . This is a constant gap at each fixed n but vanishes asymptotically ($q_{\text{graph}}(n) \rightarrow 0$), so it rules out reductions at any fixed dimension without by itself giving an asymptotic bound. A sharper, *non-vanishing* distinguisher exists in the *shared-sample* model. If a reduction must answer $k \geq 2$ samples with a common public

matrix C (equivalently a fixed B), stack the outputs $Y = [y^{(1)} \mid \dots \mid y^{(k)}]$ and apply a parity-check H of C ($HC = 0$): then $HY = H(CX + BE) = HBE$. Since $C = BA$ gives $\text{Col}(C) = \text{Col}(BA) \subseteq \text{Col}(B)$ and $\ker H = \text{Col}(C)$, the noise image obeys the sharp identity $\text{rank}(HB) = \text{rank}(B) - \text{rank}(BA)$, so

$$\text{rank}(HY) \leq \text{rank}(HB) \leq n \quad \text{deterministically, for every } B$$

(the naive confinement bound $2n$ is loose by a factor of two, since H already annihilates the n -dimensional $\text{Col}(C) \subseteq \text{Col}(B)$). For full-rank B the identity gives $\text{rank}(HB) = n$ exactly; the bound can collapse to 0 only when $\text{rank}(B) \leq n$, which forces $\text{Col}(B) = \text{Col}(C)$ and hence $y \in \text{Col}(C)$ for every output — i.e. the already-provable $W = 0$ spike — so neither branch escapes detection. Real LPN, whose noise is full-entropy (and whose rate $\rightarrow 1/2$ under Lemma 9.11), has $\text{rank}(HY) \rightarrow \min(m - n, k)$; for $m \geq 2n$ and $k > n$ this exceeds n with probability $1 - o(1)$, a detection advantage that does *not* vanish in n . Thus the shared-sample (fixed- B) case is closed asymptotically by this *multi-sample* ($k \geq 2$) statistic — but for the single block of Lemma 9.12 ($k = 1$, one noise vector e) the stack has a single column and $\text{rank}(HY) \in \{0, 1\}$ is trivial, leaving that object to its low-dimensional-support signal (statistical distance $\rightarrow 1$ at fixed n ; asymptotic-in- n rate open). The genuine residual is the *fresh- B* model, where each sample uses an independent basis $A^{(i)}$ and matrix $B^{(i)}$, so the noises $B^{(i)}e^{(i)}$ no longer share one $\leq 2n$ -dimensional space and the stacked-rank invariant does not apply; there, exact computation (uniform- B , row-correlated, column-pair, and k -sample rank families, $n = 2$, $m \leq 80$) shows every tested distinguisher leaks only at a rate $\rightarrow 0$ in n . Whether the fresh- B marginal-adaptive reduction is asymptotically distinguishable remains the open core.

10.1 Limitations of the Present Work

We state explicitly the gaps a reader should weigh against the results above.

1. **No worst-case foundation.** LSN hardness is a primitive assumption: there is no reduction from a worst-case problem in the style of Regev’s reduction for LWE, and no non-vacuous reduction from low-noise LPN or from LWE is known (Open Problems 2–3).
2. **SQ bounds are evidence, not proof.** Exponential SQ lower bounds also hold for LPN and LWE; they bound a restricted (if natural and historically predictive) class of algorithms and do not distinguish hardness families. The unconditional bound of Theorem 6.4 moreover protects only a promise subfamily.
3. **Two results are conditional.** The $2^{2n-O(1)}$ SQ bound depends on the pencil-extremality conjecture (Conjecture 6.5), and the marginal-adaptive barrier (Theorem 9.14) depends on Lemma 9.12, which remains unproven; both dependencies are stated where they occur.
4. **Moment results close every fixed-order multi-secret composition; the $k = \Theta(n)$ level is open.** The closed forms of Section 7 are exact for every order j , Proposition 7.7 controls the variance multiplier at the maximal block, Proposition 7.8 recovers the full exact law of the quadrant count (with the proven TV rate $2^{-(n+1)}$), and Theorem 7.12 closes the joint composition of every fixed-size secret tuple ($k \leq n$; Theorems 7.9 and 7.11 are $k = 2, 3$). What remains open is the $k = \Theta(n)$ regime and the SQ statements built on these joint laws.
5. **Quantum hardness is conjectural.** We rule out the natural quantum attack families (Section 8) but prove no quantum query lower bound; Conjecture 8.1 has the same epistemic status as the corresponding conjectures for LPN and LWE.

A Full Proof of Pairwise Correlation

Here we give a self-contained derivation of Lemma 6.1. Let D_0 be the distribution on $\mathbb{F}_2^{2n} \times \mathbb{F}_2$ in which a is uniform and $b \sim \text{Bernoulli}(p)$ independently. For a fixed Lagrangian L , the LSN distribution D_L has density

$$D_L(a, b) = \frac{1}{2^{2n}} \left((1-p) \mathbf{1}_{b=\mathbf{1}_L(a)} + p \mathbf{1}_{b \neq \mathbf{1}_L(a)} \right).$$

The likelihood ratio is

$$\frac{D_L(a, b)}{D_0(a, b)} = 1 + \mathbf{1}_L(a) \cdot \beta \cdot \left(\mathbf{1}_{b=1} - \mathbf{1}_{b=0} \frac{p}{1-p} \right),$$

where $\beta = (1-2p)/p$. Therefore

$$\ell_L(a, b) = \mathbf{1}_L(a) \cdot \beta \cdot \left(\mathbf{1}_{b=1} - \mathbf{1}_{b=0} \frac{p}{1-p} \right).$$

For two secrets L, L' with $j = \dim(L \cap L')$,

$$\begin{aligned} \langle D_L, D_{L'} \rangle &= \mathbb{E}_{D_0}[\ell_L \ell_{L'}] \\ &= \mathbb{E}_a[\mathbf{1}_L(a) \mathbf{1}_{L'}(a)] \cdot \beta^2 \mathbb{E}_b \left[\left(\mathbf{1}_{b=1} - \mathbf{1}_{b=0} \frac{p}{1-p} \right)^2 \right] \\ &= 2^{j-2n} \cdot \beta^2 \left(p + (1-p) \frac{p^2}{(1-p)^2} \right) \\ &= 2^{j-2n} \cdot \frac{(1-2p)^2}{p(1-p)}. \end{aligned}$$

Setting $p = 1/4$ gives the coefficient $(1/2)^2 / (1/4 \cdot 3/4) = (1/4) / (3/16) = 4/3$ exactly.

B Q-Binomial Distance Distribution

The number of k -dimensional subspaces of $\mathbb{F}_2^n$ is the Gaussian binomial coefficient

$$\begin{bmatrix} n \\ k \end{bmatrix}_2 = \prod_{i=0}^{k-1} \frac{2^{n-i} - 1}{2^{k-i} - 1}.$$

To count pairs of Lagrangians with intersection dimension k , fix a k -dimensional isotropic subspace W . The number of Lagrangians containing W equals the number of Lagrangians in the symplectic quotient $W^\perp/W \cong \mathbb{F}_2^{2(n-k)}$, namely $\prod_{i=1}^{n-k} (2^i + 1)$. Summing over all k -dimensional subspaces $W \subseteq L$ and using the identity $\prod_{i=1}^{n-k} (2^i + 1) = 2^{(n-k)(n-k+1)/2} \prod_{i=1}^{n-k} (1 + 2^{-i})$, the exact probability is

$$\Pr[j = k] = \frac{\begin{bmatrix} n \\ k \end{bmatrix}_2 \cdot 2^{(n-k)(n-k+1)/2}}{\prod_{i=1}^n (2^i + 1)}.$$

The moment generating function $C_n = \mathbb{E}[2^j]$ converges rapidly to 2 as $n \rightarrow \infty$.

C Full Proof of the Affine-Coset Bias Bound

Let $N \subset \mathbb{F}_2^{2n}$ be a uniform random Lagrangian. For $\lambda \in (0, 1)$ define the weight enumerator $W_N(\lambda) := \sum_{v \in N} \lambda^{|v|}$. In Lemma 9.4 we need $\lambda = 1/2$.

Exact moments. Write $p := \Pr[v \in N \setminus \{0\}] = 1/(2^n + 1)$ and $q := \Pr[v, v' \in N \setminus \{0\} \mid v \neq v', \Omega(v, v') = 0] = 1/((2^{n-1} + 1)(2^n + 1))$. Let $X_v := \mathbf{1}_{v \in N}$ and $W := W_N(1/2) - 1 = \sum_{v \neq 0} X_v 2^{-|v|}$. Then

$$\mathbb{E}[W] = \frac{(3/2)^{2n} - 1}{2^n + 1}, \quad \text{Var}[W] = p(1-p)D + qS_0 - p^2T,$$

where $D = (5/4)^{2n} - 1$, $T = ((3/2)^{2n} - 1)^2 - D$, and $S_0 = \frac{1}{2}(T + C)$ with

$$C = \sum_{\substack{v \neq v' \\ v, v' \neq 0}} (-1)^{\Omega(v, v')} 2^{-|v| - |v'|} = (7/4)^{2n} - 2(3/2)^{2n} - (5/4)^{2n} + 2.$$

Block factorisation of C . Decompose $v = (a, b)$, $v' = (x, y)$ with $a, b, x, y \in \mathbb{F}_2^n$. The symplectic form factorises coordinate-wise: $\Omega(v, v') = \sum_{i=1}^n (a_i y_i + b_i x_i)$. Hence

$$\sum_{v, v' \in \mathbb{F}_2^{2n}} (-1)^{\Omega(v, v')} 2^{-|v| - |v'|} = \prod_{i=1}^n \underbrace{\sum_{a, b, x, y \in \{0, 1\}} (-1)^{ay + bx} 2^{-(a+b+x+y)}}_{= 49/16 = (7/4)^2} = (7/4)^{2n}.$$

Removing the terms with $v = 0$ or $v' = 0$ (each contributes $(3/2)^{2n}$) and adding back the double-counted $(0, 0)$ term yields C as above.

Asymptotic decay. Keeping only dominant exponentials in each symbol,

$$\begin{aligned} p(1-p)D &= (25/32)^n + O((25/64)^n), \\ qS_0 &= (81/64)^n + (49/64)^n + O((9/16)^n), \\ p^2T &= (81/64)^n + O((9/16)^n). \end{aligned}$$

The $(81/64)^n$ terms cancel exactly because $\frac{q}{2} - p^2 = -1/(2(2^{n-1} + 1)(2^n + 1)^2)$. Therefore

$$\text{Var}[W] = (25/32)^n + (49/64)^n + O((9/16)^n) + O((25/64)^n).$$

Since $\mathbb{E}[W] = (9/8)^n(1 + o(1))$, we obtain $\text{Var}[W]/\mathbb{E}[W]^2 = O((50/81)^n)$.

Chebyshev concentration. Choose $\varepsilon_n = (50/81)^{n/4}$. Then

$$\Pr[|W - \mathbb{E}[W]| \geq \varepsilon_n \mathbb{E}[W]] \leq \frac{\text{Var}[W]}{\varepsilon_n^2 \mathbb{E}[W]^2} = O((50/81)^{n/2}) = 2^{-\Omega(n)}.$$

On this high-probability event,

$$\frac{W_N(1/2) - 1}{2^n} = \frac{(9/8)^n}{2^n} (1 + o(1)) = (9/16)^n (1 + o(1)),$$

which yields the bound of Lemma 9.4.

D Reduction Barrier Summary

This appendix summarizes the SQ/feature-map barriers; for the full linear-reduction landscape (transport, entropy-continuity, reachability, and the open marginal-adaptive cell) see Table 2 and Section 9.

Linear. For any linear classifier $h(a) = \langle w, a \rangle + t$, the advantage

$$\text{Adv}(h) = \left| \Pr_{D_L}[h(a) = 1] - \Pr_{D_0}[h(a) = 1] \right|$$

is zero because $\Pr_{D_L}[h(a) = 1] = \Pr_{D_0}[h(a) = 1]$.

Polynomial degree $< n$. The exact polynomial representation of $\mathbf{1}_L$ has degree exactly n and 2^n monomials. Any polynomial of degree $d < n$ cannot vanish on L^c , so its correlation with $\mathbf{1}_L$ is at most 2^{-n} .

Adaptive linear SQ. For real-linear queries $q(a, b) = \langle w, a \rangle + cb$ we have $\mathbb{E}_{D_L}[q] = c(p + 2^{-n}(1 - 2p))$, independent of L . For $\mathbb{F}_2$ -linear queries the maximum advantage is $(1 - 2p)2^{-n}$ (Theorem 6.8). In either case the advantage is exponentially small in n , so adaptive linear SQ algorithms require exponentially many samples.

E Extension to $\mathbb{F}_q$

All results in the main text are stated over $\mathbb{F}_2$. Here we sketch the extension to arbitrary finite fields $\mathbb{F}_q$, which widens the parameter space and allows a trade-off between block length n and field size q .

E.1 Symplectic Spaces over $\mathbb{F}_q$

Let q be a prime power and $V = \mathbb{F}_q^{2n}$ equipped with a non-degenerate alternating bilinear form $\Omega(x, y) = x^T J y$ where $J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$. A subspace $L \subset V$ is *Lagrangian* if it is maximal isotropic; then $\dim L = n$ and $|\text{Lagr}(2n, \mathbb{F}_q)| = \prod_{i=1}^n (q^i + 1)$.

For fixed L and $j = \dim(L \cap L')$, the number of Lagrangians L' with intersection dimension j is

$$N_j(n, q) = \begin{bmatrix} n \\ j \end{bmatrix}_q \cdot q^{(n-j)(n-j+1)/2},$$

where $\begin{bmatrix} n \\ j \end{bmatrix}_q$ is the Gaussian binomial coefficient. Summing over j recovers $|\text{Lagr}(2n, \mathbb{F}_q)|$.

E.2 Exact Correlation over $\mathbb{F}_q$

The label remains binary ($b \in \mathbb{F}_2$) even though the ambient space is $\mathbb{F}_q^{2n}$, so the SQ machinery of Feldman *et al.* applies without modification. The likelihood ratio calculation is identical with 2 replaced by q :

Lemma E.1 ($\mathbb{F}_q$ pairwise correlation). *For $L, L' \in \text{Lagr}(2n, \mathbb{F}_q)$ with $j = \dim(L \cap L')$,*

$$\langle D_L, D_{L'} \rangle = \frac{(1 - 2p)^2}{p(1 - p)} \cdot \frac{q^j}{q^{2n}}.$$

Lemma E.2 ($\mathbb{F}_q$ average correlation). *Let $C_{n,q} = \mathbb{E}[q^j]$ computed via the q -binomial distribution. Then*

$$\rho_{\text{avg}} = \frac{(1 - 2p)^2}{p(1 - p)} \cdot C_{n,q} \cdot q^{-2n}.$$

Numerically, $C_{n,q} \rightarrow 2$ as $n \rightarrow \infty$ for every q (the limit is independent of the field size).

Table 3: LSN parameters over $\mathbb{F}_q$ for $p = 1/4$.

Security	$\mathbb{F}_2$: n	$\mathbb{F}_3$: n	$\mathbb{F}_5$: n	$\mathbb{F}_7$: n
80-bit	41	26	18	15
128-bit	65	41	28	24
192-bit	97	62	42	35
256-bit	129	82	56	46

E.3 Security Parameters over $\mathbb{F}_q$

The SQ lower bound becomes $q_{\text{queries}} \geq q^{2n-O(1)}$. One “bit” of security corresponds to $\log_2(q)$ field operations, so the effective bit-security is $2n \log_2(q)$. Table 3 gives the block lengths n needed for target security levels.

Larger q reduces the required n but increases the cost of each sample (field arithmetic in $\mathbb{F}_q$ versus $\mathbb{F}_2$). The optimal choice depends on the target platform.

E.4 Open Questions

1. Extend to q -ary labels $y \in \mathbb{F}_q$ with noise over $\mathbb{F}_q$; this requires q -ary SQ lower bounds.
2. The proof holds for $q = p^k$ with $k > 1$ because symplectic theory is field-agnostic.
3. In characteristic 2, alternating is equivalent to symmetric; in odd characteristic they differ. The intersection distribution formula remains valid in all characteristics.

Acknowledgements

During the preparation of this work, the author used large language models (Claude, Kimi, and Codex) to assist with literature review, mathematical exploration, code prototyping, and manuscript drafting. After using these tools, the author reviewed and edited the content as needed and takes full responsibility for the accuracy, integrity, and final content of the work.

Data and code availability. All supporting material — Python enumeration and verification scripts, Rust implementations (cryptanalysis screens, decoder validation, and a reference scaffold), SageMath symbolic checks, exact-fraction data and test vectors, and the L^AT_EX source — is available at <https://github.com/Gattochoo/lsn-pq> and archived at [doi:10.5281/zenodo.20646796](https://doi.org/10.5281/zenodo.20646796). In particular, the closed forms of Section 7 are reproducible from the enumeration and symbolic-verification scripts.

Conflict of interest. The author declares no competing interests.

References

- [AAC+22] C. Aguilar Melchor, N. Aragon, S. Bettaleb, L. Bidoux, O. Blazy, J.-C. Deneuville, P. Gaborit, E. Persichetti, G. Zémor, *et al.* HQC. NIST PQC Round 4 Submission, 2022.
- [ABD+21] M. Albrecht, D. Bernstein, T. Chou, C. Cid, J. Gilcher, T. Lange, V. Maram, I. von Maurich, R. Misoczki, R. Niederhagen, K. Paterson, E. Persichetti, C. Peters, P. Schwabe, N. Sendrier, J. Szefer, C. Tjhai, M. Tomlinson, and W. Wang. Classic McEliece. NIST PQC Round 4 Submission, 2021.
- [ABW10] A. Akavia, S. Goldwasser, and V. Vaikuntanathan. Simultaneous hardcore bits and cryptography against memory attacks. In *TCC*, pages 474–495. Springer, 2010.

- [Ale11] M. Alekhnovich. More on average case vs approximation complexity. *Computational Complexity*, 20(4):755–786, 2011.
- [BFJ+94] A. Blum, M. Furst, J. Jackson, M. Kearns, Y. Mansour, and S. Rudich. Weakly learning DNF and characterizing statistical query learning using Fourier analysis. In *STOC*, pages 253–262. ACM, 1994.
- [BFKL94] A. Blum, M. Furst, M. Kearns, and R. Lipton. Cryptographic primitives based on hard learning problems. In *CRYPTO*, pages 278–291. Springer, 1994.
- [BKW03] A. Blum, A. Kalai, and H. Wasserman. Noise-tolerant learning, the parity problem, and the statistical query model. *J. ACM*, 50(4):506–519, 2003.
- [BPR12] A. Banerjee, C. Peikert, and A. Rosen. Pseudorandom functions and lattices. In *EUROCRYPT*, pages 719–737. Springer, 2012.
- [BCGI18] E. Boyle, G. Couteau, N. Gilboa, and Y. Ishai. Compressing vector OLE. In *ACM CCS*, pages 896–912, 2018.
- [CRSS98] A. R. Calderbank, E. M. Rains, P. W. Shor, and N. J. A. Sloane. Quantum error correction via codes over $GF(4)$. *IEEE Trans. Inform. Theory*, 44(4):1369–1387, 1998.
- [DKS17] I. Diakonikolas, D. M. Kane, and A. Stewart. Statistical query lower bounds for robust estimation of high-dimensional Gaussians and Gaussian mixtures. In *FOCS*, pages 73–84. IEEE, 2017.
- [DGM14] N. Döttling, J. Müller-Quade, and A. Nascimento. IND-CCA secure cryptography based on a variant of the LPN problem. In *ASIACRYPT*, pages 485–508. Springer, 2014.
- [EKM17] A. Esser, R. Kübler, and A. May. LPN decoded. In *CRYPTO*, pages 486–514. Springer, 2017.
- [FGR+17] V. Feldman, E. Grigorescu, L. Reyzin, S. S. Vempala, and Y. Xiao. Statistical algorithms and a lower bound for detecting planted cliques. *Journal of the ACM*, 64(2):8:1–8:37, 2017.
- [GJL14] Q. Guo, T. Johansson, and C. Löndahl. Solving LPN using covering codes. In *ASIACRYPT*, pages 1–20. Springer, 2014.
- [GL22] A. Gollakota and D. Liang. On the hardness of PAC-learning stabilizer states with noise. *arXiv:2102.13312*, 2022.
- [Got97] D. Gottesman. Stabilizer codes and quantum error correction. PhD thesis, Caltech, 1997.
- [Gro96] L. K. Grover. A fast quantum mechanical algorithm for database search. In *STOC*, pages 212–219. ACM, 1996.
- [HKL+12] S. Heyse, E. Kiltz, V. Lyubashevsky, C. Paar, and K. Pietrzak. Lapin: An efficient authentication protocol based on ring-LPN. In *FSE*, pages 346–365. Springer, 2012.
- [Kea98] M. Kearns. Efficient noise-tolerant learning from statistical queries. *J. ACM*, 45(6):983–1006, 1998.
- [Kir11] P. Kirchner. Improved generalized birthday attack. *Cryptology ePrint Archive*, Report 2011/377, 2011.

- [KLP+25] A. Khesin, J. Lu, A. Poremba, S. Ramkumar, and V. Vaikuntanathan. Hardness of decoding random quantum stabilizer codes. *arXiv:2509.20697*, 2025.
- [LF06] E. Levieil and P.-A. Fouque. An improved LPN algorithm. In *SCN*, pages 348–359. Springer, 2006.
- [LPQR26] J. Lu, A. Poremba, Y. Quek, and S. Ramkumar. Post-quantum cryptography from quantum stabilizer decoding. *arXiv:2603.19110*, 2026.
- [Lyu05] V. Lyubashevsky. The parity problem in the presence of noise, decoding random linear codes, and the subset sum problem. In *APPROX-RANDOM*, pages 378–389. Springer, 2005.
- [Mac71] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford University Press, 1979.
- [McE78] R. J. McEliece. A public-key cryptosystem based on algebraic coding theory. *DSN Progress Report*, 44:114–116, 1978.
- [NC00] M. A. Nielsen and I. L. Chuang. *Quantum Computation and Quantum Information*. Cambridge University Press, 2000.
- [PQS26] A. Poremba, Y. Quek, and P. Shor. Learning stabilizers with noise. *arXiv:2410.18953*, 2025.
- [Pra62] E. Prange. The use of information sets in decoding cyclic codes. *IRE Trans. Inform. Theory*, 8(5):5–9, 1962.
- [Reg05] O. Regev. On lattices, learning with errors, random linear codes, and cryptography. *J. ACM*, 56(6):34, 2005.
- [Sho94] P. W. Shor. Algorithms for quantum computation: Discrete logarithms and factoring. In *FOCS*, pages 124–134. IEEE, 1994.