

Trigonometric functions

$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$
 $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$
 $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$
 $\sin(2\alpha) = 2 \sin \alpha \cos \alpha; \tan(2\alpha) = \frac{2 \tan \alpha}{1 - \tan^2 \alpha}$
 $\cos(2\alpha) = \cos^2 \alpha - \sin^2 \alpha =$
 $= 2 \cos^2 \alpha - 1 = 1 - 2 \sin^2 \alpha$

Hyperbolic functions

$\sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y$
 $\cosh(x + y) = \cosh x \cosh y + \sinh x \sinh y$
 $\tanh(x + y) = \frac{\tanh x + \tanh y}{1 + \tanh x \tanh y}$

Areas

triangle: $\sqrt{p(p-a)(p-b)(p-c)}$

Combinatorics

$D_{n,k} = \frac{n!}{(n-k)!}$

$P_n^{(m_1,m_2,\dots)} = \frac{n!}{m_1!m_2! \dots}$

$C_{n,k} = \binom{n}{k} = \frac{n!}{k!(n-k)!}$

$C'_{n,k} = \binom{n+k-1}{k}$

Miscellaneous

$A.B\overline{C} = \frac{ABC-AB}{9 \times C \quad 0 \times B}$

$\sqrt{a \pm \sqrt{b}} = \sqrt{\frac{a + \sqrt{a^2 - b}}{2}} \pm \sqrt{\frac{a - \sqrt{a^2 - b}}{2}}$

$\sum_{i=0}^n a^i = \frac{1-a^{n+1}}{1-a}$

$\sum_{x=1}^n x^3 = (\sum_{x=1}^n x)^2 = \frac{1}{4}n^2(n+1)^2$

Derivatives

$\tan' x = 1 + \tan^2 x$
 $\cot' x = -1 - \cot^2 x$
 $\operatorname{atan}' x = -\operatorname{acot}' x = \frac{1}{1+x^2}$
 $\operatorname{asin}' x = -\operatorname{acos}' x = \frac{1}{\sqrt{1-x^2}}$
 $(a^x)' = a^x \ln a$
 $\log'_a x = \frac{1}{x \ln a}$
 $\cosh' x = \sinh x$
 $\tanh' x = 1 - \tanh^2 x$
 $\operatorname{atanh}' x = \operatorname{acoth}' x = \frac{1}{1-x^2}$
 $\operatorname{asinh}' x = \frac{1}{\sqrt{x^2+1}}$
 $\operatorname{acosh}' x = \frac{1}{\sqrt{x^2-1}}$
 $(f^{-1})' = \frac{1}{f'(f^{-1})}$
 $\left(\frac{1}{x}\right)' = -\frac{\dot{x}}{x^2}$
 $\left(\frac{x}{y}\right)' = \frac{\dot{x}y - x\dot{y}}{y^2}$
 $(x^y)' = x^y(\dot{y} \ln x + y\frac{\dot{x}}{x})$

Integrals

$\int x^a = \frac{x^{a+1}}{a+1}$
 $\int a^x = \frac{a^x}{\ln a}$
 $\int \frac{1}{x} = \ln |x|$
 $\int \tan x = -\ln |\cos x|$
 $\int \cot x = \ln |\sin x|$
 $\int \frac{1}{\sin x} = \ln \left| \tan \frac{x}{2} \right|$
 $\int \frac{1}{\cos x} = \ln \left| \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right|$
 $\int \ln x = x(\ln x - 1)$
 $\int \tanh x = \ln \cosh x$
 $\int \coth x = \ln |\sinh x|$
 $\int \frac{1}{\sqrt{a^2-x^2}} = \operatorname{asin} \frac{x}{a}$
 $\int \frac{1}{a^2+x^2} = \frac{1}{a} \operatorname{atan} \frac{x}{a}$
 $\int xy = x \int y - \int (\dot{x} \int y)$
 $\int e^{yx} = e^{yx} \left(\frac{y}{x} - \frac{1}{y^2} \right)$

Differential equations

$\dot{x} + \dot{a}x = b : x = e^{-a} \left(\int be^a + c_1 \right)$
 $a\ddot{x} + b\dot{x} + cx = 0 : x = c_1e^{z_1t} + c_2e^{z_2t}$
 $\ddot{x} = -\omega^2x : x = c_1 \sin(\omega t) + c_2 \cos(\omega t)$
 $x\ddot{x} = k\dot{x}^2 : x = c_2 \sqrt[1-k]{(1-k)t + c_1}$
 $\dot{x} + ax^2 = b : x = \sqrt{\frac{b}{a}} \tanh \left(\sqrt{ab}(c_1 + t) \right)$
 $\ddot{x} + \gamma \dot{x} + \omega_0^2 x = fe^{-i\omega t} : x = \frac{fe^{-i\omega t}}{\omega_0^2 - \omega^2 - i\gamma\omega}$

Taylor

$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$
 $\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$
 $\tan x = x + \frac{x^3}{3} + \frac{2}{15}x^5 + \frac{17}{315}x^7 + O(x^9)$
 $\frac{1}{\sin x} = \frac{1}{x} + \frac{x}{6} + \frac{7}{360}x^3 + \frac{31}{15120}x^5 + O(x^7)$
 $\frac{1}{\cos x} = 1 + \frac{x^2}{2} + \frac{5}{24}x^4 + \frac{61}{720}x^6 + \frac{277}{8064}x^8 + O(x^{10})$
 $\frac{1}{\tan x} = \frac{1}{x} - \frac{x}{3} - \frac{x^3}{45} - \frac{2}{945}x^5 + O(x^7)$
 $\operatorname{asin} x = x + \frac{x^3}{6} + \frac{3}{40}x^5 + \frac{5}{112}x^7 + O(x^9)$
 $\operatorname{acos} x = \frac{\pi}{2} - \operatorname{asin} x$
 $\operatorname{atan} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$
 $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$
 $\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots$
 $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \dots$
 $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$
 $\sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots$
 $\cosh x = 1 + \frac{x^2}{2} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots$
 $\tanh x = x - \frac{x^3}{3} + \frac{2}{15}x^5 - \frac{17}{315}x^7 + O(x^9)$
 $\frac{1}{\sinh x} = \frac{1}{x} - \frac{x}{6} + \frac{7}{360}x^3 - \frac{31}{15120}x^5 + O(x^7)$
 $\frac{1}{\cosh x} = 1 - \frac{x^2}{2} + \frac{5}{24}x^4 - \frac{61}{720}x^6 + \frac{277}{8064}x^8 + O(x^{10})$
 $\frac{1}{\tanh x} = \frac{1}{x} + \frac{x}{3} - \frac{x^3}{45} + \frac{2}{945}x^5 + O(x^7)$
 $(1+x)^n = 1 + nx + \frac{n(n-1)}{2}x^2 + O(x^3)$
 $(1+x)^x = 1 + x^2 - \frac{x^3}{2} + \frac{5}{6}x^4 - \frac{3}{4}x^5 + O(x^6)$
 $x! = 1 - \gamma x + \left(\frac{\gamma^2}{2} + \frac{\pi^2}{12} \right) x^2 + O(x^3)$

Vectors

$\varepsilon_{ijk} = \begin{cases} 0 & i = j \vee j = k \vee k = i \\ 1 & i + 1 \equiv j \wedge j + 1 \equiv k \\ -1 & i \equiv j + 1 \wedge j \equiv k + 1 \end{cases}$
 $\varepsilon_{ijk}\varepsilon_{ilm} = \delta_{jl}\delta_{km} - \delta_{jm}\delta_{kl}$
 $\vec{a} \times \vec{b} = \varepsilon_{ijk}a_jb_k\hat{e}_i$
 $(\vec{a} \otimes \vec{b})_{ij} = a_ib_j$
 $(\vec{a} \times \vec{b})\vec{c} = (\vec{c} \times \vec{a})\vec{b}$
 $(\vec{a} \times \vec{b}) \times \vec{c} = -(\vec{b}\vec{c})\vec{a} + (\vec{a}\vec{c})\vec{b}$
 $(\vec{a} \times \vec{b})(\vec{c} \times \vec{d}) = (\vec{a}\vec{c})(\vec{b}\vec{d}) - (\vec{a}\vec{d})(\vec{b}\vec{c})$
 $|\vec{u} \times \vec{v}|^2 = u^2v^2 - (\vec{u}\vec{v})^2$
 $\vec{\nabla} = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right); \square = \frac{\partial^2}{\partial t^2} - \nabla^2$
 $\vec{\nabla} V = \frac{\partial V}{\partial \rho} \hat{\rho} + \frac{1}{\rho} \frac{\partial V}{\partial \phi} \hat{\phi} + \frac{\partial V}{\partial z} \hat{z}$

$$\begin{aligned}\vec{\nabla}\vec{v} &= \frac{1}{\rho}\frac{\partial(\rho v_\rho)}{\partial\rho} + \frac{1}{\rho}\frac{\partial v_\phi}{\partial\phi} + \frac{\partial v_z}{\partial z} \\ \vec{\nabla}\times\vec{v} &= \left(\frac{1}{\rho}\frac{\partial v_z}{\partial\phi} - \frac{\partial v_\phi}{\partial z}\right)\hat{\rho} + \\ &+ \left(\frac{\partial v_\rho}{\partial z} - \frac{\partial v_z}{\partial\rho}\right)\hat{\phi} + \frac{1}{\rho}\left(\frac{\partial(\rho v_\phi)}{\partial\rho} - \frac{\partial v_\rho}{\partial\phi}\right) \\ \nabla^2 V &= \frac{1}{\rho}\frac{\partial}{\partial\rho}\left(\rho\frac{\partial V}{\partial\rho}\right) + \frac{1}{\rho^2}\frac{\partial^2 V}{\partial\phi^2} + \frac{\partial^2 V}{\partial z^2} \\ \vec{\nabla}V &= \frac{\partial V}{\partial r}\hat{r} + \frac{1}{r}\frac{\partial V}{\partial\theta}\hat{\theta} + \frac{1}{r\sin\theta}\frac{\partial V}{\partial\varphi}\hat{\varphi} \\ \vec{\nabla}\vec{v} &= \frac{1}{r^2}\frac{\partial(r^2 v_r)}{\partial r} + \frac{1}{r\sin\theta}\frac{\partial(v_\theta\sin\theta)}{\partial\theta} + \frac{1}{r\sin\theta}\frac{\partial v_\varphi}{\partial\varphi} \\ \vec{\nabla}\times\vec{v} &= \frac{1}{r\sin\theta}\left(\frac{\partial(v_\varphi\sin\theta)}{\partial\theta} - \frac{\partial v_\theta}{\partial\varphi}\right)\hat{r} + \\ &+ \frac{1}{r}\left(\frac{1}{\sin\theta}\frac{\partial v_r}{\partial\varphi} - \frac{\partial(rv_\varphi)}{\partial r}\right)\hat{\theta} + \frac{1}{r}\left(\frac{\partial(rv_\theta)}{\partial r} - \frac{\partial v_r}{\partial\theta}\right)\hat{\varphi} \\ \nabla^2 V &= \frac{\frac{\partial}{\partial r}(r^2\frac{\partial V}{\partial r})}{r^2} + \frac{\frac{\partial}{\partial\theta}(\sin\theta\frac{\partial V}{\partial\theta})}{r^2\sin\theta} + \frac{\frac{\partial^2 V}{\partial\varphi^2}}{r^2\sin^2\theta} \\ \vec{\nabla}\left(\vec{\nabla}\times\vec{v}\right) &= \vec{\nabla}\times\vec{\nabla}\vec{V} = 0 \\ \vec{\nabla}(f\vec{v}) &= (\vec{\nabla}f)\vec{v} + f\vec{\nabla}\vec{v} \\ \vec{\nabla}\times(f\vec{v}) &= \vec{\nabla}f\times\vec{v} + f\vec{\nabla}\times\vec{v} \\ \vec{\nabla}\times(\vec{\nabla}\times\vec{v}) &= -\nabla^2\vec{v} + \vec{\nabla}(\vec{\nabla}\cdot\vec{v}) \\ \vec{\nabla}(\vec{v}\times\vec{w}) &= \vec{w}(\vec{\nabla}\times\vec{v}) - \vec{v}(\vec{\nabla}\times\vec{w}) \\ \vec{\nabla}\times(\vec{v}\times\vec{w}) &= (\vec{\nabla}\cdot\vec{w} + \vec{w}\cdot\vec{\nabla})\vec{v} - (\vec{\nabla}\cdot\vec{v} + \vec{v}\cdot\vec{\nabla})\vec{w}\end{aligned}$$

$$\begin{aligned}\frac{1}{2}\vec{\nabla}v^2 &= (\vec{v}\cdot\vec{\nabla})\vec{v} + \vec{v}\times(\vec{\nabla}\times\vec{v}) \\ \int\vec{\nabla}\vec{v}\mathrm{d}^3x &= \oint\vec{v}\mathrm{d}\vec{S}; \int(\vec{\nabla}\times\vec{v})\mathrm{d}\vec{S} = \oint\vec{v}\mathrm{d}\vec{l} \\ \int(f\nabla^2g - g\nabla^2f)\mathrm{d}^3x &= \oint_S(f\frac{\partial g}{\partial n} - g\frac{\partial f}{\partial n})\mathrm{d}S \\ \oint\vec{v}\times\mathrm{d}\vec{S} &= -\int(\vec{\nabla}\times\vec{v})\mathrm{d}^3x \\ \delta(\vec{r}-\vec{r}_0) &= \frac{\delta(r-r_0)\delta(\theta-\theta_0)\delta(\varphi-\varphi_0)}{r^2\sin\theta_0} \\ \nabla^2\frac{1}{|\vec{r}-\vec{r}_0|} &= -4\pi\delta(\vec{r}-\vec{r}_0) \\ \delta(g(x)) &= \frac{\delta(x-x_i)}{|g'(x_i)|}; g(x_i) = 0 \\ \langle\mathrm{Re}(ae^{-i\omega t})\mathrm{Re}(be^{-i\omega t})\rangle &= \frac{1}{2}\mathrm{Re}(a\bar{b})\end{aligned}$$

Statistics

$$\begin{aligned}P(E\cap E_1) &= P(E_1)\cdot P(E|E_1) \\ \Delta x_{\mathrm{hist}} &\approx \frac{x_{\mathrm{max}}-x_{\mathrm{min}}}{\sqrt{N}} \\ P(x\leq k) &= F(k) = \int_{-\infty}^k p(x) \\ \mathrm{median} &= F^{-1}(\tfrac{1}{2}) \\ E[f(x)] &= \int_{-\infty}^{\infty} f(x)p(x) \\ \mu &= E[x] = \int_{-\infty}^{\infty} xp(x) \\ \alpha_n &= E[x^n] \\ M_n &= E[(x-\mu)^n] \\ \sigma^2 &= M_2 = E[x^2] - \mu^2 \\ \mathrm{FWHM} &\approx 2\sigma \\ \gamma_1 &= \frac{M_3}{\sigma^3}, \gamma_2 = \frac{M_4}{\sigma^4}\end{aligned}$$

$$\begin{aligned}\phi[y](t) &= E[e^{ity}] \\ \phi[y_1+\lambda y_2] &= \phi[y_1]\phi[\lambda y_2] \\ \alpha_n &= i^{-n}\frac{\partial^n t}{\partial\phi[x]^n}\Big|_{t=0} \\ h\geq 0: P(h\geq k) &\leq \frac{E[h]}{k} \\ P(|x-\mu|>k\sigma) &\leq \frac{1}{k^2} \\ B(n,p,k) &= \binom{n}{k}p^k(1-p)^{n-k} \\ \mu_B &= np, \sigma_B^2 = np(1-p) \\ P(\mu,k) &= \frac{\mu^k}{k!}e^{-\mu}, \sigma_P^2 = \mu \\ u(x,a,b) &= \frac{1}{b-a}, x\in[a;b] \\ \mu_u &= \frac{b+a}{2}, \sigma_u^2 = \frac{(b-a)^2}{12} \\ \varepsilon(x,\lambda) &= \lambda e^{-\lambda x}, x\geq 0\end{aligned}$$

$$\begin{aligned}\mu_\varepsilon &= \frac{1}{\lambda}, \sigma_\varepsilon^2 = \frac{1}{\lambda^2} \\ g(x,\mu,\sigma) &= \frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2} \\ \mathrm{FWHM}_g &= 2\sigma\sqrt{2\ln 2} \\ z &= \frac{x-\mu}{\sigma}; \mu, \sigma[z] = 0, 1 \\ \chi^2 &= \sum_{i=1}^n z_i^2 \\ \wp_n(x) &= \frac{2^{-\frac{n}{2}}}{\Gamma(\frac{n}{2})}x^{\frac{n}{2}-1}e^{-\frac{x}{2}} \\ \mu_\varphi &= n, \sigma_\varphi^2 = 2n \\ n\geq 30: \wp_n(x) &\approx g(x,n,\sqrt{2n}) \\ n\geq 8: p[\sqrt{2\chi^2}] &\approx g(\sqrt{2n-1}, 1) \\ S(x,n) &= \frac{\Gamma(\frac{n+1}{2})}{\sqrt{n\pi}\Gamma(\frac{n}{2})}\left(1+\frac{x}{n}\right)^{-\frac{n+1}{2}} \\ \mu_S &= 0, \sigma_S^2 = \frac{n}{n-2}\end{aligned}$$

$$\begin{aligned}p[z\sqrt{\frac{n}{\chi^2}}] &= S(n) \\ n\geq 35: S(x,n) &\approx g(x,0,1) \\ c(x,a) &= \frac{a}{\pi}\frac{1}{a^2+x^2} \\ \sigma_{xy} &= E[xy] - \mu_x\mu_y \leq \sigma_x\sigma_y \\ \rho &= \frac{\sigma_{xy}}{\sigma_x\sigma_y}, |\rho| \leq 1 \\ \mu[f(x_1,\dots)] &\approx f(\mu_1,\dots) \\ \sigma^2[f(x_1,\dots)] &\approx \sigma_{x_ix_j}\frac{\partial f}{\partial x_i}\Big|_{\mu_i}\frac{\partial f}{\partial x_j}\Big|_{\mu_j} \\ \mu &\approx m = \frac{1}{n}\sum_{i=1}^n x_i \\ \sigma^2 &\approx s^2 = \frac{1}{n-1}\sum_{i=1}^n (x_i-m)^2 \\ s_m^2 &= \frac{s^2}{n} \\ p[\frac{m-\mu}{s_m}] &= S(n)\end{aligned}$$

Fit

$$\begin{aligned}f(x) &= mx+q, \quad f(x) = a \\ f(x) &= bx \\ m &= \frac{\frac{\sum\frac{1}{\Delta y^2}\cdot\sum\frac{xy}{\Delta y^2}-\sum\frac{x}{\Delta y^2}\cdot\sum\frac{y}{\Delta y^2}}{\sum\frac{1}{\Delta y^2}\cdot\sum\frac{x^2}{\Delta y^2}-(\sum\frac{x}{\Delta y^2})^2}} \\ \Delta m^2 &= \frac{\sum\frac{1}{\Delta y^2}}{\sum\frac{1}{\Delta y^2}\cdot\sum\frac{x^2}{\Delta y^2}-(\sum\frac{x}{\Delta y^2})^2} \\ q &= \frac{\sum\frac{y}{\Delta y^2}\cdot\sum\frac{x^2}{\Delta y^2}-\sum\frac{x}{\Delta y^2}\cdot\sum\frac{xy}{\Delta y^2}}{\sum\frac{1}{\Delta y^2}\cdot\sum\frac{x^2}{\Delta y^2}-(\sum\frac{x}{\Delta y^2})^2} \\ \Delta q^2 &= \frac{\sum\frac{x^2}{\Delta y^2}}{\sum\frac{1}{\Delta y^2}\cdot\sum\frac{x^2}{\Delta y^2}-(\sum\frac{x}{\Delta y^2})^2} \\ a &= \frac{\sum\frac{y}{\Delta y^2}}{\sum\frac{1}{\Delta y^2}}, \Delta a^2 = \frac{1}{\sum\frac{1}{\Delta y^2}} \\ b &= \frac{\sum\frac{xy}{\Delta y^2}}{\sum\frac{x^2}{\Delta y^2}}, \Delta b^2 = \frac{1}{\sum\frac{x^2}{\Delta y^2}}\end{aligned}$$

Kinematics

$$\begin{aligned}\frac{1}{R} &= \left|\frac{v_xa_y-v_ya_x}{v^3}\right| \\ \vec{\omega} &= \dot{\varphi}\cos\theta\hat{r} - \dot{\varphi}\sin\theta\hat{\theta} + \dot{\theta}\hat{\varphi} \\ \dot{\vec{w}} &= \frac{\mathrm{d}(\vec{w}\hat{r})}{\mathrm{d}t}\hat{r} + \frac{\mathrm{d}(\vec{w}\hat{\theta})}{\mathrm{d}t}\hat{\theta} + \frac{\mathrm{d}(\vec{w}\hat{\varphi})}{\mathrm{d}t}\hat{\varphi} + \vec{\omega}\times\vec{w} \\ \theta\equiv\frac{\pi}{2}\rightarrow\dot{\vec{r}} &= \dot{r}\hat{r} + r\dot{\varphi}\hat{\varphi} \\ \theta\equiv\frac{\pi}{2}\rightarrow\ddot{\vec{r}} &= (\ddot{r}-r\dot{\varphi}^2)\hat{r} + (r\ddot{\varphi}+2\dot{r}\dot{\varphi})\hat{\varphi} \\ \vec{r} &= \dot{r}\hat{r} + r\dot{\theta}\hat{\theta} + r\dot{\varphi}\sin\theta\hat{\varphi} \\ \langle\vec{r},\hat{r}\rangle &= \dot{r} - r\dot{\theta}^2 - r\dot{\varphi}^2\sin^2\theta \\ \langle\vec{r},\hat{\theta}\rangle &= r\ddot{\theta} + 2\dot{r}\dot{\theta} - r\dot{\varphi}^2\sin\theta\cos\theta \\ \langle\vec{r},\hat{\varphi}\rangle &= r\ddot{\varphi}\sin\theta + 2\dot{r}\dot{\varphi}\sin\theta + 2r\dot{\theta}\dot{\varphi}\cos\theta\end{aligned}$$

$$\begin{aligned}\vec{A} &= \ddot{\vec{r}} + \vec{A}_T + \vec{\omega}\times(\vec{\omega}\times\vec{r}) + \dot{\vec{\omega}}\times\vec{r} + 2\vec{\omega}\times\dot{\vec{r}} \\ \hat{z} &= \hat{r}\cos\theta - \hat{\theta}\sin\theta \\ \hat{y} &= \hat{\theta}\cos\theta + \hat{\varphi}\sin\theta \\ \hat{x} &= \hat{\varphi}\cos\theta - \hat{r}\sin\theta\end{aligned}$$

Mechanics

$$\begin{aligned}\dot{\alpha} &= \frac{\mathrm{d}}{\mathrm{d}t}\alpha(\beta,t) = \frac{\partial\alpha}{\partial\beta}\dot{\beta} + \frac{\partial\alpha}{\partial t} \\ \vec{p} &:= m\dot{\vec{r}}; \vec{F} = \dot{\vec{p}}; \frac{\mathrm{d}(mT)}{\mathrm{d}t} = \vec{F}\cdot\vec{p} \\ M &:= \sum_i m_i; \vec{R} := \frac{m_i\vec{r}_i}{M} \\ T &= \frac{1}{2}M\dot{\vec{R}}^2 + \frac{1}{2}m_i(\dot{\vec{r}}_i - \dot{\vec{R}})^2 \\ \vec{L} &= \vec{R}\times M\dot{\vec{R}} + (\vec{r}_i - \vec{R})\times m_i(\dot{\vec{r}}_i - \dot{\vec{R}}) \\ \vec{\tau}_O &= \dot{\vec{L}}_O + \vec{v}_O\times\vec{p} \\ \tau_1 &= I_1\omega_1 + (I_3 - I_2)\omega_3\omega_2 \\ \mathcal{L}(q,\dot{q},t) &= T - V + \frac{\mathrm{d}}{\mathrm{d}t}f(q,t) \\ S[q] &= \int_{t_1}^{t_2}\mathcal{L}(q,\dot{q},t)\mathrm{d}t \\ \frac{\partial}{\partial\epsilon}S[q+\epsilon] \Big|_{\epsilon=0}^{\epsilon(t_1)=\epsilon(t_2)=0} &= 0 \\ p &:= \frac{\partial\mathcal{L}}{\partial\dot{q}}; \dot{p} = \frac{\partial\mathcal{L}}{\partial q} \\ \mathcal{H}(q,p,t) &= \dot{q}p - \mathcal{L} \\ \dot{q} &= \frac{\partial\mathcal{H}}{\partial p}; \dot{p} = -\frac{\partial\mathcal{H}}{\partial q} \\ \frac{\mathrm{d}\mathcal{H}}{\mathrm{d}t} &= \frac{\partial\mathcal{H}}{\partial t} = -\frac{\partial\mathcal{L}}{\partial t} \\ \{u,v\} &= \frac{\partial u}{\partial q}\frac{\partial v}{\partial p} - \frac{\partial u}{\partial p}\frac{\partial v}{\partial q} \\ \frac{\mathrm{d}u}{\mathrm{d}t} &= \{u,\mathcal{H}\} + \frac{\partial u}{\partial t} \\ \eta &= (q,p); \Gamma = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ \dot{\eta} &= \Gamma\frac{\partial\mathcal{H}}{\partial\eta}; \{u,v\} = \frac{\partial u}{\partial\eta}\Gamma\frac{\partial v}{\partial\eta}\end{aligned}$$

Inertia

$$\begin{aligned}\text{point: } &mr^2 \\ \text{two points: } &\mu d^2 \\ \text{rod: } &\frac{1}{12}mL^2 \\ \text{disk: } &\frac{1}{2}mr^2 \\ \text{tetrahedron: } &\frac{1}{20}ms^2 \\ \text{octahedron: } &\frac{1}{10}ms^2 \\ \text{sphere: } &\frac{2}{3}mr^2 \\ \text{ball: } &\frac{2}{5}mr^2 \\ \text{cone: } &\frac{3}{10}mr^2 \\ \text{torus: } &m(R^2 + \frac{3}{4}r^2) \\ \text{ellipsoid: } &I_a = \frac{1}{5}m(b^2+c^2) \\ \text{rectangulus: } &\frac{1}{12}m(a^2+b^2)\end{aligned}$$

Kepler

$$\begin{aligned}\langle U \rangle &\approx -2\langle T \rangle \\ U_{\mathrm{eff}} &= U + \frac{L^2}{2mr^2} \\ \frac{1}{\mu} &= \frac{1}{m_1} + \frac{1}{m_2} \\ \vec{r} &= \vec{r}_1 - \vec{r}_2, \alpha = Gm_1m_2 \\ T &= \frac{1}{2}M\dot{\vec{R}}^2 + \frac{1}{2}\mu\dot{\vec{r}}^2 \\ \vec{L} &= \vec{R}\times M\dot{\vec{R}} + \vec{r}\times\mu\dot{\vec{r}} \\ k &= \frac{L^2}{\mu\alpha}, \varepsilon = \sqrt{1 + \frac{2EL^2}{\mu\alpha^2}} \\ r &= \frac{k}{1+\varepsilon\cos\theta} \\ a &= \frac{k}{|1-\varepsilon^2|} = \frac{\alpha}{2|E|} \\ a^3\omega^2 &= G(m_1+m_2) = \frac{\alpha}{\mu} \\ \vec{A} &= \mu\dot{\vec{r}}\times\vec{L} - \mu\alpha\hat{r}, \dot{\vec{A}} = 0\end{aligned}$$

Inequalities

$$\begin{aligned}|a|-|b| &\leq |a+b| \leq |a|+|b| \\ x>-1: 1+nx &\leq (1+x)^n \\ \frac{|a^n-b^n|}{|a-b|<1} &\leq n(1+|b|)^{n-1} \\ \sqrt[p]{\sum(a_i+b_i)^p} &\leq \sqrt[p]{\sum a_i^p} + \sqrt[p]{\sum b_i^p} \\ \sum a_ib_i &\leq (\sum a_i^p)^{\frac{1}{p}}(\sum b_i^{\frac{p}{p-1}})^{\frac{p-1}{p}} \\ x^py^q &\leq \left(\frac{px+qy}{p+q}\right)^{p+q} \\ \sqrt[p]{\frac{1}{n}\sum a_i^{p\leq q}} &\leq \sqrt[q]{\frac{1}{n}\sum a_i^q} \\ \sum\left(\frac{a_1+\dots+a_i}{i}\right)^p &\leq \left(\frac{p}{p-1}\right)^p\sum a_i^p \\ x\geq 0, |\ddot{x}| &\leq M: |\dot{x}| \leq \sqrt{2Mx} \\ \frac{1}{1+x} &< \ln\left(1+\frac{1}{x}\right) < \frac{1}{x}\end{aligned}$$

Vector spaces
 $(V, \mathbb{K}, +, \cdot)$ vector space; \mathbb{K} field
 $\exists \vec{0} \in V : \vec{v} + \vec{0} = \vec{v}$
 $\cdot : \mathbb{K} \times V \rightarrow V; \quad \lambda \cdot (\vec{v} + \vec{w}) = \lambda \vec{v} + \lambda \vec{w}$
 $0_{\mathbb{K}} \cdot \vec{v} = \vec{0}, 1_{\mathbb{K}} \cdot \vec{v} = \vec{v}$

$\lambda \in \mathbb{K}, \vec{v}, \vec{w} \in V \Rightarrow \vec{v} + \vec{w} \in V, \lambda \vec{v} \in V$
 $\dim(U + V) = \dim U + \dim V - \dim(U \cap V)$
 ℓ linear : $\ell(\vec{v} + \lambda \vec{w}) = \ell(\vec{v}) + \lambda \ell(\vec{w})$
 $\ker \ell = \{ \vec{v} \in V \mid \ell(\vec{v}) = 0 \}$
 $\dim V = \dim \ell(V) + \dim(V \cap \ker \ell)$

$\langle, \rangle : V \times V \rightarrow \mathbb{K}; \quad \langle \vec{v}, \vec{w} \rangle = \langle \vec{w}, \vec{v} \rangle$
 $\langle \vec{v} + \lambda \vec{w}, \vec{u} \rangle = \langle \vec{v}, \vec{u} \rangle + \lambda \langle \vec{w}, \vec{u} \rangle$
 $\| \| : V \rightarrow \mathbb{K}; \quad \| \vec{v} \| = 0 \rightarrow \vec{v} = \vec{0}$
 $\| \lambda \vec{v} \| = |\lambda| \| \vec{v} \|; \quad \| \vec{v} + \vec{w} \| \leq \| \vec{v} \| + \| \vec{w} \|^$

Symbols
 $A \quad B \quad \Gamma \quad \Delta \quad E \quad Z \quad H \quad \Theta \quad I \quad K \quad \Lambda \quad M$
 $\alpha \quad \beta \quad \gamma \quad \delta \quad \epsilon/\varepsilon \quad \zeta \quad \eta \quad \theta/\vartheta \quad \iota \quad \kappa \quad \lambda \quad \mu$

$N \quad \Xi \quad O \quad \Pi \quad P \quad \Sigma \quad T \quad \Upsilon \quad \Phi \quad X \quad \Psi \quad \Omega$
 $\nu \quad \xi \quad o \quad \pi/\varpi \quad \rho/\varrho \quad \sigma/\varsigma \quad \tau \quad \upsilon \quad \phi/\varphi \quad \chi \quad \psi \quad \omega$

Constants, units
 $\pi = 3.142$
 $e = 2.718$
 $\gamma = 5.772 \cdot 10^{-1}$
 $G = 6.674 \cdot 10^{-11} \frac{\text{m}^3}{\text{kg s}^2}$

$R = 8.314 \frac{\text{J}}{\text{mol K}}$
 $R = 8.206 \cdot 10^{-2} \frac{1 \text{atm}}{\text{mol K}}$
 $N_{\text{A}} = 6.022 \cdot 10^{23} \frac{1}{\text{mol}}$
 $k_{\text{B}} = 1.381 \cdot 10^{-23} \frac{\text{J}}{\text{K}}$

$c = 2.998 \cdot 10^8 \frac{\text{m}}{\text{s}}$
 $q_{\text{e}} = 1.602 \cdot 10^{-19} \text{ A s}$
 $m_{\text{e}} = 9.109 \cdot 10^{-31} \text{ kg}$
 $m_{\text{p}} = 1.673 \cdot 10^{-27} \text{ kg}$

$m_{\text{n}} = 1.675 \cdot 10^{-27} \text{ kg}$
 $\text{amu} = 1.661 \cdot 10^{-27} \text{ kg}$
 $h = 6.626 \cdot 10^{-34} \text{ J s}$
 $\varepsilon_0 = 8.854 \cdot 10^{-12} \frac{\text{C}^2 \text{s}^2}{\text{kg m}^3}$

$\mu_0 = 1.257 \cdot 10^{-6} \frac{\text{N}}{\text{A}^2}$
 $\mu_{\text{B}} = 9.274 \cdot 10^{-24} \text{ A m}^2$
 $\alpha = 7.297 \cdot 10^{-3}$
 $\text{eV} = 1 \cdot 10^{-12} \text{ erg}$

Chemistry
 $H = U + pV$
 $\text{d}p = 0 \rightarrow \Delta H = \text{heat transfer}$
 $G = H - TS$
 $a_i \text{A}_i \rightarrow b_j \text{B}_j$
 $\Delta H_{\text{r}}^{\circ} = b_j \Delta H_{\text{f}}^{\circ}(\text{B}_j) - a_i \Delta H_{\text{f}}^{\circ}(\text{A}_i)$
 $\forall i, j : v_{\text{r}} = - \frac{1}{a_i} \frac{\Delta[\text{A}_i]}{\Delta t} = \frac{1}{b_j} \frac{\Delta[\text{B}_j]}{\Delta t}$

$\exists k, (m_i) : v_{\text{r}} = k[\text{A}_i]^{m_i}$
 $k = Ae^{-\frac{E_{\text{a}}}{RT}}$ (Arrhenius)
 $a_{(\ell)} = \gamma \frac{[\text{X}]}{[\text{X}]_0}, [\text{X}]_0 = 1 \frac{\text{mol}}{\text{l}}$
 $a_{(g)} = \gamma \frac{p}{p_0}, p_0 = 1 \text{ atm}$
 $K = \frac{\prod a_{\text{B}_j}^{b_j}}{\prod a_{\text{A}_i}^{a_i}}, K_{\text{c}} = \frac{\prod [\text{B}_j]^{b_j}}{\prod [\text{A}_i]^{a_i}}$
 $K_p = \frac{\prod p_{\text{B}_j}^{b_j}}{\prod p_{\text{A}_i}^{a_i}}, K_{\text{n}} = \frac{\prod n_{\text{B}_j}^{b_j}}{\prod n_{\text{A}_i}^{a_i}}$

$K_{\chi} = \frac{\prod \chi_{\text{B}_j}^{b_j}}{\prod \chi_{\text{A}_i}^{a_i}}, \chi = \frac{n}{n_{\text{tot}}}$
 $K_{\text{c}} = K_p(RT)^{\sum a_i - \sum b_j}$
 $K_{\text{c}} = K_{\text{n}} V^{\sum a_i - \sum b_j}$
 $K_{\chi} = K_{\text{n}} n_{\text{tot}}^{\sum a_i - \sum b_j}$
 $\Delta G_{\text{r}}^{\circ} = -RT \ln K$
 $Q = K(t) = \frac{\prod a_{\text{B}_j}^{b_j}(t)}{\prod a_{\text{A}_i}^{a_i}(t)}$

$\Delta G = RT \ln \frac{Q}{K}$
 $\ln \frac{K_2}{K_1} = - \frac{\Delta H^{\circ}}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$
 $K_{\text{w}} = [\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$
 $\Delta E = \Delta E^{\circ} - \frac{RT}{n_{\text{e}} N_{\text{A}} q_{\text{e}}} \ln Q$ (Nerst)
 $(\text{std}) \Delta E = \Delta E^{\circ} - \frac{0.059}{n_{\text{e}}} \log_{10} Q$
 $\text{pH} = -\log_{10} [\text{H}_3\text{O}^+]$
 $K_{\text{a}} = \frac{[\text{A}^-][\text{H}_3\text{O}^+]}{[\text{AH}]}$

Thermodynamics
 $\text{d}L = p \text{d}V$

$\text{d}Q = \text{d}U + \text{d}L$
 $\text{d}S = \frac{\text{d}Q}{T}$

$C_V = \left(\frac{\text{d}Q}{\text{d}T} \right)_V$

$C_p = \left(\frac{\text{d}Q}{\text{d}T} \right)_p$

$\gamma = \frac{C_p}{C_v}$

Ideal gas
 $pV = nRT$

$c_V, c_p = \frac{C_V, C_p}{n}, \quad c_V = \frac{\text{dof}}{2} R, \quad c_p = c_V + R$
 $c_V = \frac{R}{\gamma - 1}, \quad c_p = \frac{\gamma}{\gamma - 1} R$

$\text{d}Q = 0 : pV^{\gamma}, TV^{\gamma - 1}, p^{\frac{1}{\gamma} - 1} T \text{ const.}$

Statistical mechanics
 $Z = \frac{1}{h^N} \int \text{d}q_1 \cdots \text{d}q_N \int \text{d}p_1 \cdots \text{d}p_N e^{-\beta \mathcal{H}}$

$U = - \frac{\partial}{\partial \beta} \log Z; \quad \beta = \frac{1}{k_{\text{B}} T}; \quad C = \frac{\partial U}{\partial T}$

$F(T, V) = U - TS = - \frac{\log Z}{\beta}$
 $S = - \frac{\partial F}{\partial T}$

Electronics
(MKS)
 $\left(\frac{V}{I} \right) = \left(\frac{V_0}{I_0} \right) e^{i \omega t}$

$Z = \frac{V}{I}$
 $Z_{\text{R}} = R$

$Z_{\text{C}} = -i \frac{1}{\omega C}$
 $Z_{\text{L}} = i \omega L$

$Z_{\text{series}} = \sum_k Z_k$
 $\frac{1}{Z_{\text{parallel}}} = \sum_k \frac{1}{Z_k}$

$\sum_{\text{loop}} V_k = 0$
 $\sum_{\text{node}} I_k = 0$

$\mathcal{E} = -LI'$
 $L = \frac{\Phi_B}{I}$

Relativity
 $\beta = \frac{v}{c} = \tanh \chi$
 $\gamma = \frac{1}{\sqrt{1 - \beta^2}} = \cosh \chi$
 $\vec{p} = \gamma m \vec{v}$
 $\mathcal{E} = \gamma mc^2$
free particle: $\mathcal{L} = \mathcal{E}$

$\frac{\text{d} \vec{p}}{\text{d} t} = \vec{F}$
 $\left(\begin{smallmatrix} ct' \\ x' \end{smallmatrix} \right) = \gamma \left(\begin{smallmatrix} 1 & -\beta \\ -\beta & 1 \end{smallmatrix} \right) \left(\begin{smallmatrix} ct \\ x \end{smallmatrix} \right)$
 $\chi'' = \chi' + \chi$
 $V'_{\parallel} = \frac{V_{\parallel} - v}{1 - \frac{v V_{\parallel}}{c^2}}$

$V'_{\perp} = \frac{1}{\gamma} \frac{V_{\perp}}{1 - \frac{v V_{\parallel}}{c^2}}$
 $\frac{V'}{c} = 1 - \frac{(1 - \frac{V^2}{c^2})(1 - \frac{v^2}{c^2})}{\left(1 - \frac{v V_{\parallel}}{c^2} \right)^2}$
 $\text{d} \tau = \frac{1}{\gamma} \text{d} t$
 $x^{\mu} = (ct, \vec{x})$

$v^{\mu} = \frac{\text{d} x^{\mu}}{\text{d} \tau} = \gamma(c, \vec{v})$
 $a^{\mu} = \frac{\text{d} v^{\mu}}{\text{d} \tau} = \gamma \left(\frac{\text{d} \gamma}{\text{d} t} c, \frac{\text{d}(\gamma \vec{v})}{\text{d} t} \right)$
 $p^{\mu} = m v^{\mu} = \left(\frac{\mathcal{E}}{c}, \vec{p} \right)$
 $\partial_{\mu} = \frac{\partial}{\partial x^{\mu}} = \left(\frac{1}{c} \frac{\partial}{\partial t}, \vec{\nabla} \right)$
 $g_{\mu \nu} = \left(\begin{smallmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{smallmatrix} \right)$

$x_{\mu} = g_{\mu \nu} x^{\nu}$
 $\partial_{\mu} \partial^{\mu} = \square$
 $p^{\mu} p_{\mu} = (mc)^2$
 $v^{\mu} a_{\mu} = 0$
 $M \rightarrow \sum_i m_i$
 $E_1^{\text{max}} = \frac{M^2 + m_1^2 - \sum_{i \neq 1} m_i^2}{2M} c^2$

Electrostatics (CGS)
 $\vec{F}_{12} = q_1 q_2 \frac{\vec{r}_2 - \vec{r}_1}{|\vec{r}_1 - \vec{r}_2|^3}; \quad \vec{E}_1 = \frac{\vec{E}_{12}}{q_2}; \quad V(\vec{r}) = \int \text{d}^3 r' \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|}; \quad \rho_q = \delta(\vec{r} - \vec{r}_q)$
 $\oint \vec{E} \text{d} \vec{S} = 4\pi \int \rho \text{d}^3 x; \quad -\nabla^2 V = \vec{\nabla} \cdot \vec{E} = 4\pi \rho; \quad \vec{\nabla} \times \vec{E} = 0$
 $U = \frac{1}{8\pi} \int E^2 \text{d}^3 x; \quad \tilde{U} = \frac{1}{2} \frac{q_i q_j}{|\vec{r}_i - \vec{r}_j|} = \frac{1}{8\pi} \sum_{i \neq j} \int \vec{E}_i \cdot \vec{E}_j \text{d}^3 x$
 $V(\vec{r}) = \int \rho G_{\text{D}}(\vec{r}) \text{d}^3 x - \frac{1}{4\pi} \oint_S V \frac{\partial G_{\text{D}}}{\partial n} \text{d} S$
 $V(\vec{r}) = \langle V \rangle_S + \int \rho G_{\text{N}}(\vec{r}) \text{d}^3 x + \frac{1}{4\pi} \oint_S \frac{\partial V}{\partial n} G_{\text{N}}(\vec{r}) \text{d} S$
 $\nabla_y^2 G(\vec{x}, \vec{y}) = -4\pi \delta(\vec{x} - \vec{y}); \quad G_{\text{D}}(\vec{x}, \vec{y})|_{\vec{y} \in S} = 0; \quad \frac{\partial G_{\text{N}}}{\partial n} \Big|_{\vec{y} \in S} = -\frac{4\pi}{S}$
 $U_{\text{sphere}} = \frac{3}{5} \frac{Q^2}{R}; \quad \vec{p} = \int \text{d}^3 r \rho \vec{r}; \quad \vec{E}_{\text{dip}} = \frac{3(\vec{p} \hat{r}) \hat{r} - \vec{p}}{r^3}; \quad V_{\text{dip}} = \frac{\vec{p} \cdot \hat{r}}{r^2}$
 $Q_{ij} = \int \text{d}^3 r \rho(\vec{r}) (3r_i r_j - \delta_{ij} r^2); \quad V_{\text{quad}} = \frac{1}{6 r^5} Q_{ij} (3r_i r_j - \delta_{ij} r^2)$
 $V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l - \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta)$

$V(r, \theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \left(A_{lm} r^l + \frac{B_{lm}}{r^{l+1}} \right) Y_{lm}(\theta, \varphi)$
 $\frac{1}{|\vec{r} - \vec{r}'|} = \sum_{l=0}^{\infty} \frac{\min(r, r')^l}{\max(r, r')^{l+1}} P_l \left(\frac{\vec{r} \cdot \vec{r}'}{rr'} \right)$
 $P_l(x) = \frac{1}{2^l l!} \frac{\text{d}^l}{\text{d} x^l} (x^2 - 1)^l; \quad f = \sum_{l=0}^{\infty} c_l P_l : c_l = \frac{2l+1}{2} \int_{-1}^1 f P_l$
 $P_l(1) = 1; \quad \langle P_n | P_m \rangle = \frac{2\delta_{nm}}{2n+1}; \quad \langle Y_{lm} | Y_{l'm'} \rangle = \delta_{ll'} \delta_{mm'}$
 $P_0 = 1; \quad P_1 = x; \quad P_2 = \frac{3x^2 - 1}{2}; \quad Y_{00} = \frac{1}{\sqrt{4\pi}}; \quad Y_{10} = \sqrt{\frac{3}{4\pi}} \cos \theta$
 $Y_{11} = -\sqrt{\frac{3}{8\pi}} \sin \theta e^{i \varphi}; \quad Y_{20} = \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1)$
 $Y_{21} = -\sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{i \varphi}; \quad Y_{22} = \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{2i \varphi}$
 $P_{lm}(x) = \frac{(-1)^m}{2^l l!} (1 - x^2)^{\frac{m}{2}} \frac{\text{d}^{l+m}}{\text{d} x^{l+m}} (x^2 - 1)^l, \quad |m| \leq l$

$$Y_{lm}(\theta, \varphi) = \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} e^{im\varphi} P_{lm}(\cos \theta); Y_{l,-m} = (-1)^m \bar{Y}_{lm}$$

$$P_l\left(\frac{\vec{r}\vec{r}'}{rr'}\right) = \frac{4\pi}{2l+1} \sum_{m=-l}^l \bar{Y}_{lm}(\theta', \varphi') Y_{lm}(\theta, \varphi)$$

Magnetostatics (CGS)

$$\vec{\nabla} \vec{J} = -\frac{\partial \rho}{\partial t} = 0; I = \int \vec{J} d\vec{S}$$

$$\text{solenoid: } B = 4\pi \frac{I_s}{c}$$

$$d\vec{F} = \frac{I d\vec{l}}{c} \times \vec{B} = d^3x \frac{\vec{J}}{c} \times \vec{B}; \vec{F}_q = q \frac{\vec{r}}{c} \times \vec{B}$$

$$d\vec{B} = \frac{I d\vec{l}}{c} \times \frac{\vec{r}}{r^3}; \vec{B}_q = q \frac{\vec{r}}{c} \times \frac{\vec{r}}{r^3}$$

Electromagnetism (CGS)

$$\text{Faraday: } \mathcal{E} = -\frac{1}{c} \frac{d\Phi_B}{dt}; \int d^3x \vec{J} = \dot{\vec{p}}$$

$$\vec{\nabla} \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t}; \vec{\nabla} \vec{E} = 4\pi \rho; \vec{\nabla} \vec{J} = -\frac{\partial \rho}{\partial t}$$

$$\vec{\nabla} \times \vec{B} = 4\pi \frac{\vec{J}}{c} + \frac{1}{c} \frac{\partial \vec{E}}{\partial t}; \vec{\nabla} \vec{B} = 0$$

$$d\vec{F} = d^3x (\rho \vec{E} + \frac{\vec{J}}{c} \times \vec{B}); \vec{F}_q = q(\vec{E} + \frac{\dot{\vec{r}}}{c} \times \vec{B})$$

$$u = \frac{E^2 + B^2}{8\pi}; \vec{S} = \frac{c}{4\pi} \vec{E} \times \vec{B}; \vec{g} = \frac{\vec{S}}{c^2}$$

$$\mathbf{T}^E = \frac{1}{4\pi} (\vec{E} \otimes \vec{E} - \frac{1}{2} E^2); \mathbf{T} = \mathbf{T}^E + \mathbf{T}^B$$

$$-\frac{\partial u}{\partial t} = \vec{J} \vec{E} + \vec{\nabla} \vec{S}; -\frac{\partial \vec{g}}{\partial t} = \rho \vec{E} + \frac{\vec{J}}{c} \times \vec{B} - \vec{\nabla} \mathbf{T}$$

$$\vec{B} = \vec{\nabla} \times \vec{A}; \vec{E} = -\vec{\nabla} \phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t}$$

$$-\nabla^2 \phi - \frac{1}{c} \frac{\partial}{\partial t} \vec{\nabla} \vec{A} = 4\pi \rho$$

$$\vec{\nabla} (\vec{\nabla} \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t}) - \nabla^2 \vec{A} + \frac{1}{c} \frac{\partial^2 \vec{A}}{\partial t^2} = 4\pi \frac{\vec{J}}{c}$$

$$(\phi, \vec{A}) \cong (\phi - \frac{1}{c} \frac{\partial \chi}{\partial t}, \vec{A} + \vec{\nabla} \chi)$$

E.M. in matter (CGS)

$$\vec{\nabla} \vec{D} = 4\pi \rho_{\text{ext}}; \vec{\nabla} \times \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t}$$

$$\vec{\nabla} \vec{B} = 0; \vec{\nabla} \times \vec{H} = 4\pi \frac{\vec{J}_{\text{ext}}}{c} + \frac{1}{c} \frac{\partial \vec{D}}{\partial t}$$

$$\vec{P} = \frac{\langle \vec{p} \rangle}{V}; \vec{M} = \frac{\langle \vec{m} \rangle}{V}$$

$$\rho_{\text{pol}} = -\vec{\nabla} \vec{P}; \sigma_{\text{pol}} = \hat{n} \vec{P}; \frac{\vec{J}_{\text{mag}}}{c} = \vec{\nabla} \times \vec{M}$$

$$\vec{D}_{\text{pol}} = \vec{E} + 4\pi \vec{P}; \vec{H}_{\text{mag}} = \vec{B} - 4\pi \vec{M}$$

$$\text{static linear isotropic: } \vec{P} = \chi \vec{E}$$

$$\text{static linear: } P_i = \chi_{ij} E_j$$

$$\text{static linear: } \varepsilon = 1 + 4\pi \chi$$

$$\text{static: } \Delta D_{\perp} = 4\pi \sigma_{\text{ext}}; \Delta E_{\parallel} = 0$$

$$\text{static linear: } u = \frac{1}{8\pi} \vec{E} \vec{D}$$

$$V(r > \text{diam supp } \rho, \theta, \varphi) = \sum_{l=0}^{\infty} \frac{4\pi}{2l+1} \sum_{m=-l}^l q_{lm}[\rho] \frac{Y_{lm}(\theta, \varphi)}{r^{l+1}}$$

$$q_{lm}[\rho] = \int_0^{\infty} r^2 dr \int_0^{2\pi} d\varphi \int_0^{\pi} \sin \theta d\theta r^l \rho(r, \theta, \varphi) \bar{Y}_{lm}(\theta, \varphi)$$

$$\vec{B} = \vec{\nabla} \times \vec{A}; \vec{A} = \int d^3r' \frac{\vec{J}}{c} \frac{1}{|\vec{r} - \vec{r}'|} + \vec{\nabla} A_0$$

$$\vec{B} = \int d^3r' \frac{\vec{J}}{c} \times \frac{\vec{r} - \vec{r}'}{|\vec{r} - \vec{r}'|^3}$$

$$\varphi = \frac{I}{c} \Omega, \vec{B} = -\vec{\nabla} \varphi$$

$$\vec{\nabla} \vec{A} = 0 \rightarrow \nabla^2 \vec{A} = -4\pi \frac{\vec{J}}{c}$$

$$(\phi, \vec{A}) = \int d^3r' \frac{(\rho, \frac{\vec{J}}{c})(\vec{r}', t - \frac{1}{c} |\vec{r} - \vec{r}'|)}{|\vec{r} - \vec{r}'|}$$

$$\text{Coulomb gauge: } \vec{\nabla} \vec{A} = 0$$

$$\text{Lorenz gauge: } \vec{\nabla} \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t} = 0$$

$$\vec{E}'_{\parallel} = \vec{E}_{\parallel}; \vec{B}'_{\parallel} = \vec{B}_{\parallel}$$

$$\vec{E}'_{\perp} = \gamma(\vec{E}_{\perp} + \frac{\vec{v}}{c} \times \vec{B})$$

$$\vec{B}'_{\perp} = \gamma(\vec{B}_{\perp} - \frac{\vec{v}}{c} \times \vec{E})$$

$$\text{plane wave: } \begin{cases} \vec{E} = \vec{E}_0 e^{i(\vec{k}\vec{r} - \omega t)} \\ \vec{B} = \hat{k} \times \vec{E} \\ \omega = ck \end{cases}$$

$$\vec{B}_{\text{diprad}} = \frac{1}{c^2} \frac{\ddot{\vec{p}} \times \hat{r}}{r} \Big|_{t_{\text{rit}}}; \vec{E}_{\text{diprad}} = \vec{B}_{\text{diprad}} \times \hat{r}$$

$$\text{Larmor: } P_{\text{diprad}} = \frac{2}{3c^3} |\ddot{\vec{p}}|^2$$

$$\vec{A}_{\text{dm}} = \frac{1}{c} \frac{\dot{\vec{m}} \times \hat{r}}{r} \Big|_{t_{\text{rit}}}$$

$$\Delta U_{\text{dielectric}} = \frac{1}{2} \int d^3r \vec{P} \vec{E}_0$$

$$\text{plane capacitor: } C = \frac{\varepsilon}{4\pi} \frac{S}{d}$$

$$\text{atomic polarizability: } \vec{p} = \alpha \vec{E}$$

$$\text{non-interacting gas: } \vec{p} = \alpha \vec{E}_0; \chi = n\alpha$$

$$\text{hom. cubic isotropic: } \chi = \frac{1}{\frac{1}{n\alpha} - \frac{4\pi}{3}}$$

$$\text{Clausius-Mossotti: } \frac{\varepsilon-1}{\varepsilon+2} = \frac{4\pi}{3} n\alpha$$

$$\chi = \frac{4\pi}{3} \frac{n p_0^2}{kT}; \vec{E}_e = \vec{E} + \frac{4\pi}{3} \vec{P}$$

$$\vec{J} \vec{E} = -\vec{\nabla} \left(\frac{c}{4\pi} \vec{E} \times \vec{H} \right) - \frac{1}{4\pi} \left(\vec{E} \frac{\partial \vec{D}}{\partial t} + \vec{H} \frac{\partial \vec{B}}{\partial t} \right)$$

$$n = \sqrt{\varepsilon \mu}; k = n \frac{\omega}{c}$$

$$\text{plane wave: } B = nE$$

$$\vec{J}_c = \sigma \vec{E}; \varepsilon_{\sigma} = 1 + i \frac{4\pi \sigma}{\omega}$$

$$\vec{\nabla} \vec{B} = 0; \vec{\nabla} \times \vec{B} = 4\pi \frac{\vec{J}}{c}; \oint \vec{B} d\vec{l} = 4\pi \frac{I}{c}$$

$$\vec{m} = \frac{1}{2} \int d^3r' (\vec{r}' \times \frac{\vec{J}}{c}) = \frac{1}{2c} \frac{q}{m} \vec{L} = \frac{S}{c}$$

$$\vec{A}_{\text{dm}} = \frac{\vec{m} \times \vec{r}}{r^3}; \vec{\tau} = \vec{m} \times \vec{B}$$

$$\vec{F}_{\text{dm dm}} = -\vec{\nabla}_R \frac{\vec{m} \vec{m}' - 3(\vec{m} \hat{R})(\vec{m}' \hat{R})}{R^3}$$

$$\text{loop axis: } \vec{B} = \hat{z} \frac{2\pi R^2}{(z^2 + R^2)^{3/2}} \frac{I}{c}$$

$$\text{L.W.: } (\phi, \vec{A}) = \frac{q(1, \frac{\vec{v}}{c})}{[r - \frac{\vec{v}\vec{r}}{c}]_{t_{\text{rit}}}}; t_{\text{rit}} = t - \frac{r}{c} \Big|_{t_{\text{rit}}}$$

$$A^{\mu} = (\phi, \vec{A}); J^{\mu} = (c\rho, \vec{J})$$

$$\text{Lorenz gauge: } \partial_{\alpha} A^{\alpha} = 0$$

$$F^{\mu\nu} = \partial^{\mu} A^{\nu} - \partial^{\nu} A^{\mu}$$

$$F^{\mu\nu} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -B_z & B_y \\ E_y & B_z & 0 & -B_x \\ E_z & -B_y & B_x & 0 \end{pmatrix}$$

$$\mathcal{F}^{\mu\nu} = \frac{1}{2} \varepsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}$$

$$\partial_{\alpha} F^{\alpha\nu} = 4\pi \frac{J^{\nu}}{c}; \partial_{\alpha} \mathcal{F}^{\alpha\nu} = 0; \frac{dp^{\mu}}{d\tau} = q F^{\mu\alpha} v_{\alpha}$$

$$F^{\alpha\beta} F_{\alpha\beta} = 2(B^2 - E^2); F^{\alpha\beta} \mathcal{F}_{\alpha\beta} = 4\vec{E} \vec{B}$$

$$\Theta^{\mu\nu} = \frac{1}{4\pi} (F^{\mu}_{\alpha} F^{\alpha\nu} + \frac{1}{4} g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta})$$

$$\Theta^{\mu\nu} = \begin{pmatrix} u & c\vec{g} \\ c\vec{g} & -\mathbf{T} \end{pmatrix}$$

$$\partial_{\alpha} \Theta^{\alpha\nu} = \frac{1}{c} J_{\alpha} F^{\alpha\nu}$$

$$\omega_p^2 = 4\pi \frac{nq^2}{m}; \omega_{\text{cyclo}} = \frac{qB}{mc}$$

$$\text{I: } u = \frac{1}{8\pi} (\vec{E} \vec{D} + \vec{H} \vec{B})$$

$$\text{I: } \langle S_z \rangle = \frac{c}{n} \langle u \rangle$$

$$\text{II: } u = \frac{1}{8\pi} \left(\frac{\partial}{\partial \omega} (\varepsilon_{\omega} \omega) E^2 + \frac{\partial}{\partial \omega} (\mu_{\omega} \omega) H^2 \right)$$

$$\text{II: } \langle S_z \rangle = v_g \langle u \rangle; v_g = \frac{\partial \omega}{\partial k}$$

$$\text{III: } \langle W \rangle = \frac{\omega}{8\pi} (\text{Im } \varepsilon_{\omega} |E_0^2| + \text{Im } \mu_{\omega} |H_0^2|)$$

$$\text{Fresnel TE (S): } \frac{E_t}{E_i} = \frac{2}{1 + \frac{k_{tz}}{k_{iz}}}; \frac{E_r}{E_i} = \frac{1 - \frac{k_{tz}}{k_{iz}}}{1 + \frac{k_{tz}}{k_{iz}}}$$

$$\text{TM (P): } \frac{E_t}{E_i} = \frac{2}{\frac{n_2}{n_1} + \frac{n_1}{n_2} \frac{k_{tz}}{k_{iz}}}; \frac{E_r}{E_i} = \frac{\frac{n_2}{n_1} - \frac{n_1}{n_2} \frac{k_{tz}}{k_{iz}}}{\frac{n_2}{n_1} + \frac{n_1}{n_2} \frac{k_{tz}}{k_{iz}}}$$

$$\text{Fresnel: } k_{tz} = \pm \sqrt{\varepsilon_2 \left(\frac{\omega}{c} \right)^2 - k_x^2}, \text{Im } k_{tz} > 0$$