

TWISTED RESTRICTED CONFORMAL BLOCKS OF VERTEX OPERATOR ALGEBRAS

MATH-PHYSICS JOINT SEMINAR

Xu Gao

Oct. 31st, 2024

Tongji University

- 1 Conformal blocks
- 2 Twisted conformal blocks
- 3 Twisted fusion rules
- 4 Restricted conformal blocks
- 5 Twisted Zhu's algebra and coherence

CONFORMAL BLOCKS

Idea: One have various *sectors* attached to a *pointed curve* (which gives rise to a structure named *conformal blocks*), and study how they interact as the pointed curve varies (via the *correlators*).

Idea: One have various *sectors* attached to a *pointed curve* (which gives rise to a structure named *conformal blocks*), and study how they interact as the pointed curve varies (via the *correlators*).

Representation theoretical interpretation:

Vertex operator algebra \rightsquigarrow Conformal field theory

Idea: One have various *sectors* attached to a *pointed curve* (which gives rise to a structure named *conformal blocks*), and study how they interact as the pointed curve varies (via the *correlators*).

Representation theoretical interpretation:

Vertex operator algebra \rightsquigarrow Conformal field theory

- modules of the vertex operator algebra \rightsquigarrow sectors;

Idea: One have various *sectors* attached to a *pointed curve* (which gives rise to a structure named *conformal blocks*), and study how they interact as the pointed curve varies (via the *correlators*).

Representation theoretical interpretation:

Vertex operator algebra \rightsquigarrow Conformal field theory

- modules of the vertex operator algebra \rightsquigarrow sectors;
- vertex/intertwining operators \rightsquigarrow correlators.

Idea: One have various *sectors* attached to a *pointed curve* (which gives rise to a structure named *conformal blocks*), and study how they interact as the pointed curve varies (via the *correlators*).

Representation theoretical interpretation:

Vertex operator algebra \rightsquigarrow Conformal field theory

- modules of the vertex operator algebra \rightsquigarrow sectors;
- vertex/intertwining operators \rightsquigarrow correlators.

Borcherds 1986; Frenkel, Lepowsky, and Meurman 1988; Beilinson, Feigin, and Mazur 1991; Dong and Lepowsky 1993; Frenkel, Huang, and Lepowsky 1993; Lepowsky and Li 2004; Zhu 1994, 1996; Frenkel and Ben-Zvi 2004; Nagatomo and Tsuchiya 2005

DEFINITION: A *vertex operator algebra* (VOA) is a graded vector space $V = \bigoplus_{k=0}^{\infty} V_k$ with a *vacuum vector* $\mathbf{1} \in V_0$, a *Virasoro element* $\omega \in V_2$, and a *vertex operator*

$$Y(-, z): V \longrightarrow \text{End}(V)[[z^{\pm 1}]],$$

satisfying certain axioms.

DEFINITION: A *vertex operator algebra* (VOA) is a graded vector space $V = \bigoplus_{k=0}^{\infty} V_k$ with a *vacuum vector* $\mathbf{1} \in V_0$, a *Virasoro element* $\omega \in V_2$, and a *vertex operator*

$$Y(-, z): V \longrightarrow \text{End}(V)[[z^{\pm 1}]],$$

satisfying certain axioms.

DEFINITION: An *(admissible) module* of a vertex operator algebra V is a graded vector space $M = \bigoplus_{k=0}^{\infty} M(k)$ with a *vertex operator*

$$Y_M(-, z): V \longrightarrow \text{End}(M)[[z^{\pm 1}]]$$

satisfying certain axioms.

DEFINITION: An *(admissible) module* of a vertex operator algebra V is a graded vector space $M = \bigoplus_{k=0}^{\infty} M(k)$ with a *vertex operator*

$$Y_M(-, z): V \longrightarrow \text{End}(M)[[z^{\pm 1}]]$$

satisfying certain axioms.

DEFINITION: An *intertwining operator* among V -modules M^1 , M^2 , and M^3 is a linear map

$$I(-, z): M^1 \longrightarrow \text{Hom}(M^2, M^3)[[z^{\pm 1}]]z^{-h}$$

that is compatible with the V -module structures.

Algebro-geometric interpretation:

Given an n -tuple of V -modules, one constructs (and studies) a **vector bundle** \mathbb{V} with a **projectively flat connection** ∇ on the moduli space $\overline{\mathcal{M}}_{g,n}$ of stable pointed curves.

Algebraic-geometric interpretation:

Given an n -tuple of V -modules, one constructs (and studies) a **vector bundle** \mathbb{V} with a **projectively flat connection** ∇ on the moduli space $\overline{\mathcal{M}}_{g,n}$ of stable pointed curves.

DEFINITION: Given a stable n -pointed curve (C, \mathbf{p}_\bullet) and an n -tuple of V -modules M^1, \dots, M^n , a **conformal block** associated to these data is a linear functional

$$M^1 \otimes \dots \otimes M^n \longrightarrow \mathbb{C}$$

invariant under the action of the **chiral Lie algebra** $\mathcal{L}_{(C, \mathbf{p}_\bullet)}(V)$.

DEFINITION: Given a stable n -pointed curve (C, \mathbf{p}_\bullet) and an n -tuple of V -modules M^1, \dots, M^n , a **conformal block** associated to these data is a linear functional

$$M^1 \otimes \dots \otimes M^n \longrightarrow \mathbb{C}$$

invariant under the action of the **chiral Lie algebra** $\mathcal{L}_{(C, \mathbf{p}_\bullet)}(V)$.

THEOREM (Frenkel & Ben-Zvi 04', Nagatomo & Tsuchiya 05'): *The vector space of conformal blocks associated to*

$$(\mathbb{P}^1, \infty, 1, 0, (M^3)', M^1, M^2)$$

is isomorphic to the space of intertwining operators among M^1 , M^2 , and M^3 .

THEOREM (Damiolini, Gibney, and Tarasca 21'–24'): Given:

- V a rational, C_2 -cofinite VOA with $V_0 = \mathbb{C}\mathbf{1}$, and
- M^\bullet an n -tuple of f.g. admissible V -modules,

\rightsquigarrow a vector bundle $[M^\bullet]$ of finite rank with a projectively flat connection ∇ on $\overline{\mathcal{M}}_{g,n}$, whose dual bundle classifies the conformal blocks.

THEOREM (Damiolini, Gibney, and Tarasca 21'–24'): *Those vector bundles $([M^\bullet], \nabla)$ are compatible with various moduli spaces in the sense that:*

THEOREM (Damiolini, Gibney, and Tarasca 21'–24'): *Those vector bundles $([M^\bullet], \nabla)$ are compatible with various moduli spaces in the sense that:*

1. *Propagation of vacua* $[M^\bullet]_{(C, p_\bullet)} \cong [M^\bullet \otimes V]_{(C, p_\bullet \sqcup q)}$;

$$\begin{array}{c} M^1 \quad \dots \quad M^n \\ \bullet \quad \dots \quad \bullet \\ \hline p_1 \quad \dots \quad p_n \end{array} \cong \begin{array}{c} M^1 \quad \dots \quad M^n \quad V \\ \bullet \quad \dots \quad \bullet \quad \bullet \\ \hline p_1 \quad \dots \quad p_n \quad q \end{array}$$

THEOREM (Damiolini, Gibney, and Tarasca 21'–24'): *Those vector bundles $([M^\bullet], \nabla)$ are compatible with various moduli spaces in the sense that:*

1. *Propagation of vacua* $[M^\bullet]_{(C, p_\bullet)} \cong [M^\bullet \otimes V]_{(C, p_\bullet \sqcup q)}$;
2. *Decomposition* $[M^\bullet \otimes N^\bullet]_{(C \sqcup C', p_\bullet \sqcup p'_\bullet)} \cong [M^\bullet]_{(C, p_\bullet)} \otimes [N^\bullet]_{(C', p'_\bullet)}$;

$$\begin{array}{ccc}
 M^1 & \dots & M^n \\
 \bullet & \text{---} & \bullet \\
 p_1 & \dots & p_n
 \end{array}
 \qquad
 \begin{array}{ccc}
 N^1 & \dots & N^m \\
 \bullet & \text{---} & \bullet \\
 p'_1 & \dots & p'_m
 \end{array}$$


THEOREM (Damiolini, Gibney, and Tarasca 21'–24'): Those vector bundles $([M^\bullet], \nabla)$ are compatible with various moduli spaces in the sense that:

1. *Propagation of vacua* $[M^\bullet]_{(C, p_\bullet)} \cong [M^\bullet \otimes V]_{(C, p_\bullet \sqcup q)}$;
2. *Decomposition* $[M^\bullet \otimes N^\bullet]_{(C \sqcup C', p_\bullet \sqcup p'_\bullet)} \cong [M^\bullet]_{(C, p_\bullet)} \otimes [N^\bullet]_{(C', p'_\bullet)}$;
3. *Factorization* $[M^\bullet]_{(C, p_\bullet)} \cong \bigoplus_W [M^\bullet \otimes W \otimes W']_{(\tilde{C}, p_\bullet \sqcup q_\pm)}$



THEOREM (Damiolini, Gibney, and Tarasca 21'–24'): *Those vector bundles $([M^\bullet], \nabla)$ are compatible with various moduli spaces in the sense that:*

1. *Propagation of vacua* $[M^\bullet]_{(C, p_\bullet)} \cong [M^\bullet \otimes V]_{(C, p_\bullet \sqcup q)}$;
2. *Decomposition* $[M^\bullet \otimes N^\bullet]_{(C \sqcup C', p_\bullet \sqcup p'_\bullet)} \cong [M^\bullet]_{(C, p_\bullet)} \otimes [N^\bullet]_{(C', p'_\bullet)}$;
3. *Factorization* $[M^\bullet]_{(C, p_\bullet)} \cong \bigoplus_W [M^\bullet \otimes W \otimes W']_{(\tilde{C}, p_\bullet \sqcup q_\pm)}$
4. *Formal smoothing* $[M^\bullet[[q]]]_{(C, p_\bullet)} \cong [M^\bullet]_{(C_0, p_\bullet(0))}[[q]]$



TWISTED CONFORMAL BLOCKS

Idea: One can also have **twisted** sectors attached to a pointed orbifold (which gives rise to *twisted conformal blocks*), and study how they interact as the orbifold varies (via the *twisted correlators*).

Idea: One can also have **twisted** sectors attached to a pointed orbifold (which gives rise to *twisted conformal blocks*), and study how they interact as the orbifold varies (via the *twisted correlators*).

Representation theoretical interpretation:

Idea: One can also have **twisted** sectors attached to a pointed orbifold (which gives rise to *twisted conformal blocks*), and study how they interact as the orbifold varies (via the *twisted correlators*).

Representation theoretical interpretation:

- *twisted* modules \rightsquigarrow *twisted* sectors;

Idea: One can also have **twisted** sectors attached to a pointed orbifold (which gives rise to *twisted conformal blocks*), and study how they interact as the orbifold varies (via the *twisted correlators*).

Representation theoretical interpretation:

- **twisted** modules \rightsquigarrow *twisted* sectors;
- **twisted** vertex/intertwining operators \rightsquigarrow *twisted* correlators.

Idea : One can also have **twisted** sectors attached to a pointed **orbifold** (which gives rise to *twisted conformal blocks*), and study how they interact as the orbifold varies (via the *twisted correlators*).

Representation theoretical interpretation:

- **twisted** modules \rightsquigarrow *twisted* sectors;
- **twisted** vertex/intertwining operators \rightsquigarrow *twisted* correlators.

Frenkel, Lepowsky, and Meurman 1988; Dong 1994; Xu 1995; Dong, Li, and Mason 1998; Barron, Dong, and Mason 2002; Dong and Xu 2006; Huang 2010; Dong, Ren, and Xu 2017; Huang 2018; Huang and Yang 2019...

DEFINITION: Let g be an automorphism of V having order T . An *(admissible) g -twisted module* of a vertex operator algebra V is a graded vector space $M = \bigoplus_{k \in \frac{1}{T}\mathbb{Z}} M(k)$ with a *vertex operator*

$$Y_M(-, z): V \longrightarrow \text{End}(M)[[z^{\pm 1/T}]]$$

satisfying certain axioms.

DEFINITION: Let g be an automorphism of V having order T . An *(admissible) g -twisted module* of a vertex operator algebra V is a graded vector space $M = \bigoplus_{k \in \frac{1}{T}\mathbb{Z}} M(k)$ with a *vertex operator*

$$Y_M(-, z): V \longrightarrow \text{End}(M)[[z^{\pm 1/T}]]$$

satisfying certain axioms.

DEFINITION: An *twisted intertwining operator* among twisted V -modules M^1 , M^2 , and M^3 is a linear map

$$I(-, z): M^1 \longrightarrow \text{Hom}(M^2, M^3)\{z\}$$

that is compatible with the twisted V -module structures.

Algebro-geometric interpretation:

Given an n -tuple of **twisted** V -modules, one constructs (and studies) a **vector bundle** \mathbb{V} with a **projectively flat connection** ∇ on the moduli space $\overline{\mathcal{M}}_{g,n}^G$ of stable **G -covers**.

Algebra-geometric interpretation:

Given an n -tuple of **twisted** V -modules, one constructs (and studies) a **vector bundle** \mathbb{V} with a **projectively flat connection** ∇ on the moduli space $\overline{\mathcal{M}}_{g,n}^G$ of stable **G -covers**.

DEFINITION: Given a stable G -cover $(\mathfrak{X}, \mathbf{p}_\bullet)$ and an n -tuple of twisted V -modules M^1, \dots, M^n , a **twisted conformal block** associated to these data is a linear functional

$$M^1 \otimes \dots \otimes M^n \longrightarrow \mathbb{C}$$

invariant under the action of the **chiral Lie algebra** $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$.

DEFINITION: Given a stable G -cover $(\mathfrak{X}, \mathbf{p}_\bullet)$ and an n -tuple of twisted V -modules M^1, \dots, M^n , a *twisted conformal block* associated to these data is a linear functional

$$M^1 \otimes \dots \otimes M^n \longrightarrow \mathbb{C}$$

invariant under the action of the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$.

Remark. \mathfrak{X} is a stack, points of it are equipped with cyclic stabilizers. For each point \mathbf{p} , the sector is given by a g -twisted V -module, where g generates its stabilizer.

DEFINITION: A *stable G -cover* $(\mathfrak{X}, \mathbf{p}_\bullet)$ is


- a stacky curve \mathfrak{X} ,
- a principal G -bundle $\pi: P \rightarrow \mathfrak{X}$, and
- an n -tuple of points $\mathbf{p}_\bullet = (p_1, \dots, p_n)$ on \mathfrak{X} ,

satisfying certain stability conditions.

DEFINITION: A *stable G -cover* $(\mathfrak{X}, \mathbf{p}_\bullet)$ is

- a stacky curve \mathfrak{X} ,
- a principal G -bundle $\pi: P \rightarrow \mathfrak{X}$, and
- an n -tuple of points $\mathbf{p}_\bullet = (p_1, \dots, p_n)$ on \mathfrak{X} ,

satisfying certain stability conditions.



Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$?

Recall that (Damianini, Gibney, and Tarasca 21'–24'):
a VOA V

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$?

Recall that (Damiolini, Gibney, and Tarasca 21'–24'):

a VOA $V \rightsquigarrow$ an ind-bundle \mathcal{V} with flat connection on a nodal curve C

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .
- a morphism $G \rightarrow \text{Aut}(V) \implies (\mathcal{V}, \nabla)$ is G -equivariant.

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .
- a morphism $G \rightarrow \text{Aut}(V) \implies (\mathcal{V}, \nabla)$ is G -equivariant.
- (\mathcal{V}, ∇) descends to (\mathcal{V}^G, ∇) on \mathfrak{X} .

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .
- a morphism $G \rightarrow \text{Aut}(V) \implies (\mathcal{V}, \nabla)$ is G -equivariant.
- (\mathcal{V}, ∇) descends to (\mathcal{V}^G, ∇) on \mathfrak{X} .
- $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V) := h_{\text{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathbf{p}_\bullet) = H^0(\mathfrak{X} \setminus \mathbf{p}_\bullet, \mathcal{V}^G \otimes \omega_{\mathfrak{X}} / \text{Im } \nabla)$.

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .
- a morphism $G \rightarrow \text{Aut}(V) \implies (\mathcal{V}, \nabla)$ is G -equivariant.
- (\mathcal{V}, ∇) descends to (\mathcal{V}^G, ∇) on \mathfrak{X} .
- $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V) := \text{h}_{\text{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathfrak{p}_{\bullet}) = H^0(\mathfrak{X} \setminus \mathfrak{p}_{\bullet}, \mathcal{V}^G \otimes \omega_{\mathfrak{X}} / \text{Im } \nabla)$.

Q: How does $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V)$ act on (twisted) V -modules?

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .
- a morphism $G \rightarrow \text{Aut}(V) \implies (\mathcal{V}, \nabla)$ is G -equivariant.
- (\mathcal{V}, ∇) descends to (\mathcal{V}^G, ∇) on \mathfrak{X} .
- $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V) := \text{h}_{\text{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathfrak{p}_{\bullet}) = H^0(\mathfrak{X} \setminus \mathfrak{p}_{\bullet}, \mathcal{V}^G \otimes \omega_{\mathfrak{X}} / \text{Im } \nabla)$.

Q: How does $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V)$ act on (twisted) V -modules?

$$\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V) \xrightarrow{\text{res}} \mathcal{L}_{\mathfrak{p}}(V) (:= \text{h}_{\text{dR}}(\mathcal{V}^G, \nabla)(D_{\mathfrak{p}}^{\times})).$$

Q: How to obtain the *chiral Lie algebra* $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V)$?

- a VOA $V \rightsquigarrow (\mathcal{V}, \nabla)$ on P .
- a morphism $G \rightarrow \text{Aut}(V) \implies (\mathcal{V}, \nabla)$ is G -equivariant.
- (\mathcal{V}, ∇) *descends to* (\mathcal{V}^G, ∇) on \mathfrak{X} .
- $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V) := \mathfrak{h}_{\text{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathfrak{p}_{\bullet}) = H^0(\mathfrak{X} \setminus \mathfrak{p}_{\bullet}, \mathcal{V}^G \otimes \omega_{\mathfrak{X}} / \text{Im } \nabla)$.

Q: How does $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V)$ act on (twisted) V -modules?

$$\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V) \xrightarrow{\text{res}} \mathcal{L}_{\mathfrak{p}}(V) (:= \mathfrak{h}_{\text{dR}}(\mathcal{V}^G, \nabla)(D_{\mathfrak{p}}^{\times})).$$

The latter is \cong to the Lie algebra $L_g(V)$ of *g -twisted* vertex modes, where g is a generator of the stabilizer of \mathfrak{p} .

TWISTED FUSION RULES

DEFINITION: The *(twisted) fusion rule* is the dimension of the vector space of (twisted) intertwining operators.

DEFINITION: The *(twisted) fusion rule* is the dimension of the vector space of (twisted) intertwining operators.

THEOREM (G., Liu & Zhu 24'): *The vector space of twisted conformal blocks associated to*

$$([C_w/\mu_T], \infty, w, 0, (M^3)', M^1, M^2)$$

is isomorphic to the space of twisted intertwining operators among the untwisted module M^1 and g -twisted modules M^2 and M^3 .

DEFINITION: The *(twisted) fusion rule* is the dimension of the vector space of (twisted) intertwining operators.

THEOREM (G., Liu & Zhu 24'): *The vector space of twisted conformal blocks associated to*

$$([C_w/\mu_T], \infty, w, 0, (M^3)', M^1, M^2)$$

is isomorphic to the space of twisted intertwining operators among the untwisted module M^1 and g -twisted modules M^2 and M^3 .

Note: $[C_w/\mu_T]$ is totally ramified.

DEFINITION: The *(twisted) fusion rule* is the dimension of the vector space of (twisted) intertwining operators.

PROPOSAL: *The vector space of twisted intertwining operators among M^1 , M^2 , and M^3 is given by the space of twisted conformal blocks associated to the data*

$$(\mathbb{P}_{g_3^{-1}, g_1, g_2}^1, \infty, 1, 0, (M^3)', M^1, M^2),$$

where $g_1 g_2 = g_3$.

DEFINITION: The *(twisted) fusion rule* is the dimension of the vector space of (twisted) intertwining operators.

Q: How to compute the twisted fusion rules?

DEFINITION: The *(twisted) fusion rule* is the dimension of the vector space of (twisted) intertwining operators.

Q: How to compute the twisted fusion rules? More generally, how to compute the dimension of a space of twisted conformal blocks?

RESTRICTED CONFORMAL BLOCKS

Recall: $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_{\bullet})}(V) = \mathfrak{h}_{\mathrm{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathfrak{p}_{\bullet})$.

Recall: $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet)}(V) = \mathbf{h}_{\mathrm{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathfrak{p}_\bullet)$.

We can constrain it by twisting the \mathcal{D} -module (\mathcal{V}^G, ∇) by a divisor Δ .

$$\Delta_{m_\bullet} := - \sum_{i=1}^n (L_{(0)} - m_i - 1)[\mathfrak{p}_i].$$

Recall: $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V) = \mathbf{h}_{\mathrm{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathbf{p}_\bullet)$.

We can constrain it by twisting the \mathcal{D} -module (\mathcal{V}^G, ∇) by a divisor Δ .

$$\Delta_{m_\bullet} := - \sum_{i=1}^n (L_{(0)} - m_i - 1)[\mathbf{p}_i].$$

NOTATION: For any $m_\bullet \in (\mathbb{Z} \cup \{\infty\})^n$, denote

$$\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)_{\leq m_\bullet} := \mathbf{h}_{\mathrm{dR}}(\mathcal{V}^G(\Delta_{m_\bullet}), \nabla)(\mathfrak{X}),$$

Recall: $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V) = \mathbf{h}_{\mathrm{dR}}(\mathcal{V}^G, \nabla)(\mathfrak{X} \setminus \mathbf{p}_\bullet)$.

We can constrain it by twisting the \mathcal{D} -module (\mathcal{V}^G, ∇) by a divisor Δ .

$$\Delta_{m_\bullet} := - \sum_{i=1}^n (L_{(0)} - m_i - 1)[\mathbf{p}_i].$$

NOTATION: For any $m_\bullet \in (\mathbb{Z} \cup \{\infty\})^n$, denote

$$\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)_{\leq m_\bullet} := \mathbf{h}_{\mathrm{dR}}(\mathcal{V}^G(\Delta_{m_\bullet}), \nabla)(\mathfrak{X}),$$

Remark. $(\mathcal{V}^G(\Delta_{m_\bullet}), \nabla) = (\mathcal{V}(\pi^* \Delta_{m_\bullet} - \mathfrak{K}), \nabla)^G$.

| EXAMPLE: $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$ and $\mathbf{p}_\bullet = (1, 0, \infty)$

EXAMPLE: $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$ and $\mathbf{p}_\bullet = (1, 0, \infty) \rightsquigarrow \pi^* \Delta_{m_\bullet} - \mathfrak{R}$ reads

$$-(T(L_{(0)} - m_0) - 1)\pi^*[0] - (L_{(0)} - m_1 - 1)\pi^*[1] - (T(L_{(0)} - m_\infty) - 1)\pi^*[\infty].$$

EXAMPLE: $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$ and $\mathbf{p}_\bullet = (1, 0, \infty) \rightsquigarrow \pi^* \Delta_{m_\bullet} - \mathfrak{R}$ reads

$$-(T(L_{(0)} - m_0) - 1)\pi^*[0] - (L_{(0)} - m_1 - 1)\pi^*[1] - (T(L_{(0)} - m_\infty) - 1)\pi^*[\infty].$$

On the other hand, $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V) = V \otimes \mathbb{C}[z^{\pm 1/T}, (z-1)^{-1}] dz / \text{Im } \nabla$.

EXAMPLE: $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$ and $\mathbf{p}_\bullet = (1, 0, \infty) \rightsquigarrow \pi^* \Delta_{m_\bullet} - \mathfrak{R}$ reads

$$-(T(L_{(0)} - m_0) - 1)\pi^*[0] - (L_{(0)} - m_1 - 1)\pi^*[1] - (T(L_{(0)} - m_\infty) - 1)\pi^*[\infty].$$

On the other hand, $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V) = V \otimes \mathbb{C}[z^{\pm 1/T}, (z-1)^{-1}] dz / \text{Im } \nabla$.

Hence, $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)_{\leq m_\bullet}$ is spanned by

$$\left\{ \left[u \otimes z^{\frac{r}{T} + a} (z-1)^b dz \right] \mid \begin{array}{l} u \in V_k \cap V^r \\ a \geq k - 1 - \frac{r}{T} - m_0 \\ b \geq k - 1 - \frac{r}{T} - m_1 \\ a + b \leq k - 1 - \frac{r}{T} + m_\infty \end{array} \right\}$$

We consider $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$ and the multi-index $\infty \sqcup m$ asserting ∞ to each \mathbf{p}_i and m to each \mathbf{q}_j .

We consider $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$ and the multi-index $\infty \sqcup m$ asserting ∞ to each \mathbf{p}_i and m to each \mathbf{q}_j .

NOTATION: $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq m} := \mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq \infty \sqcup m}$

We consider $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$ and the multi-index $\infty \sqcup m$ asserting ∞ to each \mathbf{p}_i and m to each \mathbf{q}_j .

NOTATION: $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq m} := \mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq \infty \sqcup m}$

LEMMA: Suppose M is a (twisted) V -module attached to some \mathbf{q}_j . Then, $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{<0} (:= \bigcup_{m < 0} \mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq m})$ acts trivially on its bottom level $M(0)$. Hence, $M(0)$ is an $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_0$ -module.

We consider $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$ and the multi-index $\infty \sqcup m$ asserting ∞ to each \mathbf{p}_i and m to each \mathbf{q}_j .

NOTATION: $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq m} := \mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq \infty \sqcup m}$

LEMMA: Suppose M is a (twisted) V -module attached to some \mathbf{q}_j . Then, $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{<0} (:= \bigcup_{m < 0} \mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq m})$ acts trivially on its bottom level $M(0)$. Hence, $M(0)$ is an $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_0$ -module. Conversely, any $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_0$ -module (and any $A_g(V)$ -module) can be viewed as an $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq 0}$ -module.

DEFINITION: Given a stable G -cover $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$ and a pair of families of (twisted) V -modules M^\bullet and $A_{g_\diamond}(V)$ -modules U^\diamond , a *twisted restricted conformal block* associated to these data is a linear functional

$$M^1 \otimes \cdots \otimes M^m \otimes U^1 \otimes \cdots \otimes U^n \longrightarrow \mathbb{C}$$

invariant under the action of the Lie subalgebra $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)_{\leq 0}$.

DEFINITION: Given a stable G -cover $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$ and a pair of families of (twisted) V -modules M^\bullet and $A_{g_\diamond}(V)$ -modules U^\diamond , a *twisted restricted conformal block* associated to these data is a linear functional

$$M^1 \otimes \cdots \otimes M^m \otimes U^1 \otimes \cdots \otimes U^n \longrightarrow \mathbb{C}$$

invariant under the action of the Lie subalgebra $\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet)}(V)_{\leq 0}$.

EXAMPLE: Given an $(m+n)$ -tuple of (twisted) V -module $M^\bullet \sqcup N^\diamond$,

$$\mathrm{Conf}(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond, M^\bullet \sqcup N^\diamond) \quad \mathrm{ResConf}(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond, M^\bullet \sqcup N^\diamond(0)).$$

EXAMPLE: Given an $(m + n)$ -tuple of (twisted) V -module $M^\bullet \sqcup N^\diamond$,
 $\text{Conf}(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond, M^\bullet \sqcup N^\diamond) \xrightarrow{\pi} \text{ResConf}(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond, M^\bullet \sqcup N^\diamond(0)).$

EXAMPLE: Given an $(m + n)$ -tuple of (twisted) V -module $M^\bullet \sqcup N^\diamond$,
 $\text{Conf}(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond, M^\bullet \sqcup N^\diamond) \xrightarrow{\pi} \text{ResConf}(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond, M^\bullet \sqcup N^\diamond(0)).$

THEOREM (G., Liu, Zhu 24'): If N^\diamond are lowest-weight modules, then π is *injective*. If N^\diamond are further the *generalized Verma modules* of their bottom levels, then π is an *isomorphism*.

EXAMPLE (G., Liuu, and Zhu 23'): Let g be a VOA-automorphism of V of order T and consider the data:

- $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$, $\mathbf{p}_\bullet = (1)$, and $\mathbf{q}_\diamond = (0, \infty)$;
- attach a **untwisted** V -module M^1 at $\mathbf{1}$;
- attach a left $A_g(V)$ -module U^2 at $\mathbf{0}$;
- attach a left $A_{g^{-1}}(V)$ -module U^3 at ∞ ;

EXAMPLE (G., Liiu, and Zhu 23'): Let g be a VOA-automorphism of V of order T and consider the data:

- $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$, $\mathbf{p}_\bullet = (1)$, and $\mathbf{q}_\diamond = (0, \infty)$;
- attach a untwisted V -module M^1 at 1;
- attach a left $A_g(V)$ -module U^2 at 0;
- attach a left $A_{g^{-1}}(V)$ -module U^3 at ∞ ;

Then, the space of twisted restricted conformal block is the dual of $U^2 \otimes M^1 \otimes U^3/J$, where J has an explicit description.

EXAMPLE (G., Liiu, and Zhu 23'): Let g be a VOA-automorphism of V of order T and consider the data:

- $\mathfrak{X} = [\mathbb{P}^1/\mu_T]$, $\mathbf{p}_\bullet = (1)$, and $\mathbf{q}_\diamond = (0, \infty)$;
- attach a untwisted V -module M^1 at 1;
- attach a left $A_g(V)$ -module U^2 at 0;
- attach a left $A_{g^{-1}}(V)$ -module U^3 at ∞ ;

Then, the space of **twisted restricted conformal block** is the dual of $U^2 \otimes M^1 \otimes U^3/J$, where J has an explicit description. From which, the space is the dual of

$$U^3 \otimes_{A_g(V)} A_g(M^1) \otimes_{A_g(V)} U^2.$$

TWISTED ZHU'S ALGEBRA AND COHERENCE

Basic set-up: $(\mathfrak{X}, p_{\bullet} \sqcup q_{\diamond})$.

Basic set-up: $(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)$.

LEMMA: $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)_{\leq \star}$ form an exhaustive, separated, and split filtration on $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)$.

Basic set-up: $(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)$.

LEMMA: $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)_{\leq \star}$ form an exhaustive, separated, and split filtration on $\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)$.

LEMMA (Damiolini, Gibney, and Krashen 23'): From a split-filtered associative algebra $(U_{\leq \star}, U_\star)$, one ends up with a complete semi-normed split-filtered associative algebra $(\widehat{U}_{\leq \star}^f, \widehat{U}_\star^g)$. In such a pair, any homogeneous ideal $I \triangleleft \widehat{U}_\star^g$ induces a pair of closed ideals $(\overline{I}^f, \overline{I}^g)$, and we obtain another pair of semi-normed split-filtered associative algebra $(\widehat{U}_{\leq \star}^f / \overline{I}^f, \widehat{U}_\star^g / \overline{I}^g)$.

LEMMA (Damiolini, Gibney, and Krashen 23'): *From a split-filtered associative algebra $(U_{\leqslant \star}, U_{\star})$, one ends up with a complete semi-normed split-filtered associative algebra $(\widehat{U}_{\leqslant \star}^f, \widehat{U}_{\star}^g)$. In such a pair, any homogeneous ideal $I \triangleleft \widehat{U}_{\star}^g$ induces a pair of closed ideals (\bar{I}^f, \bar{I}^g) , and we obtain another pair of semi-normed split-filtered associative algebra $(\widehat{U}_{\leqslant \star}^f / \bar{I}^f, \widehat{U}_{\star}^g / \bar{I}^g)$.*

DEFINITION: Take $U_{\leqslant \star} = U\left(\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_{\bullet} \sqcup \mathbf{q}_{\diamond})}(V)_{\leqslant \star}\right)$ and I to be generated by the Jacobi relations. The resulted semi-normed split-filtered associative algebra $\mathcal{U}_{(\mathfrak{X}, \mathbf{p}_{\bullet} \sqcup \mathbf{q}_{\diamond})}(V)$ is called the *(twisted) universal enveloping algebra of V w.r.t. the data $(\mathfrak{X}, \mathbf{p}_{\bullet} \sqcup \mathbf{q}_{\diamond})$.*

DEFINITION: The *(twisted) Zhu's algebra of V w.r.t. the data $(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)$* is

$$\mathcal{A}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V) := \mathcal{U}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)_0 / \mathbf{N}^1.$$

DEFINITION: The *(twisted) Zhu's algebra of V w.r.t. the data $(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)$* is

$$\mathcal{A}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V) := \mathcal{U}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_0 / \mathbf{N}^1.$$

LEMMA: Suppose \mathfrak{X} is smooth and \mathbf{q}_\diamond contains all the stacky points. Then

$$\mathcal{A}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V) \cong \bigotimes_{j \in \diamond} \mathcal{A}_{\mathbf{q}_j}(V) \cong \bigotimes_{j \in \diamond} \mathcal{A}_{g_j}(V).$$

EXAMPLE: $\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) \cong A_g(V) \otimes A_{g-1}(V).$

EXAMPLE: $\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) \cong A_g(V) \otimes A_{g-1}(V)$. Hence, the space of coinvariants is

$$(M^1)_{\mathcal{L}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) < 0} \otimes_{\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V)} (U^2 \otimes U^3)$$

EXAMPLE: $\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) \cong A_g(V) \otimes A_{g-1}(V)$. Hence, the space of coinvariants is

$$\begin{aligned} (M^1)_{\mathcal{L}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) < 0} \otimes_{\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V)} (U^2 \otimes U^3) \\ \cong A^{\text{op}}(M^1) \otimes_{A_g(V) \otimes A_{g-1}(V)} (U^2 \otimes U^3) \end{aligned}$$

EXAMPLE: $\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) \cong A_g(V) \otimes A_{g-1}(V)$. Hence, the space of coinvariants is

$$\begin{aligned} (M^1)_{\mathcal{L}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V) < 0} &\otimes_{\mathcal{A}_{([\mathbb{P}^1/\mu_T], (1) \sqcup (0, \infty))}(V)} (U^2 \otimes U^3) \\ &\cong A^{\text{op}}(M^1) \otimes_{A_g(V) \otimes A_{g-1}(V)} (U^2 \otimes U^3) \\ &= U^3 \otimes_{A_g(V)} A(M^1) \otimes_{A_g(V)} U^2. \end{aligned}$$

THEOREM (G., Liu, Zhu 24'): Suppose $M^\bullet \sqcup N^\diamond$ is an $(m+n)$ -tuple of (twisted) V -module *generated by their finite-dimensional bottom levels*. Suppose q_\diamond contains all the stacky points. Then, the space of coinvariants $(M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup q_\diamond)}(V)}$ is *finite-dimensional*.

THEOREM (G., Liu, Zhu 24'): Suppose $M^\bullet \sqcup N^\diamond$ is an $(m+n)$ -tuple of (twisted) V -module generated by their finite-dimensional bottom levels. Suppose \mathbf{q}_\diamond contains all the stacky points. Then, the space of coinvariants $(M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)}$ is finite-dimensional.

Proof. We have surjective map

$$(M^\bullet \otimes N^\diamond(0))_{\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)_{\leq 0}} \twoheadrightarrow (M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, \mathbf{p}_\bullet \sqcup \mathbf{q}_\diamond)}(V)}.$$

□

THEOREM (G., Liu, Zhu 24'): Suppose $M^\bullet \sqcup N^\diamond$ is an $(m+n)$ -tuple of (twisted) V -module generated by their finite-dimensional bottom levels. Suppose \mathfrak{q}_\diamond contains all the stacky points. Then, the space of coinvariants $(M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)}$ is finite-dimensional.

Proof. We have surjective map

$$(M^\bullet \otimes N^\diamond(0))_{\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)_{\leq 0}} \longrightarrow (M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)}.$$

We also have

$$L.H.S. \cong (M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)_{< 0}} \otimes_{\mathcal{A}_{(\mathfrak{X}, \mathfrak{p}_\bullet \sqcup \mathfrak{q}_\diamond)}(V)} N^\diamond(0).$$

□

Proof. We have surjective map

$$(M^\bullet \otimes N^\diamond(0))_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)_{\leq 0}} \longrightarrow (M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)}.$$

We also have

$$L.H.S. \cong (M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)_{< 0}} \otimes_{\mathcal{A}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)} N^\diamond(0).$$

Note that

$$(M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)_{< 0}} = (M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet)}(V)} = (M^\bullet)_{\mathcal{L}_{(P, \tilde{p}_\bullet)}(V)}.$$

□

Proof. We have surjective map

$$(M^\bullet \otimes N^\diamond(0))_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)_{\leq 0}} \longrightarrow (M^\bullet \otimes N^\diamond)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)}.$$

We also have

$$L.H.S. \cong (M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)_{< 0}} \otimes_{\mathcal{A}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)} N^\diamond(0).$$

Note that

$$(M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet \sqcup q_\diamond)}(V)_{< 0}} = (M^\bullet)_{\mathcal{L}_{(\mathfrak{X}, p_\bullet)}(V)} = (M^\bullet)_{\mathcal{L}_{(P, \tilde{p}_\bullet)}(V)}.$$

The coherence of $(M^\bullet)_{\mathcal{L}_{(P, \tilde{p}_\bullet)}(V)}$ has been proven by (Damiolini, Gibney, and Tarasca 21'–24'). □

Thank you!