

Transcendence of Periods

Xu Gao

2019

Outline

- 1 What is a period?
- 2 Transcendence of periods
- 3 Higher periods on Calabi-Yau varieties
- 4 Periods on a remarkable hypersurface

Motivation

- Integration of a meromorphic function depends on the choice of paths.

Motivation

- Integration of a meromorphic function depends on the choice of paths.
- Difference is given by integration of the form

$$\int_{\gamma} \omega,$$

where γ is a 1-cycle and ω is a closed 1-form.

Motivation

- Integration of a meromorphic function depends on the choice of paths.
- Difference is given by integration of the form

$$\int_{\gamma} \omega,$$

where γ is a 1-cycle and ω is a closed 1-form.

- Restrictions may need:

everything should be defined over $\overline{\mathbb{Q}}$.

Motivation

- Integration of a meromorphic function depends on the choice of paths.
- Difference is given by integration of the form

$$\int_{\gamma} \omega,$$

where γ is a 1-cycle and ω is a closed 1-form.

- Restrictions may need:

everything should be defined over $\overline{\mathbb{Q}}$.

- Generalization is worth to consider:

integration of a closed n -form along a n -cycle.

What is a period?

X a smooth algebraic variety (over $\overline{\mathbb{Q}}$)

What is a period?

- X a smooth algebraic variety (over $\overline{\mathbb{Q}}$)
- X^{an} the associated complex manifold of X

What is a period?

- X a smooth algebraic variety (over $\overline{\mathbb{Q}}$)
- X^{an} the associated complex manifold of X
- ω a closed algebraic n -form on X ,
which defines a holomorphic form on X^{an}

What is a period?

- X a smooth algebraic variety (over $\overline{\mathbb{Q}}$)
- X^{an} the associated complex manifold of X
- ω a closed algebraic n -form on X ,
which defines a holomorphic form on X^{an}
- γ a n -cycle on X^{an}

What is a period?

- X a smooth algebraic variety (over $\overline{\mathbb{Q}}$)
- X^{an} the associated complex manifold of X
- ω a closed algebraic n -form on X ,
which defines a holomorphic form on X^{an}
- γ a n -cycle on X^{an}

Definition (Kontsevich and Zagier, 2001)

A **n -period** of X is the integration

$$\int_{\gamma} \omega.$$

Examples

Here are some examples:

Examples

Here are some examples:

0 is a period

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.
- For example, $\mathbb{C} = (\mathbb{G}_a)^{\text{an}}$ is contractible.

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.
- For example, $\mathbb{C} = (\mathbb{G}_a)^{\text{an}}$ is contractible.
- Periods of \mathbb{G}_a are $\oint dz = 0$.

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.
- For example, $\mathbb{C} = (\mathbb{G}_a)^{\text{an}}$ is contractible.
- Periods of \mathbb{G}_a are $\oint dz = 0$.

$2\pi\sqrt{-1}$ is a period

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.
- For example, $\mathbb{C} = (\mathbb{G}_a)^{\text{an}}$ is contractible.
- Periods of \mathbb{G}_a are $\oint dz = 0$.

$2\pi\sqrt{-1}$ is a period

- $\frac{dz}{z}$ gives rise to a non-exact holomorphic form on $\mathbb{C}^\times = (\mathbb{G}_m)^{\text{an}}$.

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.
- For example, $\mathbb{C} = (\mathbb{G}_a)^{\text{an}}$ is contractible.
- Periods of \mathbb{G}_a are $\oint dz = 0$.

$2\pi\sqrt{-1}$ is a period

- $\frac{dz}{z}$ gives rise to a non-exact holomorphic form on $\mathbb{C}^\times = (\mathbb{G}_m)^{\text{an}}$.
- $H_1(\mathbb{C}^\times, \mathbb{Z})$ is generated by the unit circle.

Examples

Here are some examples:

0 is a period

- $H_1(M, \mathbb{Z}) = 0$ if M is contractible.
- For example, $\mathbb{C} = (\mathbb{G}_a)^{\text{an}}$ is contractible.
- Periods of \mathbb{G}_a are $\oint dz = 0$.

$2\pi\sqrt{-1}$ is a period

- $\frac{dz}{z}$ gives rise to a non-exact holomorphic form on $\mathbb{C}^\times = (\mathbb{G}_m)^{\text{an}}$.
- $H_1(\mathbb{C}^\times, \mathbb{Z})$ is generated by the unit circle.
- Periods of \mathbb{G}_m are multiples of $\oint \frac{dz}{z} = 2\pi\sqrt{-1}$.

Periods of an elliptic curve E

Periods of an elliptic curve E

- $H^0(E, \Omega^1)$ is spanned by $\omega = \frac{dx}{y}$.

Periods of an elliptic curve E

- $H^0(E, \Omega^1)$ is spanned by $\omega = \frac{dx}{y}$.
- $H_1(E^{\text{an}}, \mathbb{Z})$ has two generators α, β .

Periods of an elliptic curve E

- $H^0(E, \Omega^1)$ is spanned by $\omega = \frac{dx}{y}$.
- $H_1(E^{\text{an}}, \mathbb{Z})$ has two generators α, β .
- The $\overline{\mathbb{Q}}$ -vector space \mathcal{P} of periods of E is spanned by

$$\omega_1 = \int_{\alpha} \omega \quad \text{and} \quad \omega_2 = \int_{\beta} \omega.$$

Periods of an elliptic curve E

- $H^0(E, \Omega^1)$ is spanned by $\omega = \frac{dx}{y}$.
- $H_1(E^{\text{an}}, \mathbb{Z})$ has two generators α, β .
- The $\overline{\mathbb{Q}}$ -vector space \mathcal{P} of periods of E is spanned by

$$\omega_1 = \int_{\alpha} \omega \quad \text{and} \quad \omega_2 = \int_{\beta} \omega.$$

- The lattice $\Lambda = \mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2$ gives rise to the isomorphism

$$E^{\text{an}} \cong \mathbb{C}/\Lambda.$$

Outline

- 1 What is a period?
- 2 Transcendence of periods**
- 3 Higher periods on Calabi-Yau varieties
- 4 Periods on a remarkable hypersurface

Transcendence of periods

Observation

$2\pi\sqrt{-1}, \omega_1, \omega_2$ are unlikely to be algebraic.

Transcendence of periods

Observation

$2\pi\sqrt{-1}, \omega_1, \omega_2$ are unlikely to be algebraic.

So we have the following conjecture:

Conjecture

The 1-periods are either 0 or transcendental.

Transcendence of periods

Observation

$2\pi\sqrt{-1}, \omega_1, \omega_2$ are unlikely to be algebraic.

So we have the following conjecture:

Conjecture

The 1-periods are either 0 or transcendental.

More radically, one can conjecture the following:

Conjecture

The n -periods ($n \geq 1$) are either 0 or transcendental.

Known results

According to Waldschmidt (2006), the following results are known.

Known results

According to Waldschmidt (2006), the following results are known.

- (Lindemann, 1882) π is transcendental.

Known results

According to Waldschmidt (2006), the following results are known.

- (Lindemann, 1882) π is transcendental.
- (Schneider, 1935) ω_1, ω_2 are transcendental.

According to Waldschmidt (2006), the following results are known.

- (Lindemann, 1882) π is transcendental.
- (Schneider, 1935) ω_1, ω_2 are transcendental.
- (Schneider, 1937) $\tau = \omega_2/\omega_1$ is algebraic if and only if the elliptic curve E has complex multiplication.

According to Waldschmidt (2006), the following results are known.

- (Lindemann, 1882) π is transcendental.
- (Schneider, 1935) ω_1, ω_2 are transcendental.
- (Schneider, 1937) $\tau = \omega_2/\omega_1$ is algebraic if and only if the elliptic curve E has complex multiplication.
- (Schneider, 1941) On an abelian variety, not all periods are algebraic.

According to Waldschmidt (2006), the following results are known.

- (Lindemann, 1882) π is transcendental.
- (Schneider, 1935) ω_1, ω_2 are transcendental.
- (Schneider, 1937) $\tau = \omega_2/\omega_1$ is algebraic if and only if the elliptic curve E has complex multiplication.
- (Schneider, 1941) On an abelian variety, not all periods are algebraic.
- (Lang, 1966) The *Schneider-Lang Theorem*, which uniformizes proofs of above results.

Schneider-Lang Theorem

(Lang, 1966)

Schneider-Lang Theorem

(Lang, 1966)

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Suppose that

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Suppose that

- the ring $K[f_1, \dots, f_m]$ is stable under differentiation and

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Suppose that

- the ring $K[f_1, \dots, f_m]$ is stable under differentiation and
- has transcendence degree at least 2.

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Suppose that

- the ring $K[f_1, \dots, f_m]$ is stable under differentiation and
- has transcendence degree at least 2.

Then

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Suppose that

- the ring $K[f_1, \dots, f_m]$ is stable under differentiation and
- has transcendence degree at least 2.

Then

- there are only finitely many complex numbers,

Schneider-Lang Theorem

(Lang, 1966)

Let

- K be a number field and
- f_1, \dots, f_m be meromorphic functions of finite order.

Suppose that

- the ring $K[f_1, \dots, f_m]$ is stable under differentiation and
- has transcendence degree at least 2.

Then

- there are only finitely many complex numbers,
- at which f_1, \dots, f_m simultaneously assume values in K .

Known results

According to Waldschmidt (2006), the following results are known.

- (Lindemann, 1882) π is transcendental.
- (Schneider, 1935) ω_1, ω_2 are transcendental.
- (Schneider, 1937) $\tau = \omega_2/\omega_1$ is algebraic if and only if the elliptic curve E has complex multiplication.
- (Schneider, 1941) On an abelian variety, not all periods are algebraic.
- (Lang, 1966) The *Schneider-Lang Theorem*, which uniformizes proofs of above results.
- (Wüstholz, 1987, 1989) The *Analytic Subgroup Theorem*, which implies the transcendence and linearly independence of 1-periods of smooth projective varieties.

Analytic subgroup theorem

(Wüstholz, 1989)

Analytic subgroup theorem

(Wüstholz, 1989)

Analytic subgroup theorem

(Wüstholz, 1989)

Let

- G be an algebraic group and

Analytic subgroup theorem

(Wüstholz, 1989)

Let

- G be an algebraic group and
- H an analytic subgroup of G^{an} ,

Analytic subgroup theorem

(Wüstholz, 1989)

Let

- G be an algebraic group and
- H an analytic subgroup of G^{an} ,

both defined over $\overline{\mathbb{Q}}$.

Analytic subgroup theorem

(Wüstholz, 1989)

Let

- G be an algebraic group and
- H an analytic subgroup of G^{an} ,

both defined over $\overline{\mathbb{Q}}$.

- If H contains a non-trivial algebraic point,

Analytic subgroup theorem

(Wüstholz, 1989)

Let

- G be an algebraic group and
- H an analytic subgroup of G^{an} ,

both defined over $\overline{\mathbb{Q}}$.

- If H contains a non-trivial algebraic point,
- then it contains a non-trivial algebraic subgroup.

1-periods on smooth projective varieties

Let X be a smooth projective variety defined over $\overline{\mathbb{Q}}$.

1-periods on smooth projective varieties

Let X be a smooth projective variety defined over $\overline{\mathbb{Q}}$.

Transcendence (Wüstholz, 1987)

The 1-periods of X are either zero or transcendental.

1-periods on smooth projective varieties

Let X be a smooth projective variety defined over $\overline{\mathbb{Q}}$.

Transcendence (Wüstholz, 1987)

The 1-periods of X are either zero or transcendental.

Linearly Independence (Wüstholz, 1987)

Let \mathcal{V} be the $\overline{\mathbb{Q}}$ -vector space spanned by $1, \pi$ and all 1-periods of X . Then

$$\dim_{\overline{\mathbb{Q}}} \mathcal{V} = 2 + \sum_{i=1}^k \frac{2 (\dim A_i)^2}{[\mathrm{End}_0 A_i : \mathbb{Q}]},$$

where A_i are simple abelian subvarieties of the Albanese variety of X .

Outline

- 1 What is a period?
- 2 Transcendence of periods
- 3 Higher periods on Calabi-Yau varieties**
- 4 Periods on a remarkable hypersurface

Periods on Calabi-Yau varieties

Study higher periods on good varieties

Periods on Calabi-Yau varieties

Study higher periods on good varieties

- where the lower periods are ruled out.

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and
- $H^j(X, \mathcal{O}_X) = 0$ for all $0 < j < \dim X$.

Periods on Calabi-Yau varieties

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and
- $H^j(X, \mathcal{O}_X) = 0$ for all $0 < j < \dim X$.

Examples

Periods on Calabi-Yau varieties

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and
- $H^j(X, \mathcal{O}_X) = 0$ for all $0 < j < \dim X$.

Examples

- Elliptic curves.

Periods on Calabi-Yau varieties

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and
- $H^j(X, \mathcal{O}_X) = 0$ for all $0 < j < \dim X$.

Examples

- Elliptic curves.
- K3 surfaces.

Periods on Calabi-Yau varieties

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and
- $H^j(X, \mathcal{O}_X) = 0$ for all $0 < j < \dim X$.

Examples

- Elliptic curves.
- K3 surfaces.
- Smooth hypersurfaces of degree $n + 1$ in \mathbb{P}^n .

Periods on Calabi-Yau varieties

Definition

A **Calabi-Yau variety** X is a smooth proper variety such that

- the canonical sheaf ω_X is trivial and
- $H^j(X, \mathcal{O}_X) = 0$ for all $0 < j < \dim X$.

Examples

- Elliptic curves.
- K3 surfaces.
- Smooth hypersurfaces of degree $n + 1$ in \mathbb{P}^n .
-

Periods on Calabi-Yau varieties

Why Calabi-Yau varieties?

Why Calabi-Yau varieties?

- The only interesting periods are given by a unique (up to scale) holomorphic form.

Periods on Calabi-Yau varieties

Why Calabi-Yau varieties?

- The only interesting periods are given by a unique (up to scale) holomorphic form.
- Hence $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$ implies $H^{n,0} = H_{\overline{\mathbb{Q}}}^{n,0} \otimes_{\overline{\mathbb{Q}}} \mathbb{C}$.

Periods on Calabi-Yau varieties

Why Calabi-Yau varieties?

- The only interesting periods are given by a unique (up to scale) holomorphic form.
- Hence $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$ implies $H^{n,0} = H_{\overline{\mathbb{Q}}}^{n,0} \otimes_{\overline{\mathbb{Q}}} \mathbb{C}$.
- This is an unusual property since in many case, or with a little bit more assumptions, it would imply that the Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Can a Hodge decomposition being defined over $\overline{\mathbb{Q}}$?

Proposition

If a rational Hodge structure has CM, in the sense that its Mumford-Tate group is abelian, then its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Can a Hodge decomposition being defined over $\overline{\mathbb{Q}}$?

Proposition

If a rational Hodge structure has CM, in the sense that its Mumford-Tate group is abelian, then its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Proposition

For a smooth projective variety, the followings are the same:

Can a Hodge decomposition being defined over $\overline{\mathbb{Q}}$?

Proposition

If a rational Hodge structure has CM, in the sense that its Mumford-Tate group is abelian, then its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Proposition

For a smooth projective variety, the followings are the same:

- 1 $\dim_{\overline{\mathbb{Q}}} \mathcal{P}$ assumes its minimum;

Can a Hodge decomposition being defined over $\overline{\mathbb{Q}}$?

Proposition

If a rational Hodge structure has CM, in the sense that its Mumford-Tate group is abelian, then its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Proposition

For a smooth projective variety, the followings are the same:

- 1 $\dim_{\overline{\mathbb{Q}}} \mathcal{P}$ assumes its minimum;
- 2 the Hodge structure $H^1(X^{\text{an}}, \mathbb{Q})$ has CM, and

Can a Hodge decomposition being defined over $\overline{\mathbb{Q}}$?

Proposition

If a rational Hodge structure has CM, in the sense that its Mumford-Tate group is abelian, then its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Proposition

For a smooth projective variety, the followings are the same:

- 1 $\dim_{\overline{\mathbb{Q}}} \mathcal{P}$ assumes its minimum;
- 2 the Hodge structure $H^1(X^{\text{an}}, \mathbb{Q})$ has CM, and
- 3 its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Periods and CM

In general, there is a conjecture:

Conjecture

A polarizable rational Hodge structure has CM if and only if its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Periods and CM

In general, there is a conjecture:

Conjecture

A polarizable rational Hodge structure has CM if and only if its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

In the case of Calabi-Yau n -varieties we consider

Conjecture

$H^n(X^{\text{an}}, \mathbb{Q})$ has CM if and only if its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Periods and CM

In general, there is a conjecture:

Conjecture

A polarizable rational Hodge structure has CM if and only if its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

In the case of Calabi-Yau n -varieties we consider

Conjecture

$H^n(X^{\text{an}}, \mathbb{Q})$ has CM if and only if its Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Stronger Conjecture

$\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$ implies $H^n(X^{\text{an}}, \mathbb{Q})$ has CM.

A recent result on this direction is the following:

A recent result on this direction is the following:

Tretkoff (2015, 2017)

Let X be a K3 surface defined over $\overline{\mathbb{Q}}$. If $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$, then the Hodge structure on $H^2(X^{\text{an}}, \mathbb{Q})_{\text{prim}}$ has CM.

A recent result on this direction is the following:

Tretkoff (2015, 2017)

Let X be a K3 surface defined over $\overline{\mathbb{Q}}$. If $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$, then the Hodge structure on $H^2(X^{\text{an}}, \mathbb{Q})_{\text{prim}}$ has CM.

This result is proven through the *Kuga-Satake correspondence*.

Proposition-Definition

Proposition-Definition

- Starting with a polarized weight 2 rational Hodge structure V such that $h^{2,0} = 1$,

Proposition-Definition

- Starting with a polarized weight 2 rational Hodge structure V such that $h^{2,0} = 1$,
- there is a polarized weight 1 rational Hodge structure on $\mathrm{Cl}^+(V)$,

Proposition-Definition

- Starting with a polarized weight 2 rational Hodge structure V such that $h^{2,0} = 1$,
- there is a polarized weight 1 rational Hodge structure on $\mathrm{Cl}^+(V)$,
- and an inclusion of Hodge structures

$$V \hookrightarrow \mathrm{Cl}^+(V) \otimes \mathrm{Cl}^+(V).$$

Proposition-Definition

- Starting with a polarized weight 2 rational Hodge structure V such that $h^{2,0} = 1$,
- there is a polarized weight 1 rational Hodge structure on $Cl^+(V)$,
- and an inclusion of Hodge structures

$$V \hookrightarrow Cl^+(V) \otimes Cl^+(V).$$

- The corresponding abelian variety $KS(V)$ of the Hodge structure $Cl^+(V)$ is called the **Kuga-Satake variety** of V .

Kuga-Satake correspondence

Proposition-Definition

- Starting with a polarized weight 2 rational Hodge structure V such that $h^{2,0} = 1$,
- there is a polarized weight 1 rational Hodge structure on $\mathrm{Cl}^+(V)$,
- and an inclusion of Hodge structures

$$V \hookrightarrow \mathrm{Cl}^+(V) \otimes \mathrm{Cl}^+(V).$$

- The corresponding abelian variety $KS(V)$ of the Hodge structure $\mathrm{Cl}^+(V)$ is called the **Kuga-Satake variety** of V .

However, similar construction is not available for higher Hodge structures.

Outline

- 1 What is a period?
- 2 Transcendence of periods
- 3 Higher periods on Calabi-Yau varieties
- 4 Periods on a remarkable hypersurface

The remarkable hypersurface

Let X be the hypersurface in \mathbb{P}^4 cut out by the polynomial

$$x_0^5 + x_1^5 + x_2^5 + x_3^5 + x_4^5 - 5x_0x_1x_2x_3x_4.$$

The remarkable hypersurface

Let X be the hypersurface in \mathbb{P}^4 cut out by the polynomial

$$x_0^5 + x_1^5 + x_2^5 + x_3^5 + x_4^5 - 5x_0x_1x_2x_3x_4.$$

This is a remarkable hypersurface,

- which is a member of the Dwork family (Dwork, 1963),

The remarkable hypersurface

Let X be the hypersurface in \mathbb{P}^4 cut out by the polynomial

$$x_0^5 + x_1^5 + x_2^5 + x_3^5 + x_4^5 - 5x_0x_1x_2x_3x_4.$$

This is a remarkable hypersurface,

- which is a member of the Dwork family (Dwork, 1963),
- which is a special determinantal hypersurface of the Horrocks–Mumford bundle (Horrocks and Mumford, 1973)

The remarkable hypersurface

Let X be the hypersurface in \mathbb{P}^4 cut out by the polynomial

$$x_0^5 + x_1^5 + x_2^5 + x_3^5 + x_4^5 - 5x_0x_1x_2x_3x_4.$$

This is a remarkable hypersurface,

- which is a member of the Dwork family (Dwork, 1963),
- which is a special determinantal hypersurface of the Horrocks–Mumford bundle (Horrocks and Mumford, 1973)
- which has 125 nodes and

The remarkable hypersurface

Let X be the hypersurface in \mathbb{P}^4 cut out by the polynomial

$$x_0^5 + x_1^5 + x_2^5 + x_3^5 + x_4^5 - 5x_0x_1x_2x_3x_4.$$

This is a remarkable hypersurface,

- which is a member of the Dwork family (Dwork, 1963),
- which is a special determinantal hypersurface of the Horrocks–Mumford bundle (Horrocks and Mumford, 1973)
- which has 125 nodes and
- whose geometry was studied by Schoen (1985, 1986).

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.
- General fibers of $X' \rightarrow \mathbb{P}^1$ are non-CM abelian surfaces.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.
- General fibers of $X' \rightarrow \mathbb{P}^1$ are non-CM abelian surfaces.
- The periods of X' and of its general fibers should be closely related. This leads to an impression that X' is unlikely to have CM.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.
- General fibers of $X' \rightarrow \mathbb{P}^1$ are non-CM abelian surfaces.
- The periods of X' and of its general fibers should be closely related. This leads to an impression that X' is unlikely to have CM.
- On the other hand, $h^{1,2} = h^{2,1} = 0$, hence if $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$, then the Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.
- General fibers of $X' \rightarrow \mathbb{P}^1$ are non-CM abelian surfaces.
- The periods of X' and of its general fibers should be closely related. This leads to an impression that X' is unlikely to have CM.
- On the other hand, $h^{1,2} = h^{2,1} = 0$, hence if $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$, then the Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.
- General fibers of $X' \rightarrow \mathbb{P}^1$ are non-CM abelian surfaces.
- The periods of X' and of its general fibers should be closely related. This leads to an impression that X' is unlikely to have CM.
- On the other hand, $h^{1,2} = h^{2,1} = 0$, hence if $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$, then the Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Hence it is worth to try the following conjecture.

Desingularization

- There is a small desingularization of $X' \longrightarrow X$.
- X' is a Calabi-Yau 3-variety whose periods will be studied.
- General fibers of $X' \rightarrow \mathbb{P}^1$ are non-CM abelian surfaces.
- The periods of X' and of its general fibers should be closely related. This leads to an impression that X' is unlikely to have CM.
- On the other hand, $h^{1,2} = h^{2,1} = 0$, hence if $\dim_{\overline{\mathbb{Q}}} \mathcal{P} = 1$, then the Hodge decomposition is defined over $\overline{\mathbb{Q}}$.

Hence it is worth to try the following conjecture.

Conjecture

At least one of the periods of X' is transcendental.

References I

Bernard Dwork. A deformation theory for the zeta function of a hypersurface. In *Proc. Internat. Congr. Mathematicians (Stockholm, 1962)*, pages 247–259. Inst. Mittag-Leffler, Djursholm, 1963.

Geoffrey. Horrocks and David Mumford. A rank 2 vector bundle on \mathbf{P}^4 with 15,000 symmetries. *Topology*, 12:63–81, 1973.

Maxim Kontsevich and Don Zagier. Periods. In *Mathematics unlimited—2001 and beyond*, pages 771–808. Springer, Berlin, 2001.

Serge Lang. *Introduction to transcendental numbers*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1966.

Ferdinand von Lindemann. Ueber die Zahl π . *Math. Ann.*, 20(2):213–225, 1882.

Theodor Schneider. Transzendenzuntersuchungen periodischer Funktionen. *J. Reine Angew. Math.*, 172:65–74, 1935.

References II

- Theodor Schneider. Arithmetische Untersuchungen elliptischer Integrale. *Math. Ann.*, 113(1):1–13, 1937.
- Theodor Schneider. Zur Theorie der Abelschen Funktionen und Integrale. *J. Reine Angew. Math.*, 183:110–128, 1941.
- Chad Schoen. Algebraic cycles on certain desingularized nodal hypersurfaces. *Math. Ann.*, 270(1):17–27, 1985.
- Chad Schoen. On the geometry of a special determinantal hypersurface associated to the Mumford-Horrocks vector bundle. *J. Reine Angew. Math.*, 364:85–111, 1986.
- Paula Tretkoff. K3 surfaces with algebraic period ratios have complex multiplication. *Int. J. Number Theory*, 11(5):1709–1724, 2015.
- Paula Tretkoff. *Periods and special functions in transcendence*. Advanced Textbooks in Mathematics. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2017.

References III

- Michel Waldschmidt. Transcendence of periods: the state of the art. *Pure Appl. Math. Q.*, 2(2):435–463, 2006.
- Gisbert Wüstholz. Algebraic groups, Hodge theory, and transcendence. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, pages 476–483. Amer. Math. Soc., Providence, RI, 1987.
- Gisbert Wüstholz. Algebraische Punkte auf analytischen Untergruppen algebraischer Gruppen. *Ann. of Math. (2)*, 129(3):501–517, 1989.

Thanks!