

p -ADIC REPRESENTATIONS AND SIMPLICIAL DISTANCE IN BRUHAT-TITS BUILDINGS

Xu Gao

March 16, 2023

University of California, Santa Cruz

TABLE OF CONTENTS

p -adic representations and stable lattices

What are Bruhat-Tits buildings?

What can we say about fixed-point sets?

Simplicial distance

Simplicial balls

Simplicial volume

p -ADIC REPRESENTATIONS AND STABLE LATTICES

p -adic representation


$$\rho: G \longrightarrow \mathrm{GL}_K(V),$$

- V is a finite-dimensional vector space over
- K , a local field with residue characteristic p .

p -adic representation

$$\rho: G \longrightarrow \mathrm{GL}_K(V),$$


- V is a finite-dimensional vector space over
- K , a local field with residue characteristic p .



p -adic representation

$$\rho: G \longrightarrow \mathrm{GL}_K(V),$$

- V is a finite-dimensional vector space over
- K , a local field with residue characteristic p .



NOTATIONS:

- K = a non-Archimedean local field;
- val = the valuation on K (assuming $\text{val}(K^\times) = \mathbb{Z}$);
- K° = the ring of integers $\{x \in K \mid \text{val}(x) \geq 0\}$;
- ϖ = a uniformizer, namely ϖK° is the maximal ideal of K° ;
- κ = the residue field $K^\circ / \varpi K^\circ$;
- q = the cardinality of κ .

- ***lattice*** = f. g. K° -submodule L of V spanning the entire V .

- ***lattice*** = f. g. K° -submodule L of V spanning the entire V .
- ***stable lattice*** = lattice stable under the action of $\rho(G)$.

- **lattice** = f. g. K° -submodule L of V spanning the entire V .
- **stable lattice** = lattice stable under the action of $\rho(G)$.
- A stable lattice $L \leadsto$ a modular representation on $L \otimes_{K^\circ} K$.
 - $L \otimes_{K^\circ} K$ depends on the choice of L ,
 - its semisimplification doesn't.

The later is called the **mod ϖ reduction** V_K of V .

Question

When does such a stable lattice exist? What can we say about the set of stable lattices?

Question

When does such a stable lattice exist? What can we say about the set of stable lattices?

1. has a stable lattice \iff ***precompact***

Question

When does such a stable lattice exist? What can we say about the set of stable lattices?

1. has a stable lattice \iff **precompact**
2. “stable” maintained through **homothety**

1. has a stable lattice \iff **precompact**
2. “stable” maintained through **homothety**

$$S(\rho)^0 := \{\text{stable lattices of } \rho\} / \text{homothety}$$

$$h(\rho) := |S(\rho)^0| \quad \text{class number}$$

$$S(\rho)^0 := \{\text{stable lattices of } \rho\} / \text{homothety}$$

$$h(\rho) := |S(\rho)^0| \quad \textbf{class number}$$

Compare with:

$$Cl(F) := \{\text{fractional ideas in } F\} / \text{homothety}.$$

- (Iwasawa, 1973) $Cl(F_n)^{(p)}$ a tower of cyclic F_n/F

- (Iwasawa, 1973) $Cl(F_n)^{(p)}$ a tower of cyclic F_n/F
- $h(\rho \otimes_K E_n)$ a tower of totally ramified E_n/K

- (Iwasawa, 1973) $Cl(F_n)^{(p)}$ a tower of cyclic F_n/F
- $h(\rho \otimes_K E_n)$ a tower of totally ramified E_n/K
- But: no natural composition law on $S(\rho)^0$

- (Iwasawa, 1973) $Cl(F_n)^{(p)}$ a tower of cyclic F_n/F
- $h(\rho \otimes_K E_n)$ a tower of totally ramified E_n/K
- But: no natural composition law on $S(\rho)^0$
- ~~algebra of Galois modules~~ geometry of Bruhat-Tits buildings

- $\mathcal{X}(V)$ = (reduced) ***Bruhat-Tits building*** of $GL_K(V)$

- $\mathcal{X}(V)$ = (reduced) ***Bruhat-Tits building*** of $GL_K(V)$
- lattices in $V \longleftrightarrow$ vertices (0-simplicies) $\mathcal{X}(V)$

- $\mathcal{X}(V)$ = (reduced) ***Bruhat-Tits building*** of $GL_K(V)$
- lattices in $V \longleftrightarrow$ vertices (0-simplicies) $\mathcal{X}(V)$
- $S(\rho)$ = fixed-point set of $\rho(G)$ convex + simplicial

- $\mathcal{X}(V)$ = (reduced) ***Bruhat-Tits building*** of $GL_K(V)$
- lattices in $V \longleftrightarrow$ vertices (0-simplicies) $\mathcal{X}(V)$
- $S(\rho)$ = fixed-point set of $\rho(G)$ convex + simplicial
- $S(\rho)^0 \longleftrightarrow$ set of vertices in $S(\rho)$

Theorem (Junecue, 2021)

(i) $S(\rho)$ is compact $\iff h(\rho)$ is finite $\iff \rho$ is irreducible.

Theorem (Junecue, 2021)

- (i) $S(\rho)$ is compact $\iff h(\rho)$ is finite $\iff \rho$ is irreducible.
- (ii) $S(\rho)$ has dimension $r(\rho) - 1$ where $r(\rho)$ is **reduction rank** (= the number of irreducible components in the reduction V_κ).

Theorem (Junecue, 2021)

- (i) $S(\rho)$ is compact $\iff h(\rho)$ is finite $\iff \rho$ is irreducible.
- (ii) $S(\rho)$ has dimension $r(\rho) - 1$ where $r(\rho)$ is **reduction rank** (= the number of irreducible components in the reduction V_κ).
- (iii) $h(\rho \otimes_K E)$ is a polynomial of $[E : K]$ for any totally ramified extension E/K $\iff \rho$ has **regular reduction**.

What if replace $GL_K(V)$ by other groups?

Example

V has a non-degenerate symplectic (resp. orthogonal) form and the action of G respects it. Then ρ lands in $Sp(V)$ (resp. $O(V)$).

What if replace $GL_K(V)$ by other groups?

Example

V has a non-degenerate symplectic (resp. orthogonal) form and the action of G respects it. Then ρ lands in $Sp(V)$ (resp. $O(V)$).

Consider ***stable (almost) self-dual lattices***

In general, we can consider group homomorphisms to the groups of K -rational points of **reductive groups**:

$$\rho: G \longrightarrow \mathcal{G}(K).$$

Expect: the study of fix-point set of $\rho(G)$ in the Bruhat-Tits building of \mathcal{G} can aid the research of ρ .

WHAT ARE BRUHAT-TITS BUILDINGS?

WHAT ARE BRUHAT-TITS BUILDINGS?

- \mathcal{G} reductive group \leadsto $\mathcal{B}(\mathcal{G})$ ***Bruhat-Tits building***
metric space + polysimplicial complex


WHAT ARE BRUHAT-TITS BUILDINGS?

- \mathcal{G} reductive group $\leadsto \mathcal{B}(\mathcal{G})$ **Bruhat-Tits building**
metric space + polysimplicial complex
- \mathcal{T} maximal split torus of $\mathcal{G} \leadsto \mathcal{A}(\mathcal{T})$ **affine apartment**
depends on **root datum** + $\text{val}(\cdot)$


WHAT ARE BRUHAT-TITS BUILDINGS?

- \mathcal{G} reductive group $\leadsto \mathcal{B}(\mathcal{G})$ **Bruhat-Tits building**
metric space + polysimplicial complex
- \mathcal{T} maximal split torus of $\mathcal{G} \leadsto \mathcal{A}(\mathcal{T})$ **affine apartment**
depends on **root datum** + $\text{val}(\cdot)$
- building = gluing apartments cleverly


EXAMPLES OF APARTMENTS




EXAMPLES OF APARTMENTS




EXAMPLES OF APARTMENTS



EXAMPLES OF APARTMENTS




EXAMPLES OF APARTMENTS




A_2

EXAMPLES OF APARTMENTS




$A_1 \times A_1$



C_2

EXAMPLES OF BUILDINGS: BRUHAT-TITS TREES



The Bruhat-Tits tree
(building of $GL_K(2)$)
with an apartment
specified by red color.

WHAT CAN WE SAY ABOUT FIXED-POINT SETS?

WHAT CAN WE SAY ABOUT FIXED-POINT SETS?

- We require ρ ***type-preserving*** isometries
(***precompact*** representations)

WHAT CAN WE SAY ABOUT FIXED-POINT SETS?

- We require ρ **type-preserving** isometries
(**precompact** representations)
- $S(\rho)$ is convex + simplicial

WHAT CAN WE SAY ABOUT FIXED-POINT SETS?

- We require ρ **type-preserving** isometries
(**precompact** representations)
- $S(\rho)$ is convex + simplicial
- $S(\rho)$ is compact $\iff h(\rho)$ is finite

WHAT CAN WE SAY ABOUT FIXED-POINT SETS?

- We require ρ **type-preserving** isometries (**precompact** representations)
- $S(\rho)$ is convex + simplicial
- $S(\rho)$ is compact $\iff h(\rho)$ is finite
- (Prasad and Yu, 2002) if $\rho(G)$ contains no **p -torsions**, then $S(\rho)$ is a Bruhat-Tits building.

We may ask the following questions

Question

- *How does $S(\rho)$ change along totally ramified extensions?*

We may ask the following questions


Question

- *How does $S(\rho)$ change along totally ramified extensions?*
- *Can we have a concrete description of $S(\rho)$?*


We may ask the following questions

Question


- *How does $S(\rho)$ change along totally ramified extensions?*
- *Can we have a concrete description of $S(\rho)$?*
- *Given an interesting convex simplicial subset of the building, can we realize it as $S(\rho)$ for some ρ ?*




The behavior of a “regular”
fixed-point set.




The behavior of a “irregular”
fixed-point set.




The behavior of a “regular”
fixed-point set.



The behavior of a “irregular”
fixed-point set.



The behavior of a “regular”
fixed-point set.



The behavior of a “irregular”
fixed-point set.


SIMPLICIAL DISTANCE

- Simplicial structure \leadsto vertices \leadsto ***incident geometry***


- Simplicial structure \leadsto vertices \leadsto ***incident geometry***
- (Garrett, 1997) Interpret Bruhat-Tits buildings as the incident geometry of ***lattices***
(where vertices = homothety classes of certain lattices)

- Simplicial structure \leadsto vertices \leadsto ***incident geometry***
- (Garrett, 1997) Interpret Bruhat-Tits buildings as the incident geometry of ***lattices***
(where vertices = homothety classes of certain lattices)
- \leadsto ***simplicial distance*** (=distance by incidents)

- Simplicial structure \leadsto vertices \leadsto ***incident geometry***
- (Garrett, 1997) Interpret Bruhat-Tits buildings as the incident geometry of ***lattices***
(where vertices = homothety classes of certain lattices)
- \leadsto ***simplicial distance*** (=distance by incidents)
- \leadsto ***simplicial balls, simplicial volume, simplicial curvature*** etc.



A path of length 3.



Simplicial ball of radius 3.

However, the incident relations in (Garrett, 1997) are complicated.
→ difficult to study simplicial distance

Question

However, the incident relations in (Garrett, 1997) are complicated.

→ difficult to study simplicial distance

Question

- *Can we have a more concrete, computable description of simplicial distance? of simplicial balls?*

However, the incident relations in (Garrett, 1997) are complicated.

↪ difficult to study simplicial distance

Question

- *Can we have a more concrete, computable description of simplicial distance? of simplicial balls?*
- *Are view simplicial balls fixed-point sets?*

However, the incident relations in (Garrett, 1997) are complicated.

→ difficult to study simplicial distance

Question

- *Can we have a more concrete, computable description of simplicial distance? of simplicial balls?*
- *Are view simplicial balls fixed-point sets?*
- *How does simplicial balls grow?*

Φ the root system: *irreducible and classical* (i.e. A_n , B_n , C_n , or D_n)


Theorem (G., 2022)

$d(x, y) - 1 =$ *maximum of numbers of walls between them;*
in particular, the simplicial distance from the origin o to x is


$$d(x, o) = \lceil a_0(x - o) \rceil,$$

where a_0 = highest root rel. Weyl chamber covering x .

SIMPLICIAL BALLS IN CLASSICAL TYPE BUILDINGS




The two blues are separated by 3 walls and have distance 4.




The blue is separated by 2 walls from the red and has distance 3.

SIMPLICIAL BALLS IN CLASSICAL TYPE BUILDINGS




The two blues are separated by 3 walls and have distance 4.




The blue has value 3 under a_0 and is of distance 3 from the red.

SIMPLICIAL BALLS IN CLASSICAL TYPE BUILDINGS



A path of length 4.



A path of length 3.

Corollary



If $\Phi = A_n$ or C_n , then $d(x, y) \leq r \iff \exists L \in x, L' \in y$ such that

$$L \supset L' \supset \varpi^r L.$$

When $\Phi = B_n$ or D_n , the simplicial distance could be shorter due to the oriflamme construction.

SIMPLICIAL BALLS

SIMPLICIAL BALLS



Simplicial balls of radius 2.

$B(x, r)$ = simplicial ball of radius r at x

Theorem (G., 2022)

$B(x, r)$ = *fixed-point set of the Moy-Prasad subgroup $P_{x,r}$.*

$B(x, r)$ = simplicial ball of radius r at x

Theorem (G., 2022)

$B(x, r)$ = *fixed-point set of the Moy-Prasad subgroup $P_{x,r}$.*

- (Moy and Prasad, 1996) **Moy-Prasad subgroups** $(P_{x,r})_{r \geq 0}$
generalizing principal congruence subgroups

$B(x, r)$ = simplicial ball of radius r at x

Theorem (G., 2022)

$B(x, r)$ = *fixed-point set of the Moy-Prasad subgroup $P_{x,r}$.*

- (Moy and Prasad, 1996) **Moy-Prasad subgroups** $(P_{x,r})_{r \geq 0}$
generalizing principal congruence subgroups
- (Yu, 2015) the machinery of **concave functions**

Bruhat-Tits theory

- ***parahoric subgroup*** $P_x \approx$ stabilizer of x

Bruhat-Tits theory

- **parahoric subgroup** $P_x \approx$ stabilizer of x
- \leadsto (connected) smooth model \mathfrak{G}_x of \mathcal{G} such that $\mathfrak{G}_x(K^\circ) = P_x$

Bruhat-Tits theory

- **parahoric subgroup** $P_x \approx$ stabilizer of x
- \leadsto (connected) smooth model \mathfrak{G}_x of \mathcal{G} such that $\mathfrak{G}_x(K^\circ) = P_x$

(Yu, 2015)

- **concave function** f = function on $\tilde{\Phi} := \Phi \cup \{0\}$ s.t.

$$\forall a, b \in \tilde{\Phi}, a + b \in \tilde{\Phi} \implies f(a) + f(b) \geq f(a + b).$$

Bruhat-Tits theory

- **parahoric subgroup** $P_x \approx$ stabilizer of x
- \leadsto (connected) smooth model \mathfrak{G}_x of \mathcal{G} such that $\mathfrak{G}_x(K^\circ) = P_x$

(Yu, 2015)

- **concave function** f = function on $\tilde{\Phi} := \Phi \cup \{0\}$ s.t.

$$\forall a, b \in \tilde{\Phi}, a + b \in \tilde{\Phi} \implies f(a) + f(b) \geq f(a + b).$$


- $\leadsto \mathfrak{G}_f$ s.t. $P_f = \mathfrak{G}_f(K^\circ)$ is bounded

- Any point $x \in \mathcal{B}(\mathcal{G})$ defines a concave function $f_x: a \mapsto -a(x)$, and we have $P_{f_x} = P_x$.

- Any point $x \in \mathcal{B}(\mathcal{G})$ defines a concave function $f_x: a \mapsto -a(x)$, and we have $P_{f_x} = P_x$.
- If $r \geq 0$, the shifting $f + r$ is again a concave function.

- Any point $x \in \mathcal{B}(\mathcal{G})$ defines a concave function $f_x: a \mapsto -a(x)$, and we have $P_{f_x} = P_x$.
- If $r \geq 0$, the shifting $f + r$ is again a concave function.
- In general, $P_f \subseteq P_{f'} \iff f \geq f'$.

- Any point $x \in \mathcal{B}(\mathcal{G})$ defines a concave function $f_x: a \mapsto -a(x)$, and we have $P_{f_x} = P_x$.
- If $r \geq 0$, the shifting $f + r$ is again a concave function.
- In general, $P_f \subseteq P_{f'} \iff f \geq f'$.
- In particular, $(P_{f_x+r})_{r \geq 0}$ are the *Moy-Prasad subgroups*.



$$\dots \subset P_{\bullet, 2} \subset P_{\bullet} \subset P_{\bullet} \subset P_{\bullet} \subset P_{\bullet}$$

SIMPLICIAL VOLUME


$SV(r) := |B(o, r)|$ (**simplicial volume**)

Theorem (G., 2022)

$$SV(r) = \sum_{I \subset \Delta} \frac{\mathcal{P}_{\Phi; I}(q)}{q^{\deg(\mathcal{P}_{\Phi; I})}} \sum_{x \in B(r, C, I)} \prod_{a(x) > 0} q^{\lceil a(x) \rceil}$$

- $\mathcal{P}_{\Phi; I}$ = **Poincaré polynomial**
- C = *Weyl chamber*
- $B(r, C, I) = B(o, r) \cap \text{inn}(\bigcap_{a \in I} \ker(a))$

THE INDEX SET $B(r, C, l)$



$$B(2, \text{pink}, a_1) = \{\bullet, \bullet\},$$

$$B(2, \text{pink}, \Delta) = \{\bullet\},$$

$$B(2, \text{pink}, a_2) = \{\bullet, \bullet\},$$

$$B(2, \text{pink}, \emptyset) = \{\bullet\}.$$

$$SV(r) = \sum_{I \subset \Delta} \frac{\mathcal{P}_{\Phi;I}(q)}{q^{\deg(\mathcal{P}_{\Phi;I})}} \sum_{x \in B(r,C,I)} \prod_{a(x) > 0} q^{\lceil a(x) \rceil},$$

$$SV(r) = \sum_{I \subset \Delta} \frac{\mathcal{P}_{\Phi;I}(q)}{q^{\deg(\mathcal{P}_{\Phi;I})}} \sum_{x \in B(r,C,I)} \prod_{a(x) > 0} q^{\lceil a(x) \rceil},$$

- We have a concrete description of vertices.
(However, they DO NOT form a lattice.)

$$SV(r) = \sum_{I \subset \Delta} \frac{\mathcal{P}_{\Phi;I}(q)}{q^{\deg(\mathcal{P}_{\Phi;I})}} \sum_{x \in B(r,C,I)} \prod_{a(x) > 0} q^{\lceil a(x) \rceil},$$

- We have a concrete description of vertices.
(However, they DO NOT form a lattice.)
- With such a description, we can describe the index set $B(r, C, I)$ in terms of linear inequalities.

If \mathcal{B} is of type A_n and $I = \Delta \setminus \{\ell_1, \dots, \ell_t\}$:

$$B(r, C, I) = \{0 + c_1\omega_{\ell_1} + \dots + c_t\omega_{\ell_t} \mid c_i \in \mathbb{Z}_{>0}, c_1 + \dots + c_t \leq r\},$$

where ω_i are the fundamental coweights.

If \mathcal{B} is of type A_n and $I = \Delta \setminus \{\ell_1, \dots, \ell_t\}$:

$$B(r, C, I) = \{o + c_1\omega_{\ell_1} + \dots + c_t\omega_{\ell_t} \mid c_i \in \mathbb{Z}_{>0}, c_1 + \dots + c_t \leq r\},$$

where ω_i are the fundamental coweights.

If \mathcal{B} is of type C_n and $I = \Delta \setminus \{\ell_1, \dots, \ell_t\}$:

$$B(r, C, I) = \{o + c_1\omega'_{\ell_1} + \dots + c_t\omega'_{\ell_t} \mid c_i \in \mathbb{Z}_{>0}, c_1 + \dots + c_t \leq r\},$$

where $\omega'_i = h_i^{-1}\omega_i$ with $a_0 = \sum_i h_i a_i$.

If \mathcal{B} is of type A_n and $I = \Delta \setminus \{\ell_1, \dots, \ell_t\}$:

$$B(r, C, I) = \{o + c_1\omega_{\ell_1} + \dots + c_t\omega_{\ell_t} \mid c_i \in \mathbb{Z}_{>0}, c_1 + \dots + c_t \leq r\},$$


where ω_i are the fundamental coweights.

If \mathcal{B} is of type C_n and $I = \Delta \setminus \{\ell_1, \dots, \ell_t\}$:


$$B(r, C, I) = \{o + c_1\omega'_{\ell_1} + \dots + c_t\omega'_{\ell_t} \mid c_i \in \mathbb{Z}_{>0}, c_1 + \dots + c_t \leq r\},$$

where $\omega'_i = h_i^{-1}\omega_i$ with $a_0 = \sum_i h_i a_i$.

If \mathcal{B} is of type B_n or D_n , then the description is complicated.



Vertices in B_n building



Vertices in D_n building

Theorem (G., 2022)

We have the following asymptotic dominant relation as $r \rightarrow \infty$:

$$SV(r) \asymp r^{\epsilon(n)} q^{\pi(n)r},$$

where $\epsilon(n)$ and $\pi(n)$ are in the following table.

Theorem (G., 2022)

We have the following asymptotic dominant relation as $r \rightarrow \infty$:

$$SV(r) \asymp r^{\epsilon(n)} q^{\pi(n)r},$$

where $\epsilon(n)$ and $\pi(n)$ are in the following table.

X_n	$\epsilon(n)$	$\pi(n)$
A_n (n is odd)	0	$(\frac{n+1}{2})^2$
A_n (n is even)	1	$\frac{n}{2}(\frac{n}{2} + 1)$
B_n ($n = 3$)	0	5
B_n ($n \geq 4$)	0	$\frac{n^2}{2}$
C_n ($n \geq 2$)	0	$\frac{n(n+1)}{2}$
D_n ($n = 4$)	2	6
D_n ($n \geq 5$)	1	$\frac{n(n-1)}{2}$

Theorem (G., 2022)

Leading coefficients have the following rationality properties:

1. *If \mathcal{B} is of type A_n , C_n , B_3 , or D_4 , then we have*

$$SV(r) \sim \mathbf{c}(n) \cdot r^{\epsilon(n)} q^{\pi(n)r},$$

where $\mathbf{c}(n)$ is a positive number that is a rational function of q .

Theorem (G., 2022)

Leading coefficients have the following rationality properties:

1. *If \mathcal{B} is of type A_n , C_n , B_3 , or D_4 , then we have*

$$SV(r) \sim \mathbf{c}(n) \cdot r^{\epsilon(n)} q^{\pi(n)r},$$

where $\mathbf{c}(n)$ is a positive number that is a rational function of q .

2. *If \mathcal{B} is of type B_n ($n \geq 4$) or D_n ($n \geq 5$), then we have*

$$SV(2r) \sim \mathbf{c}_0(n) \cdot r^{\epsilon(n)} q^{2\pi(n)r},$$

$$SV(2r+1) \sim \mathbf{c}_1(n) \cdot r^{\epsilon(n)} q^{2\pi(n)r},$$

where $\mathbf{c}_0(n)$ and $\mathbf{c}_1(n)$ are positive numbers that are rational functions of q .

- *Exceptional types*



- *Exceptional types*



- *Non-irreducible buildings*



- *Exceptional types*



- *Non-irreducible buildings*



- *non-split types*



- *Exceptional types*



- *Non-irreducible buildings*



- *non-split types*



- ***concave functions*** and ***fixed-point sets*** (conjectured)

- Iwasawa, Kenkichi. 1973. *On \mathbf{Z}_l -extensions of algebraic number fields*, Ann. of Math. (2) **98**, 246–326, DOI 10.2307/1970784. MR349627
- Suh, Junecue. 2021. *Stable lattices in p -adic representations i. regular reduction and schur algebra*, Journal of Algebra **575**, 192–219.
- Prasad, Gopal and Jiu-Kang Yu. 2002. *On finite group actions on reductive groups and buildings*, Invent. Math. **147**, no. 3, 545–560. MR1893005
- Garrett, Paul. 1997. *Buildings and classical groups*, Chapman & Hall, London.
- Moy, Allen and Gopal Prasad. 1996. *Jacquet functors and unrefined minimal K -types*, Comment. Math. Helv. **71**, no. 1, 98–121.
- Yu, Jiu-Kang. 2015. *Smooth models associated to concave functions in Bruhat-Tits theory*, Autour des schémas en groupes, pp. 227–258.
- Gao, Xu, *Simplicial volumes in bruhat-tits buildings of split classical type* (2022), arXiv 2210.03328