

Status Finished**Started** Thursday, 7 November 2024, 12:31 PM**Completed** Thursday, 7 November 2024, 12:39 PM**Duration** 7 mins 52 secs**Grade** 10.00 out of 10.00 (100%)**Question 1**

Correct

Mark 1.00 out of 1.00

In the context of logistic regression, which of the following statements is true regarding the logit-link function?

- ☒ a. The logit of the probability p is given by $\log\left(\frac{p}{1-p}\right) = x^T \beta$. \square
- ☐ b. The logit function transforms p to a value between -1 and 1.
- ☐ c. The probability p of an event occurring is given by $p = x^T \beta$.
- ☐ d. Logistic regression does not use the concept of maximum likelihood estimation.

Your answer is correct.

The correct answer is: The logit of the probability p is given by $\log\left(\frac{p}{1-p}\right) = x^T \beta$.

Question 2

Correct

Mark 1.00 out of 1.00

Which of the following models uses the cumulative distribution function of the standard normal distribution in its link function?

- ☐ a. Decision Tree
- ☒ b. Logistic Regression with probit-link \square
- ☐ c. Gaussian Process Regression
- ☐ d. Linear Regression
- ☐ e. Random Forest
- ☐ f. Logistic Regression with logit-link

Your answer is correct.

The correct answer is: Logistic Regression with probit-link

Question 3

Correct

Mark 1.00 out of 1.00

In a logistic regression model, let $\{y_i, x_i\}_{i=1}^n$ represent n independent observations, where y_i is a binary outcome. The likelihood function for a single observation (y_i, x_i) is given by:

$$f(y_i, x_i | \beta) = p_i^{y_i} (1 - p_i)^{1-y_i}$$

where $p_i = \frac{\exp(x_i^T \beta)}{1 + \exp(x_i^T \beta)}$. Which of the following expressions represents the joint likelihood $L(\beta; y, X)$ for the entire sample?

- ☒ a. $\prod_{i=1}^n [p_i^{y_i} (1 - p_i)^{1-y_i}]$ ☐
- ☐ b. $\prod_{i=1}^n [p_i + (1 - p_i)]$
- ☐ c. $\sum_{i=1}^n [\log(p_i^{y_i}) + \log((1 - p_i)^{1-y_i})]$
- ☐ d. $\sum_{i=1}^n [p_i^{y_i} + (1 - p_i)^{1-y_i}]$

Your answer is correct.

The correct answer is: $\prod_{i=1}^n [p_i^{y_i} (1 - p_i)^{1-y_i}]$

Question 4

Correct

Mark 1.00 out of 1.00

In logistic regression, let the probability p_i for the i -th observation be given by:

$$p_i = \frac{\exp(x_i^T \beta)}{1 + \exp(x_i^T \beta)}$$

If the log-likelihood function $\ln L(\beta; y, X)$ is maximized, the Hessian matrix of the log-likelihood function, evaluated at the maximum likelihood estimate $\hat{\beta}$, is typically:

- ☐ a. Positive definite
- ☐ b. Positive semi-definite
- ☐ c. Indefinite
- ☒ d. Negative definite ☐

Your answer is correct.

The correct answer is:

Negative definite

Question 5

Correct

Mark 1.00 out of 1.00

In the context of logistic regression with logit-link, the observed Fisher Information matrix $I(\beta)$ is given by:

$$I(\beta) = -\frac{\partial^2 \ln L(\beta; y, X)}{\partial \beta \partial \beta^T}$$

Which of the following best describes the purpose of $I(\beta)$?

- ☐ a. It gives a measure of model fit.
- ☒ b. It provides an estimate of the variance-covariance matrix of $\hat{\beta}$ for large sample sizes. □
- ☐ c. It provides an estimate of the variance-covariance matrix of $\hat{\beta}$ for small sample sizes.
- ☐ d. It represents the likelihood function evaluated at β .

Your answer is correct.

The correct answer is:

It provides an estimate of the variance-covariance matrix of $\hat{\beta}$ for large sample sizes.

Question 6

Correct

Mark 1.00 out of 1.00

In logistic regression, suppose the asymptotic distribution of the maximum likelihood estimate $\hat{\beta}$ is given by:

$$\sqrt{n}(\hat{\beta} - \beta) \sim N(0, \Omega)$$

Which of the following statements is true about the matrix Ω ?

- ☒ a. It is the variance-covariance matrix of $\hat{\beta}$. □
- ☐ b. It is the identity matrix.
- ☐ c. It is always positive definite, regardless of the data.
- ☐ d. It is a diagonal matrix with entries equal to $p_i(1 - p_i)$.

Your answer is correct.

The correct answer is: It is the variance-covariance matrix of $\hat{\beta}$.

Question 7

Correct

Mark 1.00 out of 1.00

In Linear Discriminant Analysis (LDA), suppose the class-conditional density $f_k(x)$ for class k is modeled as a multivariate Gaussian distribution:

$$f_k(x) = \frac{1}{(2\pi)^{p/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu_k)^T \Sigma^{-1} (x - \mu_k)\right)$$

If we assume $\Sigma_k = \Sigma$ for all classes k , which of the following represents the decision boundary between classes k and l ?

- ☐ a. $(x - \mu_k)^T (\mu_k - \mu_l) = (x - \mu_l)^T (\mu_k - \mu_l)$
- ☐ b. $(x - \mu_k)^T \Sigma^{-1} (x - \mu_k) = (x - \mu_l)^T \Sigma^{-1} (x - \mu_l)$
- ☒ c. $x^T \Sigma^{-1} (\mu_k - \mu_l) = \frac{1}{2}(\mu_k + \mu_l)^T \Sigma^{-1} (\mu_k - \mu_l)$ □
- ☐ d. $x^T \Sigma (\mu_k - \mu_l) = \frac{1}{2}(\mu_k - \mu_l)^T \Sigma (\mu_k - \mu_l)$

Your answer is correct.

The correct answer is:

$$x^T \Sigma^{-1} (\mu_k - \mu_l) = \frac{1}{2}(\mu_k + \mu_l)^T \Sigma^{-1} (\mu_k - \mu_l)$$

Question 8

Correct

Mark 1.00 out of 1.00

In LDA, the discriminant function $\delta_k(x)$ for class k is given by:

$$\delta_k(x) = x^T \Sigma^{-1} \mu_k - \frac{1}{2} \mu_k^T \Sigma^{-1} \mu_k + \log \pi_k$$

Which of the following represents the decision rule $G(x)$ for classifying a new observation x ?

- ☒ a. $G(x) = \arg \max_k \delta_k(x)$ □
- ☐ b. $G(x) = \arg \min_k (x^T \Sigma^{-1} \mu_k)$
- ☐ c. $G(x) = \arg \max_k \left(-\frac{1}{2} \mu_k^T \Sigma^{-1} \mu_k\right)$
- ☐ d. $G(x) = \arg \min_k \delta_k(x)$

Your answer is correct.

The correct answer is: $G(x) = \arg \max_k \delta_k(x)$

Question 9


Correct

Mark 1.00 out of 1.00

In Quadratic Discriminant Analysis (QDA), where $\Sigma_k \neq \Sigma$ for different classes k , the discriminant function $\delta_k(x)$ for class k is given by:

$$\delta_k(x) = -\frac{1}{2} \log |\Sigma_k| - \frac{1}{2} (x - \mu_k)^T \Sigma_k^{-1} (x - \mu_k) + \log \pi_k$$

What type of decision boundary is formed between any two classes k and l ?

- ☐ a. Polynomial of order 3 and above
- ☐ b. Linear
- ☐ c. Circular
- ☒ d. Quadratic 
- ☐ e. Linear Hyper-Plane

Your answer is correct.


The correct answer is: Quadratic

Question 10

Correct

Mark 1.00 out of 1.00

In LDA, suppose we estimate the parameters π_k , μ_k , and Σ from the training data. If there are f_k observations for class k in a total of n observations, what is the maximum likelihood estimate of the prior probability π_k for class k ?

- ☐ a. $\pi_k = \frac{f_k}{2n}$
- ☐ b. $\pi_k = \frac{n}{f_k}$
- ☐ c. $\pi_k = \frac{n-f_k}{n}$
- ☒ d. $\pi_k = \frac{f_k}{n}$ 

Your answer is correct.

The correct answer is:

$$\pi_k = \frac{f_k}{n}$$