

Evidence and feasibility of prone positioning in COVID-19: a resource-challenged perspective

COVID-19 has claimed more than 170 000 lives globally and has completely transformed the way we practice medicine. The majority of deaths are as a result of severe hypoxic respiratory failure and optimising oxygen delivery is the mainstay of treatment. Proning patients with COVID-19 hypoxic failure is widely practised, despite limited evidence. The majority of studies describe only surrogate outcomes such as transient improvement in hypoxemia – without assessing mortality benefit or need for ventilatory support.

Clinical research during a pandemic is very challenging and waiting for good quality RCT's is often not practical, as it may delay access to beneficial treatment options. Prior to the COVID-19 era, evidence for prone positioning during mechanical ventilation in patients with ARDS suggested improved oxygenation and mortality.

From a practical point of view, prone positioning can be challenging and not sustainable for some patients. This session will assess the evidence of prone positioning in COVID-19 and briefly explore the feasibility and safety in resourced challenged settings, where access to ventilators and high-flow oxygen delivery systems are limited.

Dr Clint Hendrikse
MB ChB, MMed, FCEM

Lecturer and Joint staff
Division of Emergency Medicine
University of Cape Town
South Africa

Emergency Physician
Mitchells Plain Hospital and Heideveld Hospital
Western Cape
South Africa