Oral toxicity prediction results for input compound



Predicted LD50: 3919mg/kg

Predicted Toxicity Class: 5

1	2	3	4	5	6

Average similarity: 82.46%

Prediction accuracy: 70.97%

	20%	40%	60%	80%

Name	kaempferol
Molweight	286.24
Number of hydrogen bond acceptors	5
Number of hydrogen bond donors	4
Number of atoms	21
Number of bonds	23
Number of rotable bonds	1
Molecular refractivity	76.01
Topological Polar Surface Area	111.13
octanol/water partition coefficient(logP)	2.28

Toxicity targets

Possible binding to toxicity targets is shown below. For more information on the targets, please click on the individual abbreviations.

no binding			probable binding

AA2AR	ADRB2	ANDR	AOFA	CRFR1	DRD3	ESR1	ESR2	GCR	HRH1	NR1I2	OPRK	OPRM	PDE4D	PGH1	PRGR

Details about possible toxicity targets:

	Toxicity Target	Avg Pharmacophore Fit	Avg Similarity Known Ligands
	Androgen Receptor	1.04%	83.77%
	Amine Oxidase A	26.87%	88.12%
	Dopamine receptor D3	0%	73.11%
	Prostaglandin G/H Synthase 1	33.6%	83%