

# Assignment 6

Gautham Bellamkonda - CS20BTECH11017

Download all latex-tikz codes from

<https://github.com/GauthamBellamkonda/AI1103/tree/main/Assignment6>

## 1 PROBLEM

Let  $X_1, X_2, \dots, X_n$  be a random sample of size  $n$  ( $\geq 2$ ) from a distribution having the probability density function

$$f(x; \theta) = \begin{cases} \frac{1}{\theta} \exp\left(-\frac{x}{\theta}\right) & x > 0, \\ 0, & \text{otherwise,} \end{cases} \quad (1.0.1)$$

where  $\theta \in (0, \infty)$ . Let  $X_{(1)} = \min \{X_1, X_2, \dots, X_n\}$  and  $T = \sum_{i=1}^n X_i$ . Then  $E(X_{(1)}|T)$  equals .....

- (A)  $\frac{T}{n^2}$
- (B)  $\frac{T}{n}$
- (C)  $\frac{(n+1)T}{2n}$
- (D)  $\frac{(n+1)^2 T}{4n^2}$

## 2 SOLUTION

### Lehmann–Scheffé theorem :

If  $T$  is a complete sufficient statistic for  $\theta$  and

$$E(g(T)) = \tau(\theta) \quad (2.0.1)$$

then  $g(T)$  is the uniformly minimum-variance unbiased estimator (UMVUE) of  $\tau(\theta)$ .

We know that

$$T = \sum_{i=1}^n X_i \quad (2.0.2)$$

is a complete and sufficient statistic. By the law of total expectation,

$$E(E(X_{(1)}|T)) = E(X_{(1)}) \quad (2.0.3)$$

By Lehmann–Scheffé theorem, with

$$\theta = X_{(1)}, \quad (2.0.4)$$

$$\tau(x) = E(x), \quad (2.0.5)$$

$$g(T) = E(X_{(1)}|T). \quad (2.0.6)$$

it follows from (2.0.3) that  $E(X_{(1)}|T)$  is the UMVUE of  $E(X_{(1)})$ .

$$\Pr(X_{(1)} > x) = \Pr(X_1 > x) \dots \Pr(X_n > x) \quad (2.0.7)$$

$$= (1 - F_{X_1}(x)) \dots (1 - F_{X_n}(x)) \quad (2.0.8)$$

$$= (1 - F_{X_1}(x))^n \quad (2.0.9)$$

$$= \exp\left(-\frac{nx}{\theta}\right) \quad (2.0.10)$$

$$F_{X_{(1)}}(x) = 1 - \exp\left(-\frac{nx}{\theta}\right) \quad (2.0.11)$$

$$f_{X_{(1)}}(x) = \frac{n}{\theta} \exp\left(-\frac{nx}{\theta}\right) \quad (2.0.12)$$

Therefore,  $X_{(1)}$  follows an exponential distribution with mean  $\frac{\theta}{n}$ .

$$E(X_{(1)}) = \frac{\theta}{n} \quad (2.0.13)$$

Note that,

$$E\left(\frac{T}{n^2}\right) = E\left(\frac{\sum_{i=1}^n X_i}{n^2}\right) \quad (2.0.14)$$

$$= \frac{E(\sum_{i=1}^n X_i)}{n^2} \quad (2.0.15)$$

$$= \sum_{i=1}^n \frac{E(X_i)}{n^2} \quad (2.0.16)$$

$$= \sum_{i=1}^n \frac{\theta}{n^2} \quad (2.0.17)$$

$$= \frac{\theta}{n} \quad (2.0.18)$$

$$= E(X_{(1)}) \quad (2.0.19)$$

Therefore, by Lehmann–Scheffé theorem, with

$$\theta = X_{(1)}, \quad (2.0.20)$$

$$\tau(x) = E(x), \quad (2.0.21)$$

$$g(T) = \frac{T}{n^2}, \quad (2.0.22)$$

it follows that  $\frac{T}{n^2}$  is UMVUE of  $E(X_{(1)})$ .

Since there exists a unique UMVUE for  $E(X_{(1)})$ , it follows that

$$E(X_{(1)}|T) = \frac{T}{n^2} \quad (2.0.23)$$

Hence, option A is correct.




Fig. 4: Theory vs Simulated plot of  $E(X_{(1)}|T)$