



INSTITUT
POLYTECHNIQUE
DE PARIS

NNT : 20XXIPPARIS

Thèse de doctorat

En cas de cotutelle internationale,
mettre ici le logo de
l'établissement de préparation de
la thèse.
(Cadre à enlever)

Logo de l'établissement de
préparation de la thèse. Ou, le cas
échéant, logo de l'établissement co-
délivrant le doctorat en cotutelle
internationale de thèse
(Cadre à enlever)

Information k -means, fragmentation and syntax analysis: A new approach to unsupervised machine learning

Thèse de doctorat de l'Institut Polytechnique de Paris
préparée à l'Ecole nationale de la statistique et de l'administration économique

École doctorale n°000 Dénomination (Sigle)
Spécialité de doctorat : Mathématiques appliquées

Thèse présentée et soutenue à Palaiseau, le 2020, par

GAUTIER APPERT

Composition du Jury :

Prénom Nom	
Statut, Établissement (Unité de recherche)	Président
Prénom Nom	
Statut, Établissement (Unité de recherche)	Rapporteur
Prénom Nom	
Statut, Établissement (Unité de recherche)	Rapporteur
Prénom Nom	
Statut, Établissement (Unité de recherche)	Examinateur
Olivier Catoni	
Directeur de recherche CNRS, ENSAE (UMR CNRS 9194)	Directeur de thèse
Prénom Nom	
Statut, Établissement (Unité de recherche)	Co-directeur de thèse
Prénom Nom	
Statut, Établissement (Unité de recherche)	Invité
Prénom Nom	
Statut, Établissement (Unité de recherche)	Invité

Titre : Information k -means, fragmentation et analyse syntaxique. Une nouvelle approche de l'apprentissage non supervisé.

Mots clés : Apprentissage non supervisé, Classification, Compression de données, bornes PAC-Bayésiennes, Chaînage, Critère des k -means.

Résumé : Le critère de l'information k -means étend le critère des k -means en utilisant la divergence de Kullback comme fonction de perte. La fragmentation est une généralisation supplémentaire permettant l'approximation de chaque signal par une combinaison de fragments.

Nous proposons un nouvel algorithme de fragmentation pour les signaux numériques se présentant comme un algorithme de compression avec perte.

A l'issue de ce traitement, chaque signal est représenté par un ensemble aléatoires de labels, servant d'entrée à une procédure d'analyse syntaxique, conçue comme un algorithme de compression sans perte.

Cet algorithme, fondé sur deux principes appliqués itérativement, la factorisation et le réétiquetage de configurations fréquentes, produit pour chaque signal un arbre syntaxique fournissant une classification hiérarchique des composantes du signal.

Nous avons testé la méthode sur des images en niveaux de gris, sur lesquelles il a été possible de détecter des configurations translatées ou transformées par une rotation. Ceci donne l'espoir d'apporter une réponse à la reconnaissance invariante par transformations fondée sur un critère de com-

pression très général.

D'un point de vue mathématique, nous avons prouvé deux types de bornes. Tout d'abord, nous avons relié notre algorithme de compression à un estimateur implicite d'un modèle statistique lui aussi implicite, à travers un lemme, prouvant que le taux de compression et le niveau de distorsion de l'un sont reliés à l'excès de risque de l'autre. Ce résultat contribue à expliquer la pertinence de nos arbres syntaxiques.

Ensuite, nous établissons des bornes de généralisation non asymptotiques et indépendantes de la dimension pour les différents critères des k -means et critères de fragmentation que nous avons introduits. Nous utilisons pour cela des inégalités PAC-Bayésiennes appliquées dans des espaces de Hilbert à noyau reproduisant.

Par exemple dans le cas des k -means classiques, nous obtenons une borne en $\mathcal{O}(k \log(k)/n)^{1/4}$ qui fournit la meilleure condition suffisante de consistance, à savoir que l'excès de risque tend vers zéro quand $k \log(k)/n$ tend vers zéro. Grâce à une nouvelle méthode de chaînage PAC-Bayésien, nous prouvons aussi une borne en $\mathcal{O}(\log(n/k) \sqrt{k \log(k)/n})$.

Title : Information k -means, fragmentation and syntax analysis. A new approach to unsupervised machine learning.

Keywords : Unsupervised machine learning, Clustering, Data compression, PAC-Bayesian bounds, Chaining, k -means criterion

Abstract : Information k -means is a new mathematical framework that extends the classical k -means criterion, using the Kullback divergence as a distortion measure. The fragmentation criterion is an even broader extension where each signal is approximated by a combination of fragments instead of a single center.

Using the fragmentation criterion as a distortion measure, we propose a new fragmentation algorithm for digital signals, conceived as a lossy data compression scheme.

Based on the output of the fragmentation algorithm, where each signal is described as a random set of labels, we describe a new syntax model, conceived as a lossless data compression scheme.

Our syntax analysis is based on two principles : factorization and relabeling of frequent patterns. It is an iterative scheme, decreasing at each step as much as possible the length of the representation of the training set. It produces for each signal a syntax tree, providing a multi-level classification of the signal components.

We tested the method on grey level digital images, where it was possible to label successfully translated patterns and rotated patterns. This lets us hope that transformation invari-

ant pattern recognition could be approached in a flexible way using a general purpose data compression criterion. From a mathematical point of view, we derived two kinds of generalization bounds. First we defined an implicit estimator based on an implicit statistical model, related to our lossy data compression scheme. We proved a lemma relating the data compression rate and the distortion level of the compression algorithm with the excess risk of the statistical estimator. This explains why our syntax trees may be meaningful.

Second, combining PAC-Bayesian lemmas with the kernel trick, we proved non asymptotic dimension-free generalization bounds for the various information k -means and information fragmentation criteria we introduced.

For instance, in the special case of the classical k -means criterion, we get a non asymptotic dimension free generalization bound of order $\mathcal{O}(k \log(k)/n)^{1/4}$ that gives the best sufficient consistency condition, namely that the excess risk goes to zero when $k \log(k)/n$ goes to zero. Using a new kind of PAC-Bayesian chaining, we also proved a bound of order $\mathcal{O}(\log(n/k) \sqrt{k \log(k)/n})$.

INFORMATION k -MEANS, FRAGMENTATION AND SYNTAX ANALYSIS:
A NEW APPROACH TO UNSUPERVISED MACHINE LEARNING

Gautier Appert

Why the awe for the Second Law ?
The Second Law of Thermodynamics
defines the ultimate purpose of life,
mind, and human striving: to deploy
energy and information to fight back
the tide of entropy and carve out
refuges of beneficial order.

Steven Pinker

Contents

General notation	5
1 Introduction	7
2 Overview	13
2.1. GENERAL IDEAS	13
2.2. GENERALIZATION BOUNDS FOR FRAGMENTATION	23
2.3. DESCRIPTION OF THE SIGNAL FRAGMENTATION ALGORITHM	25
2.4. SYNTAX ANALYSIS	27
2.5. RELATION WITH STATISTICAL ESTIMATION	33
3 Information k-means and information fragmentation algorithms	35
3.1. INFORMATION k -MEANS ALGORITHMS.	35
3.2. INFORMATION FRAGMENTATION	45
3.2.1. RECALL OF THE INFORMATION k -MEANS SETTING	45
3.2.2. FIRST GENERALIZATION: ESTIMATING A JOINT DISTRIBUTION	45
3.2.3. INFORMATION FRAGMENTATION	47
3.3. SIGNAL FRAGMENTATION	51
4 PAC-Bayesian bounds for information k-means and information fragmentation	59
4.1. A PAC-BAYESIAN BOUND FOR INFORMATION k -MEANS.	59
4.1.1. EXPLICIT BOUND IN THE INFORMATION k -MEANS SETTING	67
4.1.2. CLASSICAL k -MEANS QUANTIZATION IN A SEPARABLE HILBERT SPACE	69

4.1.3. DISCUSSION ABOUT THE BOUNDS	70
4.2. A BOUNDED CRITERION FOR INFORMATION k -MEANS.	71
4.3. A BOUNDED CRITERION FOR THE EUCLIDEAN k -MEANS	76
4.4. PAC-BAYESIAN BOUNDS FOR INFORMATION FRAGMENTATION.	78
4.5. FASTER BOUNDS	90
4.5.1. FASTER BOUNDS FOR INFORMATION k -MEANS AND CLASSICAL k -MEANS	90
4.5.2. FASTER BOUNDS FOR BOTH THE BOUNDED INFORMATION k -MEANS AND THE BOUNDED k -MEANS CRITERION	99
4.5.3. FASTER BOUNDS FOR INFORMATION FRAGMENTATION	101
5 Experiment on digital images	111
6 Conclusion	123
A Code highlights	127
References	135

General notation

We will use the following notation throughout this document.

On some measurable probability space Ω , we will consider various random variables $X : \Omega \rightarrow \mathbf{X}$, $Y : \Omega \rightarrow \mathbf{Y}$, etc. that are nothing but measurable functions. We will also consider several probability measures on Ω , and typically two measures \mathbb{P} and $Q \in \mathcal{M}_+^1(\Omega)$, where \mathbb{P} describes the usually unknown data distribution and Q describes an estimation of \mathbb{P} . Then we will use the short notation \mathbb{P}_X for the push forward measure $\mathbb{P} \circ X^{-1}$. Similarly we will let $Q_X = Q \circ X^{-1}$. In the same way $\mathbb{P}_{X,Y} \in \mathcal{M}_+^1(\mathbf{X} \times \mathbf{Y})$ will be the joint distribution of the couple (X, Y) under \mathbb{P} and $\mathbb{P}_{Y|X}$ the corresponding regular conditional probability measure of Y knowing X when it exists. We will always work under sufficient hypotheses to ensure that the decomposition

$$\mathbb{P}_{X,Y} = \mathbb{P}_X \mathbb{P}_{Y|X} \quad (1)$$

is valid, meaning that for any bounded measurable function $f(X, Y)$

$$\int f \, d\mathbb{P}_{X,Y} = \int \left(\int f \, d\mathbb{P}_{Y|X} \right) \, d\mathbb{P}_X.$$

Moreover, we will use the short notation

$$\int f \, d\mathbb{P}_{X,Y} = \mathbb{P}_{X,Y}(f),$$

so that the previous formula becomes

$$\mathbb{P}_{X,Y}(f) = \mathbb{P}_X [\mathbb{P}_{Y|X}(f)].$$

We will often use the Kullback Leibler divergence

$$\mathcal{K}(Q, \mathbb{P}) = \begin{cases} Q \left(\log \left(\frac{dQ}{d\mathbb{P}} \right) \right) & \text{when } Q \ll \mathbb{P}, \\ +\infty & \text{otherwise.} \end{cases}$$

We will always be in this memoire in a situation where the decomposition

LEMMA 1

$$\begin{aligned} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) &= \mathcal{K}(Q_X, \mathbb{P}_X) + Q_X [\mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X})] \\ &= \mathcal{K}(Q_Y, \mathbb{P}_Y) + Q_Y [\mathcal{K}(Q_{X|Y}, \mathbb{P}_{X|Y})] \end{aligned}$$

is valid.

PROOF. It follows from the decomposition (1). A precise statement and a rigorous proof dealing with measurability issues can be found in [Cat04, Appendix section 1.7 page 50]. \square

CHAPTER 1

Introduction

Over the last decades, machine learning and statistical models have been used extensively to treat large amounts of digital data of any type, such as images, speech, and texts ... One of the main problems faced by data scientists is to predict the value of one specific variable as a function of others, based on past history, consigned in a training set. The situation is usually formalized in the following way: we consider a training set

$$\mathcal{D}_n = \{(X_1, Y_1), \dots, (X_n, Y_n)\} \in (\mathcal{X} \times \mathcal{Y})^n$$

consisting in n pairs of independent and identically distributed random variables with unknown joint distribution $\mathbb{P}_{X,Y}$. The goal of the machine learning algorithm is to compute some function $f(X)$ that can predict the outcome Y associated to X . This framework is called supervised learning and can be divided into two categories depending on whether Y is a quantitative or qualitative variable. In the first case, when $Y \in \mathbb{R}^d$, we are dealing with a regression problem and in the second case when Y belongs to a finite set, we are dealing with a classification problem. The computation of the function f is usually performed using an algorithm that minimizes an empirical risk criterion

$$\inf_{f \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^n L(f(X_i), Y_i)$$

where \mathcal{F} represents a certain class of functions and $L(\cdot)$ some loss function adapted to the problem. For instance, in the classification setting, one can take $L(Y, f(X)) = \mathbb{1}(Y \neq f(X))$ and in the regression framework $L(Y, f(X)) = \|Y - f(X)\|_2^2$. The set of functions \mathcal{F} depends also on the setting, typically in the regression framework, one can consider the class of linear predictors $\mathcal{F} = \{f_\beta(X) = X^\top \beta, \beta \in \mathbb{R}^d\}$, whereas in the classification case one can take $\mathcal{F} = \{f_\beta(X) = \mathbb{1}(\mathbb{P}_\beta(Y = 1 | X) \geq 1/2)\}$, where $\mathbb{P}_\beta(Y = 1 | X) = 1/(1 + e^{-X^\top \beta})$, which corresponds to the logistic model.

However, beforehand, data scientists have usually to extract features from the variables (X_1, \dots, X_n) on the occasion of an exploratory analysis. It consists in extracting the essential information carried by the explanatory variables and in some cases in reducing the dimension of the feature space, typically using principal component analysis. This data

preprocessing step is necessary to learn afterwards a classification function f and obtain a better classification performance.

This preprocessing step is comparable to the case when the available data set

$$(X_1, \dots, X_n) \sim \mathbb{P}_X^{\otimes n}$$

does not come with attached labels (Y_1, \dots, Y_n) . The question is then simply to identify patterns or characteristic structures. Typically, one would like to cluster the data into meaningful groups, or to extract informative patterns. This task can be viewed as finding a new representation of the data that highlights its content in a more usable way.

We are touching here on the daunting matter of unsupervised learning, where we do not even know how to measure the quality of the result. Nonetheless, this is a key issue, as most of the available data are unlabelled, and as asking human experts to provide labels in order to train supervised learning algorithms is costly. This need is at the origin of this project, where we propose new ideas with a focus on digital images. We have a double interest in clustering and in changing the representation and will follow a simple guideline : use data compression as a tool for data understanding.

To be more precise, we will describe a series of transformations of the representation of an i.i.d. sample of digital images $(X_1, \dots, X_n) \sim \mathbb{P}_X^{\otimes n}$, where $X_i \in \mathbb{R}^d$. We will start with a lossy coding scheme based on labelling image regions, that can be seen as a multiple labels extension of the k -means algorithm.

This procedure will learn a classification function

$$\ell : \llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket \mapsto \llbracket 1, k \rrbracket$$

defined by the family of product sets

$$\ell^{-1}(j) = A_j \times B_j \subset \llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket$$

and a set of image fragments $C_j \in \mathbb{R}^d$, $1 \leq j \leq k$ where $\text{supp}(C_j) \subset B_j$. They define an approximation Y_i of each image X_i given by the formula

$$Y_i = \sum_{j=1}^k \mathbf{1}(i \in A_j) C_j, \quad i \in \llbracket 1, n \rrbracket,$$

or equivalently by the formula

$$Y_{i,s} = C_{\ell(i,s), s}, \quad i \in \llbracket 1, n \rrbracket, s \in \llbracket 1, d \rrbracket.$$

We will consider a coding distribution $q(\theta)$ on the parameter $\theta = (A, B, C)$, and a measure $D(\theta)$ of the distortion of the representation of (X_1, \dots, X_n) by θ , based on an entropy criterion. We will learn a representation $\widehat{\theta}(X_1, \dots, X_n)$ whose distortion is under a given level and whose ideal code length $-\log_2(q(\widehat{\theta}))$ is as small as possible. The procedure produces image fragments C_j entering into the description of a set of sample images A_j as large as

possible. From this perspective, it can be viewed as a block indexing scheme reminiscent of the Lempel Ziv algorithm. Another interesting feature of our algorithm is that it does not use the image geometry. Although it has been tested on images, it could have been used on any vectors of d measurements. It is indeed invariant by any permutation of the pixel indices and could be without modification applied to other types of digital signals, like rgb images, stereoscopic images, 3D images, video samples, speech signals etc. Moreover the image fragmentation algorithm produces a discrete representation that can be augmented with other discrete descriptors, like text annotations.

Once the fragmentation step is performed, the signal is described as a random set of labels indexing image fragments. Borrowing ideas from linguistic theory, we then learn rewriting rules to compress even more the representation. The algorithm searches repeatedly for the most frequent pair of labels to be reindexed by a single new label. Decoding is then performed by rewriting the new label into the original pair of labels. Again, this compression scheme is reminiscent of the Lempel Ziv algorithm. While in the Lempel Ziv algorithm though, a new indexed block is created by adding one bit to an old block, here a new block is made from a pair of non overlapping old blocks. This type of procedure learns automatically a set of rewriting rules, forming a context free grammar. Further compression steps are then performed by factorizing and compressing again the grammar obtained at the previous step. We obtain a grammar of the grammar that performs a kind of syntax analysis and defines syntax labels. The process can then be repeated to obtain a hierarchy of syntax labels forming a syntax tree (or rather a syntax forest in the general case).

Thus, instead of estimating a statistical model for the data, we learn a grammar through a compression algorithm. The two approaches are different, although a link can be made, considering that a lossless compression code defines a sub-probability measure, due to the Kraft inequality. A good binary code is hard to compress further, so that its distribution is close to a sequence of i.i.d. Bernoulli random variables with parameter 1/2. For this reason, this compression approach can be seen as an alternative to the selection and estimation of a statistical model based on conditional independence properties.

It avoids the difficulty of performing multiple tests of independence that was present in [Mai14]. It allows to bypass the estimation of conditional probability distributions in high dimension and overcomes the problem of model selection. To a certain extent, it is able to circumvent the statistical issues due to the curse of dimensionality. From a practical point of view, since it computes a simpler (compressed) representation of images, there is hope it is scalable and can cope with large datasets (although we did not test this yet).

Nevertheless, we will establish a link with the estimation of the joint probability distribution of random sets of labels. In particular, we will provide a sub-probability measure estimator thanks to the Kraft inequality. Indeed, according to the Kraft inequality, for any prefix binary code $(c_i)_{i \geq 1}$

$$\sum_{i \geq 1} 2^{-l(c_i)} \leq 1,$$

where $l(c_i)$ is the length in bits of c_i .

Then, we will provide an estimation of the law of one single random set using some kind of Bayesian Shtarkov type estimator [Tri16]. We will provide also an oracle inequality to measure the quality of this estimation.

Let us conclude this introduction with a presentation of k -means clustering, the algorithm at the heart of vector quantization and lossy compression that will serve us as a starting point.

The k -means algorithm was suggested by [Llo06] and it is sometimes referred to as Lloyd's algorithm. It provides a partition of a data set $\mathcal{D}_n = \{X_1, \dots, X_n \mid X_i \in \mathbb{R}^p\} \sim \mathbb{P}_X^{\otimes n}$, into k distinct non-overlapping clusters by minimizing the k -means criterion or within-cluster inertia criterion

$$\inf_{\ell: \mathbb{R}^d \rightarrow \{1, \dots, k\}} \inf_{(\mu_1, \dots, \mu_k) \in \mathbb{R}^{d \times k}} \frac{1}{n} \sum_{i=1}^n d(X_i, \mu_{\ell(i)})^2, \quad (1.1)$$

where $\ell : \mathbb{R}^p \rightarrow \{1, \dots, k\}$ denotes the labeling function or cluster assignment function, whereas $(\mu_j)_{j \in \llbracket 1, k \rrbracket}$ are the cluster centers (also called centroids) and $d(\cdot, \cdot)$ indicates some distance or dissimilarity measure. It should be pointed out that criterion (1.1) is the empirical counterpart of the theoretical k -means objective function

$$\inf_{\ell: \mathbb{R}^d \rightarrow \{1, \dots, k\}} \inf_{(\mu_1, \dots, \mu_k) \in \mathbb{R}^{d \times k}} \mathbb{P}_X [d(X, \mu_{\ell(X)})^2],$$

where $\mathbb{P}_X[\cdot]$ denotes the expectation with respect to X . Minimizing the previous criterion (1.1) requires to explore all partitions of $\{1, \dots, n\}$ into k groups. However this task represents a combinatorial optimization problem which is known to be NP-hard and hence infeasible at a practical level.

As an alternative, the k -means algorithm tries to decrease the value of the objective function in an iterative fashion by allocating each data point to the cluster with the nearest centroid and recomputing the center from this partition, see a complete description in algorithm 1 described below. In this way, the k -means algorithm decreases the value of the criterion at each step but converges generally to a local minimum.

Besides, the results obtained depend on the initialization of the k -means algorithm, that is for this reason often randomized. One way to avoid bad clustering due to bad initialization, is to repeat the operation several times and select the one for which the within-cluster inertia criterion is the smallest.

By far, the Euclidean distance $d(x, y) = \|x - y\|_2$ is the most common choice, which leads to compute centroids as empirical means inside each cluster. However, the Euclidean distance can give rise to bad clustering since the empirical mean is not robust. One way to overcome this point is to employ robust estimators for the mean of a random vector. For instance, one may use the estimator suggested in [CG18], since the resulting estimator satisfies tight concentration bounds and is straightforward to compute. An other way to

tackle this issue is to consider other distances more robust to noise and outliers, for instance the L^1 norm.

Algorithm 1 k -means

Initialization :

Pick at random k individuals in the data set that represent the initial centroids of the k clusters.

Iterate the following steps until the within-cluster inertia criterion converges :

- Put each individual in the cluster indexed by the closest centroid, that is for all $1 \leq i \leq n$

$$\ell^*(i) = \arg \min_{j \in \llbracket 1, k \rrbracket} d(x_i, \mu_j)^2.$$

- Compute the centers of gravity of each cluster, that is for all $1 \leq j \leq k$

$$\mu_j^* = \arg \min_{\mu_j} \sum_{i \in \ell^{-1}(j)} d(x_i, \mu_j)^2.$$

These centers of gravity become the new centroids.

return $((\ell^{-1}(j))_{j \in \llbracket 1, k \rrbracket}, (\mu_j)_{j \in \llbracket 1, k \rrbracket})$

In what follows we will introduce a fragmentation algorithm that uses a generalization of the k -means criterion. However, we will propose an algorithm that is very different from Lloyd's algorithm. We will indeed keep the criterion under a given level while optimizing what stands for the number k of centroids. Doing this, we will avoid the difficult question of the choice of k . We will take this opportunity to prove new dimension free generalization bounds for the k -means criterion and for the fragmentation criterion. These non asymptotic bounds decrease like $(k \log(k)/n)^{1/4}$ or like $\log(n/k) \sqrt{k \log(k)/n}$ and improve on previously known ones. They are based on PAC-Bayesian lemmas and some new kind of PAC-Bayesian chaining. Bounds for fragmentation show what quantity takes the place of the number k of centroids in this generalized framework: it is the sum of the sizes of the fragments (normalized by the image size, so that we get back k when we consider in the original k -means setting each centroid as a big fragment covering the whole pixel grid).

CHAPTER 2

Overview

2.1. GENERAL IDEAS

In this study, we will make new proposals for the unsupervised classification of signals, taking the example of digital images.

The aim of unsupervised classification, as we conceive it, is to propose various classification functions, with the hope that at least some of them may be useful to solve interesting tasks.

We will take as a starting point the k -means algorithm with Euclidean distance. Given $(X_1, \dots, X_n) \in \mathbb{R}^{d \times n}$ a training sample, and k centers $c_i, 1 \leq i \leq k$, the empirical loss of the k -means algorithm is

$$L(c_1, \dots, c_k) = \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \|X_i - c_j\|^2 = \inf_{\ell: \mathbb{R}^d \rightarrow \llbracket 1, k \rrbracket} \overline{\mathbb{P}}_X(\|X - c_{\ell(X)}\|^2),$$

where $\overline{\mathbb{P}} = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$ is the empirical measure. This empirical loss is related to the expected loss

$$\inf_{\ell: \mathbb{R}^d \rightarrow \llbracket 1, k \rrbracket} \mathbb{P}_X(\|X - c_{\ell(X)}\|^2).$$

The first thing we will do, starting from there, is to see what we get if we consider the square of the Euclidean norm as a Gaussian Kullback divergence. To make this interpretation, we add to $X \in \mathbb{R}^d$ a second random variable Y such that $\mathbb{P}_{Y|X} = \mathcal{N}(X, \sigma^2 I_d)$, where I_d is the identity matrix of size $d \times d$. We get that

$$\|X - c_{\ell(X)}\|^2 = 2\sigma^2 \mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X}),$$

where $Q_{Y|X} = \mathcal{N}(c_{\ell(X)}, \sigma^2 I_d) = Q_{Y|\ell(X)}$. It is interesting to consider $\mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X})$ rather than $\mathcal{K}(\mathbb{P}_{Y|X}, Q_{Y|X})$ that are equal in this case, because of the following property.

PROPOSITION 2 *The Euclidean k -means criterion can be viewed as an information k -means criterion due to the identity*

$$\inf_{c_1, \dots, c_k} \mathbb{P}_X(\|X - c_{\ell(X)}\|^2) = 2\sigma^2 \inf_Q \mathbb{P}_X[\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X})].$$

The important thing here is that we do not have to restrict the infimum to

$$\left\{ Q : Q_{Y|X} = \mathcal{N}(c_{\ell(X)}, \sigma^2 I_d) \right\}.$$

PROOF. This is a consequence of Proposition 11 on page 41. \square

This proposition shows that the k -means quadratic criterion is a special case of an information k -means criterion

$$\inf_{\ell: \mathbb{R}^d \rightarrow \llbracket 1, k \rrbracket} \inf_Q \mathbb{P}[\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X})],$$

and of its empirical counterpart

$$\inf_{\ell: \mathbb{R}^d \rightarrow \llbracket 1, k \rrbracket} \inf_Q \overline{\mathbb{P}}[\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X})].$$

This information k -means criterion broadens the scope of the k -means algorithm from classifying vectors to classifying distributions. In this case, the data set $X_1, \dots, X_n \sim \mathbb{P}_X^{\otimes n}$ is replaced with a set of conditional probability measures p_{X_1}, \dots, p_{X_n} . For instance in text-mining, histograms made of word counts, called *bags of words* are often used to represent documents. Similarly in computer vision, images may be represented by histograms of visual features. It is worth noticing that bag of visual features are often the result of some clustering algorithm, that may be the k -means algorithm, run on a specific set of image patches to produce a dictionary of features. For more details, one can refer to [Tsa12].

In the information k -means setting, we try to approximate $\mathbb{P}_{Y|X}$ by $Q_{Y|\ell(X)}$. This is an invitation to consider as a variant of the same idea the approximation of the joint distribution $\mathbb{P}_{X,Y}$ by $Q_{X,Y}$ such that $Q_{Y|X} = Q_{Y|\ell(X)}$.

From the disintegration theorem notice that

$$\mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) = \mathcal{K}(Q_X, \mathbb{P}_X) + Q_X[\mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X})].$$

Thus, considering the specific model

$$\mathbf{Q} = \left\{ Q_{X,Y} : Q_X = \mathbb{P}_X, Q_{Y|X} = Q_{Y|\ell(X)}, \ell(X) \in \{1, \dots, k\} \right\},$$

the information k -means can be expressed as an information projection

$$\inf_{Q_{X,Y} \in \mathbf{Q}} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) = \inf_{\ell: \mathbb{X} \rightarrow \llbracket 1, k \rrbracket} \inf_{Q_{Y|\ell(X)} \in \mathcal{M}_+^1(\mathbb{Y})} \mathbb{P}_X[\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X})].$$

The information projection, also called I-projection [Csi75], consists in projecting a probability measure P onto a set \mathbf{Q} of probability distributions, solving

$$\inf_{Q \in \mathbf{Q}} \mathcal{K}(Q, P).$$

This concept appears also in Sanov's theorem [Csi84] which provides a bound on the probability of the empirical measure $\bar{\mathbb{P}}_n$ to belong to a set of probability distributions \mathbf{Q} , informally

$$-\log\left(\mathbb{P}_X^{\otimes n}(\bar{\mathbb{P}}_n \in \mathbf{Q})\right) \sim n \inf_{Q \in \mathbf{Q}} \mathcal{K}(Q, \mathbb{P}_X).$$

The difference between maximum likelihood estimation, that can be written as

$$\hat{\theta}_{\text{MLE}} \in \arg \min_{\theta \in \Theta} \mathcal{K}(\bar{\mathbb{P}}_n, Q_\theta), \quad (2.1)$$

at least when the state space is finite, and I-projection is due to the fact that the Kullback Leibler divergence is not symmetric. In particular it is finite only if its first argument is absolutely continuous with respect to its second argument. In other words, maximum likelihood estimation tends to over-estimate the support of the data distribution, whereas I-projection under-estimate it. The difference in terms the support's estimation is very well illustrated in the Gaussian case in [Bis06], figure 10.2 and 10.3, chap 10. Besides, one can see that (2.1) is equivalent to the maximization of the expectation of a loss function

$$\hat{\theta}_{\text{MLE}} \in \arg \max_{\theta \in \Theta} \bar{\mathbb{P}}_n \left(\log \left(\frac{dQ_\theta}{d\nu} \right) \right),$$

where ν is some dominating measure ($Q_\theta \ll \nu$, for each $\theta \in \Theta$), whereas its theoretical counterpart is written as

$$\theta_{\text{MLE}}^* \in \arg \max_{\theta \in \Theta} \mathbb{P} \left(\log \left(\frac{dQ_\theta}{d\nu} \right) \right).$$

In the same manner, we will propose a loss function for the estimation of the classification parameter $\ell : \mathcal{X} \mapsto \llbracket 1, k \rrbracket$ equivalent to the minimization of the information k -means criterion. Indeed, from Lemma 1 on page 5 and Lemma 6 on page 37, one can remark that

$$\inf_{Q_{X,Y} : Q_{Y|X} = \mu_{\ell(X)}} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) = -\log \left\{ \mathbb{P}_X \left[\exp \left(-\mathcal{K}(\mu_{\ell(X)}, \mathbb{P}_{Y|X}) \right) \right] \right\}, \quad \mu \in \mathcal{M}_+^1(\mathcal{Y})^k.$$

It shows that the minimization of the information k -means criterion is related to the minimization of the expectation of a loss function $\gamma_{\mu,\ell}(X)$

$$\ell^*(\mu) \in \arg \min_{\ell : \mathcal{X} \mapsto \llbracket 1, k \rrbracket} \mathbb{P}_X(\gamma_{\mu,\ell}(X)), \quad (2.2)$$

where $\gamma_{\mu,\ell}(X) = 1 - \exp(-\mathcal{K}(\mu_{\ell(X)}, \mathbb{P}_{Y|X}))$. This loss function is completely observed (we assume that $\mathbb{P}_{Y|X}$ is known) and plays the role of $\log \left(\frac{dQ_\theta}{d\nu} \right)$ in the maximum likelihood framework. Note that the loss function $\gamma_{\mu,\ell}(X)$ belongs to the unit interval, since the Kullback divergence is non negative. We will study the excess risk

$$\mathbb{P}_X(\gamma_{\hat{\mu}, \hat{\ell}}(X)) - \mathbb{P}_X(\gamma_{\mu^*, \ell^*}(X)),$$

where

$$(\hat{\mu}, \hat{\ell}) \in \arg \min_{\mu \in \mathcal{M}_+^1(\mathcal{Y})^k, \ell : \mathcal{X} \mapsto \llbracket 1, k \rrbracket} \bar{\mathbb{P}}_X(\gamma_{\mu,\ell}(X)).$$

Furthermore, information projection appears in many machine learning algorithms, like for instance in variational Bayes methods (VB) for Bayesian estimation. Indeed, VB methods try to approximate a posterior distribution by I-projection onto a family of tractable distributions. These methods represent an alternative to slower MCMC approximations of the posterior distribution. We refer the reader to [Bis06], [BKM17], [AR20] and [ARC16] for more details on this topic. It turns out that VB methods appear also as an appealing tool for unsupervised clustering, especially to compute variational autoencoder (VAE), see [Doe16] for a complete review of the subject. It also emerged in graph variational autoencoders to perform clustering of nodes in a graph, see [Sal+19b] and [Sal+19a].

We should mention that clustering (conditional) probability distributions based on the Kullback divergence or other information criteria is not a new subject. It has been extensively used in text categorization, especially in word clustering to extract features or reduce the original space dimension. For instance, [PTL02] introduce what they call distributional clustering which consists in clustering nouns with respect to the conditional distribution of the associated verb given the noun. The grouping is performed by measuring the Kullback divergence between the conditional distribution knowing each noun and its associated centroid distribution. The centroid distribution is set to an intra-cluster average of conditional distributions that minimizes the average of the Kullback divergence.

However, in the information k -means framework, we follow a different route. We perform the grouping step by minimizing the Kullback divergence with respect to its first argument, which leads to very different centroids, computed as geometric means of conditional distributions. This is to the best of our knowledge a new addition to the literature.

The clustering of conditional distributions in [PTL02] is a particular version of a more general problem called *information bottleneck*, see [TPB01]. In the sequel, we will see that information k -means is a particular version of a more general clustering problem called information fragmentation.

Besides, in [Dhi+03], the authors propose a type of k -means algorithm that decreases a loss function based on the Jensen-Shannon entropy, written also as a loss of mutual information, leading to centers equal to weighted means of conditional distributions. In particular, they show that their entropy criterion can be expressed as a k -means type criterion using the Kullback divergence as the distortion. More formally, their criterion reduces to the following objective problem

$$\inf_{\ell: \mathbf{Q} \rightarrow \llbracket 1, k \rrbracket} \inf_{q_1, \dots, q_k} \sum_{j=1}^k \sum_{i \in \ell^{-1}(j)} \pi_i \mathcal{K}(p_i, q_j),$$

where \mathbf{Q} is a set of discrete probability distributions and $\pi_i > 0$ represents some weights associated with distribution p_i . We can notice here that centroids are computed by minimizing the Kullback divergence with respect to the second argument, so that they are leading to compute centroids as

$$q_j^* = \sum_{i \in \ell^{-1}(j)} \frac{\pi_i p_i}{\sum_{i \in \ell^{-1}(j)} \pi_i},$$

with ℓ fixed and compute the best classification function as

$$\ell^*(i) = \arg \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(p_i, q_j^*).$$

In the same way, [Cao+13] and [Wu12] go further in this direction, studying what they call *Info K-means*. In particular, [Cao+13] proposes a new algorithm to deal with the practical issues of Info- K means, that arise from computing the Kullback divergence in high dimension. They apply this algorithm to cluster a sample of digital images presenting 11 different landmarks. They first preprocess images by extracting visual features and by quantizing those features, in order to consider each image as a bag of visual features. Then, they cluster images using Info- K means with $K = 11$ and obtain promising results by recovering a large portion of the original partition given by the types of landmarks. We should point out that when we will conduct our experiments, we will not use any kind of preprocessing step such as feature selection with quantization. We will perform directly the information fragmentation algorithm on the original digital images and this represents a strong point of our approach. Besides, we refer also to [Jia+11], who propose a k -medoid algorithm to decrease a k -means loss based on the Kullback divergence in both the discrete and continuous cases, and provide in addition an estimator of the Kullback divergence in a continuous setting.

Following the ideas of [Dhi+03], [BDG04] presents a general k -means framework based on the Bregman divergence. The authors show that such criteria can be minimized iteratively. The Bregman distance encompasses many traditional similarity measures such as the Euclidean distance, the Kullback divergence, the logistic loss and many others. However, in the Kullback case, the minimization is performed with respect to the second argument, and not the first as in our proposal.

Coming back to information k -means, we have seen that if we choose freely the distribution Q_X instead of setting it to \mathbb{P}_X , we get the bounded loss function of equation (2.2) on page 15. In the quadratic case, we get the criterion

$$\inf_{\ell} \mathbb{P}_X \left[1 - \exp \left(-\frac{1}{2\sigma^2} \|X - c_{\ell(X)}\|^2 \right) \right] = \mathbb{P}_X \left[1 - \exp \left(-\frac{1}{2\sigma^2} \min_{j \in \llbracket 1, k \rrbracket} \|X - c_j\|^2 \right) \right].$$

We will see that we can state a generalization bound for this kind of robust criterion under weaker hypotheses than for the original criterion.

So far, we have described the extension of k -means to information k -means. The next extension we would like to discuss is from k -means to fragmentation. In the k -means setting, we use a center in \mathbb{R}^d to represent nearby points. This is rather crude from a classification point of view. We do not really expect whole images (or more generally whole signals) to correspond to a relevant class. We would rather like to label *parts* of images. This may be done by labeling pixels in images with different labels, producing a *fragmentation* of each image into various areas. In the usual quadratic k -means setting, we propose to approximate X by

$$Y = \sum_{j \in A_X} c_j, \text{ where } \llbracket 1, d \rrbracket = \bigsqcup_{j \in A_X} \text{supp}(c_j).$$

Here A_X replaces $\ell(X)$ and contains the labels of the components of the signal $X \in \mathbb{R}^d$. This framework can be seen as a generalization of the k -means setting that corresponds to $A_X = \{\ell(X)\}$. The quadratic criterion becomes

$$\mathbb{P}_X \left(\|X - \sum_{j \in A_X} c_j\|^2 \right).$$

Introducing π_j , the orthogonal projection on the vector space spanned by the support of c_j , it can also be written as

$$\mathbb{P}_X \left(\sum_{j \in A_X} \|\pi_j(X) - c_j\|^2 \right) = \sum_{j=1}^k \mathbb{P}_X \left(\mathbb{1}(X \in A_j) \|\pi_j(X) - c_j\|^2 \right),$$

where $A_j = \{x \in \mathbb{R}^d : j \in A_x\}$.

To give an information projection description of fragmentation, we have to introduce the pixel location S as a random variable. More precisely, we replace the representation Y of X with two random variables $S \in \llbracket 1, d \rrbracket$ and $V \in \mathbb{R}$, defined by

$$\mathbb{P}_{S|X} = \frac{1}{d} \sum_{j=1}^d \delta_j \text{ and } \mathbb{P}_{V|X, S=j} = \mathcal{N}(X_j, \sigma^2).$$

This being put, we can describe the quadratic k -means criterion as

$$\inf_Q \mathbb{P}_{X,S} \left[\mathcal{K}(Q_{V|S, \ell(X)}, \mathbb{P}_{V|S, X}) \right],$$

while the global entropy criterion is

$$\begin{aligned} \inf_{Q: Q_{S,V|X} = \mathbb{P}_S Q_{V|S, \ell(X)}} \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V}) \\ = -\log \sup_{Q: Q_{S,V|X} = \mathbb{P}_S Q_{V|S, \ell(X)}} \left\{ \mathbb{P}_X \left[\exp \left(-\mathcal{K}(Q_{S,V|\ell(X)}, \mathbb{P}_{S,V|X}) \right) \right] \right\}. \end{aligned}$$

The modification to be made to obtain the quadratic fragmentation criterion is just to make the classification function ℓ depend on S , the pixel location. We get

$$\inf_Q \mathbb{P}_{X,S} \left[\mathcal{K}(Q_{V|S, \ell(X,S)}, \mathbb{P}_{V|X, S}) \right].$$

The corresponding global entropy criterion is

$$\inf_{Q: Q_{S,V|X, \ell(X,S)} = Q_{S,V|\ell(X,S)}} \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V}).$$

It has the following interesting properties.

PROPOSITION 3 (GLOBAL FRAGMENTATION CRITERION) *Consider k centers $\rho_j \in \mathcal{M}_+^1(\mathbb{R}^d)$, $1 \leq j \leq k$. Define*

$$\mathcal{T}_2 = \left\{ B \subset \llbracket 1, k \rrbracket : \rho_i \perp \rho_j, i \neq j \in B \right\},$$

the set of (possibly partial) tilings by mutually singular probability measures ρ_j . The partial minimum

$$\inf_Q \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V})$$

taken on all probability measures Q , such that for some measurable function $\ell : \mathbb{R}^d \times \llbracket 1, d \rrbracket \rightarrow \llbracket 1, k \rrbracket$

$$Q \left[Q_{X|S,V, \ell(X,S)} = \rho_{\ell(X,S)} \right] = 1. \quad (2.3)$$

is equal to

$$\begin{aligned} - \sup_{\ell} \log \mathbb{P}_S \left(\sum_{j=1}^k \mathbb{1} \left[\rho_j(\ell_S^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \mathbf{Var}_{\rho_j}(X_S)/(2\sigma^2) \right] \right) \\ = - \log \mathbb{P}_S \left(\sup_{B \in \mathcal{T}_2} \sum_{j \in B} \exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \mathbf{Var}_{\rho_j}(X_S)/(2\sigma^2) \right] \right), \end{aligned}$$

where

$$\begin{aligned} \ell_s : \mathbb{R}^d &\rightarrow \llbracket 1, k \rrbracket \\ x &\mapsto \ell(x, s). \end{aligned}$$

For any choice of ℓ , and in particular for the optimal one, putting $W = \ell(X, S)$, the optimum in $Q_{W,S,V}$ is reached when

$$\frac{dQ_{W,S,V}}{d\mathbb{P}_{W,S} \otimes \lambda_V} = Z^{-1} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \mathbf{Var}_{\rho_W}(X_S)/(2\sigma^2) \right] g_{\sigma, \rho_W(X_S)}(V), \quad (2.4)$$

where λ_V is the Lebesgue measure on \mathbb{R} and

$$g_{\sigma, m}(v) = \frac{1}{\sigma \sqrt{2\pi}} \exp \left(-\frac{(v - m)^2}{2\sigma^2} \right).$$

In particular, for the optimal choice of $Q_{W,S,V}$, $Q_{V|S,W} = \mathcal{N}(\rho_W(X_S), \sigma^2)$ is a Gaussian probability measure and

$$\frac{dQ_{S|W}}{d\mathbb{P}_{S|W}} = Z_W^{-1} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \mathbf{Var}_{\rho_W}(X_S)/(2\sigma^2) \right].$$

On the other hand, consider k centers $\mu_{S,V}^{(j)} \in \mathcal{M}_+^1(\llbracket 1, d \rrbracket \times \mathbb{R})$, $1 \leq j \leq k$ such that

$$\mu_{V|S}^{(j)} = \mathcal{N} \left(\mu_{V|S}^{(j)}(V), \sigma^2 \right), \quad 1 \leq j \leq k.$$

Define

$$\mathcal{T}_1 = \left\{ A \subset \llbracket 1, k \rrbracket : \mu_S^{(i)} \perp \mu_S^{(j)}, i \neq j \in A \right\},$$

the set of tilings by mutually singular probability measures $\mu_S^{(j)}$ (or equivalently by mutually singular probability measures $\mu_{S,V}^{(j)}$).

The partial minimum

$$\inf_Q \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V})$$

taken on all probability measures $Q \in \mathcal{M}_+^1(\Omega)$ such that, for some measurable function $\ell : \mathbb{R}^d \times \llbracket 1, d \rrbracket \rightarrow \llbracket 1, k \rrbracket$

$$Q\left[Q_{S,V|X,\ell(X,S)} = \mu_{S,V}^{(\ell(X,S))}\right] = 1, \quad (2.5)$$

is equal to

$$\begin{aligned} & -\sup_{\ell} \log \mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_S^{(j)} \left(\ell_X^{-1}(j) \right) = 1 \right] \right. \\ & \quad \times \exp \left\{ -\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) - \mu_S^{(j)} \left[\left(\mu_{V|S}^{(j)}(V) - X_S \right)^2 / (2\sigma^2) \right] \right\} \left. \right) \\ & = -\log \mathbb{P}_X \left(\sup_{A \in \mathcal{T}_1} \sum_{j \in A} \exp \left\{ -\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) - \mu_S^{(j)} \left[\left(\mu_{V|S}^{(j)}(V) - X_S \right)^2 / (2\sigma^2) \right] \right\} \right), \end{aligned}$$

where

$$\begin{aligned} \ell_x : \llbracket 1, d \rrbracket & \rightarrow \llbracket 1, k \rrbracket \\ s & \mapsto \ell(x, s). \end{aligned}$$

For any value of ℓ , and in particular for the optimal one, considering $W = \ell(X, S)$, the minimum in $Q_{X,W}$ is reached when

$$\frac{dQ_{X,W}}{d\mathbb{P}_{X,W}} = Z^{-1} \exp \left\{ -\mathcal{K}(\mu_S^{(W)}, \mathbb{P}_{S|X,W}) - \mu_S^{(W)} \left[\left(\mu_{V|S}^{(W)}(V) - X_S \right)^2 / (2\sigma^2) \right] \right\}. \quad (2.6)$$

Alternating these two partial optimization steps, we can converge to a local minimum for the optimization problem

$$\inf_Q \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V}),$$

where the infimum is taken over probability measures $Q \in \mathcal{M}_+^1(\Omega)$ satisfying, for some measurable classification function $\ell : \mathbb{R}^d \times \llbracket 1, d \rrbracket \rightarrow \llbracket 1, k \rrbracket$,

$$Q\left[Q_{X,S,V|\ell(X,S)} = Q_{X|\ell(X,S)} \otimes Q_{S,V|\ell(X,S)}\right] = 1. \quad (2.7)$$

The proof will be given later, see Proposition 16 on page 52. The second part of the proposition shows that the entropy criterion can be viewed as an expectation with respect to \mathbb{P}_X . This expectation can be estimated by an expectation with respect to the empirical measure $\bar{\mathbb{P}}_X$. The last part of the proposition describes the pendent of Lloyd's algorithm (in the case where we replace the unknown \mathbb{P}_X by the empirical measure $\bar{\mathbb{P}}_X$).

Now that we have a criterion for fragmentation, we need an algorithm to compute a fragmentation based on this criterion.

We will use the criterion

$$\inf_Q \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V}),$$

to define a distortion function. Let X_1, \dots, X_n be an i.i.d. training set. We can represent its content by the distribution

$$\begin{aligned}\bar{\mathbb{P}}_{I, S, V} &= \left(\frac{1}{n} \sum_{i=1}^n \delta_i \right) \mathbb{P}_{S, V | X=X_I} \\ &= \left(\frac{1}{n} \sum_{i=1}^n \delta_i \right) \left(\frac{1}{d} \sum_{j=1}^d \delta_j \right) \mathcal{N}(X_{I,S}, \sigma^2) \\ &= \frac{1}{nd} \sum_{i=1}^n \sum_{j=1}^d \delta_{i,j} \mathcal{N}(X_{i,j}, \sigma^2).\end{aligned}$$

Here I is a random index ranging in $\llbracket 1, n \rrbracket$. We see immediately that (X_1, \dots, X_n) is a function of $\bar{\mathbb{P}}$, since

$$X_{i,s} = \bar{\mathbb{P}}_{V | S=s, I=i}(V), \quad i \in \llbracket 1, n \rrbracket, \quad s \in \llbracket 1, d \rrbracket.$$

Consider a finite codebook $\mathbf{C} \subset \mathbb{R}$, for instance $\mathbf{C} = \{m2^{-8} : m \in \llbracket 0, 255 \rrbracket\}$ if we are to code light intensities ranging in the unit interval $[0, 1]$ on eight bits as is usually the case. For any classification function

$$\ell : \llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket \longrightarrow \llbracket 1, k \rrbracket$$

defined by

$$\ell^{-1}(j) = A_j \times B_j, \quad 1 \leq j \leq k,$$

where $(A_j \times B_j, 1 \leq j \leq k)$ is a partition of $\llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket$ and any family of centers $(C_j, 1 \leq j \leq k) \in \mathbf{C}^{d \times k}$, where $\text{supp}(C_j) \subset B_j$, define a parameter

$$\theta = (A_j, B_j, C_j)_{j=1}^k$$

and the corresponding model

$$\begin{aligned}\mathbf{Q}_\theta = \Big\{ Q_{I, S, V} \in \mathcal{M}_+^1(\llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket \times \mathbb{R}) \\ : Q_{S, V | I, (I, S) \in A_j \times B_j} = \mathbb{P}_{S | S \in B_j} \mathcal{N}(C_{j,S}, \sigma^2), j \in \llbracket 1, k \rrbracket \Big\},\end{aligned}$$

where we recall that $\mathbb{P}_S = \frac{1}{d} \sum_{s=1}^d \delta_s$ is known and is the uniform measure on the pixel locations. To make a connection between \mathbf{Q}_θ and the model (2.5) on page 20 defined in Proposition 3 on page 18, one can see that \mathbf{Q}_θ is equivalent to impose

$$\mu_S^{(j)} = \mathbb{P}_{S | S \in B_j} \text{ and } \mu_{V|S}^{(j)} = \mathcal{N}(C_{j,S}, \sigma^2).$$

In other words, $\mathbf{Q}_\theta \subset \mathbf{Q}_\ell$, where

$$\mathbf{Q}_\ell = \Big\{ Q_{I, S, V} \in \mathcal{M}_+^1(\llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket \times \mathbb{R}) : Q_{I, S, V | \ell(I, S)} = Q_{I | \ell(I, S)} \otimes Q_{S, V | \ell(I, S)} \Big\}.$$

We define the distortion $D(\theta)$ of the representation of the sample (X_1, \dots, X_n) by the parameter θ as

$$\begin{aligned}
D(\theta) &= \inf_{Q \in \mathbf{Q}_\theta} \left\{ \mathbf{K}(Q_{I,S,V}, \bar{\mathbb{P}}_{I,S,V}) \right\} \\
&= -\log \bar{\mathbb{P}}_I \left(\sum_{j=1}^k \mathbb{1}(I \in A_j) \mathbb{P}_S(B_j) \exp \left\{ -\frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} \left[(X_{I,S} - Y_{I,S})^2 \right] \right\} \right),
\end{aligned}$$

according to Proposition 3 on page 18.

Note that it makes sense to optimize in $Q \in \mathbf{Q}_\theta$, since the quantization of X_i , given by

$$Y_{i,s} = Q_{V|S=s, I=i}(V) = \sum_{j=1}^k \mathbb{1}(i \in A_j) C_{j,s}, \quad 1 \leq i \leq n, \quad 1 \leq s \leq d,$$

does not depend on $Q \in \mathbf{Q}_\theta$, but only on θ . In fact, it does not depend on $Q_{I,S}$ so that, as a variant, we could have optimized even more in the definition of $D(\theta)$.

Note that this notion of distortion satisfies

$$\inf_{Q \in \mathbf{Q}_\theta} \mathbf{K}(Q_{I,S,V}, \bar{\mathbb{P}}_{I,S,V}) \leq D(\theta) \leq \bar{\mathbb{P}}_{I,S} \left[(X_{I,S} - Y_{I,S})^2 \right] = \frac{1}{nd} \|X_1^n - Y_1^n\|^2.$$

Given a coding distribution $q(\theta)$ and an acceptable distortion level $\eta \geq 0$, the fragmentation algorithm will compute a lossy representation $\hat{\theta}(X_1, \dots, X_n)$ with distortion $D(\hat{\theta}) \leq \eta$ and with an ideal code length $-\log(q(\hat{\theta}))$ as small as possible.

We will use a coding distribution of the form

$$q(\theta) = q(A, B, C) = q(A) q(B, C).$$

After this fragmentation step leading to the computation of $\hat{\theta}$, we will perform a syntax analysis step where we will replace the ideal code $q(A)$ by a more efficient code $\tilde{q}(A)$. This improvement will be obtained using the Bayesian Shtarkov approach. More precisely we will consider a family $q_\alpha(A)$ of coding distributions depending on a new parameter α and a prior coding distribution $\mu(\alpha)$, and we will improve on $q(A)$ by considering

$$\tilde{q}(A) = \max_{\alpha} \mu(\alpha) q_\alpha(A).$$

Since

$$\bar{q}(A) = \sum_{\alpha} \mu(\alpha) q_\alpha(A)$$

is a probability measure, \tilde{q} is a subprobability measure and thus a valid coding distribution. From a Bayesian point of view, $\tilde{q}(A)$ can also be seen as the maximum a posteriori probability estimate (MAP). Introducing

$$\hat{\alpha} \in \arg \max_{\alpha} \mu(\alpha) q_\alpha(\hat{A}),$$

we see that $\tilde{q}(\hat{A}) = \mu(\hat{\alpha}) q_{\hat{\alpha}}(\hat{A})$ and that $\hat{\alpha}$ is a function of the sample (X_1, \dots, X_n) , since this is the case for \hat{A} . When we will detail the construction of $\hat{\alpha}$ we will see that it performs some kind of syntax analysis and in particular leads to the computation of a syntax tree for each image X_i , $1 \leq i \leq n$ of the sample.

2.2. GENERALIZATION BOUNDS FOR FRAGMENTATION

The fragmentation algorithm computes a classification function

$$\ell_k : \llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket \longrightarrow \llbracket 1, k \rrbracket.$$

We can deduce from it

$$\begin{aligned} \bar{\ell}_k : \{X_1, \dots, X_n\} \times \llbracket 1, d \rrbracket &\longrightarrow \llbracket 1, k \rrbracket \\ (X_i, s) &\longmapsto \bar{\ell}(X_i, s) = \ell_k(i, s). \end{aligned}$$

A natural question is to extend $\bar{\ell}(x, s)$ to $x \notin \{X_1, \dots, X_n\}$ in some meaningful way. To this purpose, introduce the set of fragments used to represent X_i

$$\bar{A}_i = \{j \in \llbracket 1, k \rrbracket : i \in A_j\}.$$

Remark that $\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) = -\log(\mathbb{P}_S(B_j))$ and noticing that $\mu_{V|S}^{(j)}(V) = C_{j,S}$, one gets from Proposition 3 on page 18

$$\begin{aligned} D(A, B, C) &= -\log \bar{\mathbb{P}}_I \left(\sum_{j \in \bar{A}_I} \mathbb{P}_S(B_j) \exp \left[-\frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} [(C_{j,S} - X_{I,S})^2] \right] \right) \\ &\geq -\log \bar{\mathbb{P}}_X \left(\exp[-D(X, B, C)] \right), \end{aligned}$$

where $(A, B, C) = ((A_j)_{j=1}^k, (B_j)_{j=1}^k, (C_j)_{j=1}^k)$ for short and where

$$D(X, B, C) = -\log \max_{\bar{A} \in \mathcal{T}} \left(\sum_{j \in \bar{A}} \mathbb{P}_S(B_j) \exp \left\{ -\frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} [(X_S - C_{j,S})^2] \right\} \right)$$

and

$$\mathcal{T} = \{\bar{A} \subset \llbracket 1, k \rrbracket : B_i \cap B_j = \emptyset, i \neq j \in \bar{A}\}.$$

We can see $D(X, B, C)$ as the optimal distortion for a single image when it is represented by its best approximation in the codebook (B, C) of image fragments. An optimal set of fragments \bar{A}_X for X is given by the formula

$$\bar{A}_X \in \arg \max_{\bar{A} \in \mathcal{T}} \left(\sum_{j \in \bar{A}} \mathbb{P}_S(B_j) \exp \left\{ -\frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} [(X_S - C_{j,S})^2] \right\} \right). \quad (2.8)$$

It is well defined even when $X \notin \{X_1, \dots, X_n\}$.

We can then define the optimal empirical distortion of the codebook (B, C) as

$$D(B, C) = -\log \bar{\mathbb{P}}_X \left(\exp[-D(X, B, C)] \right) \leq D(A, B, C)$$

and ask for its relationship with its expected counterpart

$$\mathbf{D}(B, C) = -\log \mathbb{P}_X \left(\exp[-D(X, B, C)] \right).$$

For this we need deviation inequalities for

$$\bar{\mathbb{P}}_X \left(\exp[-D(X, B, C)] \right)$$

that are uniform with respect to the parameter (B, C) (here the parameter is the fragments codebook).

Note that the optimal expected distortion $\mathbf{D}(B, C)$ is related to the estimation of the distribution $\mathbb{P}_{X, S, V}$. Indeed, consider the model

$$\begin{aligned} \mathbf{Q}_{B, C} = & \left\{ Q_{X, S, V} \in \mathbf{M}_+^1(\mathbb{R}^d \times \llbracket 1, d \rrbracket \times \mathbb{R}) : \right. \\ & Q_{S, V | X} = \sum_{j \in \bar{A}_X} Q_{S | X}(B_j) \mathbb{P}_{S | S \in B_j} \mathcal{N}(C_{j, S}, \sigma^2) \Big\} \\ = & \left\{ Q_{X, S, V} \in \mathbf{M}_+^1(\mathbb{R}^d \times \llbracket 1, d \rrbracket \times \mathbb{R}) : \right. \\ & \left. Q_{S, V | X, \bar{\ell}(X, S)=j} = \mathbb{P}_{S | S \in B_j} \mathcal{N}(C_{j, S}, \sigma^2) \right\}, \end{aligned}$$

where \bar{A}_X , the optimal set of fragments, is defined by equation (2.8) on page 23 and where, putting $A_j = \{x \in \mathbb{R}^d : j \in \bar{A}_X\}$, $\bar{\ell}$ is defined by the formula

$$\bar{\ell}^{-1}(j) = A_j \times B_j, \quad 1 \leq j \leq k.$$

Remark that

$$\mathbf{D}(B, C) = \inf_{Q \in \mathbf{Q}_{B, C}} \mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V}).$$

Bounding $\mathbf{D}(B, C)$ in terms of $D(B, C)$ thus appears as a generalization bound for some information k -means algorithm.

We will also derive similar bounds for the more classical k -means algorithms described above.

Our proofs will be based on PAC-Bayesian lemmas. We will first rewrite the risk (of information k -means or fragmentation) using a mapping to a reproducing kernel Hilbert space. This will allow to see the risk as the expectation of the minimum of linear functions of the parameter, in a separable Hilbert space of possibly infinite dimension. We will then establish dimension free PAC-Bayesian bounds suitable to this situation. Borrowing ideas from the construction of the isonormal Gaussian process [MPS07, section 3.5], we will use the distribution of an infinite sequence of shifted Gaussian random variables both for the prior and the posterior parameter distribution. We will also use arguments from the proofs of [CG18] and [CG17], concerning the estimation of the mean of a random vector. We will prove generalization bounds going to zero as $k \log(k)/n$ goes to zero, which is better than other bounds already published. Concerning the speed of decrease of the bounds, we first prove bounds in $(k \log(k)/n)^{1/4}$ and then, introducing a more sophisticated chaining argument, bounds in $\log(n/k) \sqrt{k \log(k)/n}$. We will work with weak hypotheses and will in particular not consider the kind of margin assumptions that are necessary to get bounds

decreasing faster than $\sqrt{1/n}$ for a given value of k . See [BDL08], [Fis10], [Lev13], [Lev15] and [BFL20].

To get a $\sqrt{1/n}$ speed as in [BDL08], we take inspiration from the classical chaining procedure for bounding the expected suprema of sub-Gaussian processes (see section 13.1 in [BLM13]). We create a PAC-Bayesian version of chaining in which the concept of δ -net and δ -covering is replaced by the use of a sequence of Gaussian perturbations parametrized by a variance ranging on a logarithmic grid. We combine this PAC-Bayesian chaining with the use of the influence function ψ described in [Cat12] to decompose the excess risk into a sub-Gaussian part and an other part representing extreme values. Doing so, we will recover the speed $1/\sqrt{n}$ proved in [BDL08], but with a better dependence in k , since we get a non asymptotic bound of order $\log(n/k)\sqrt{k\log(k)/n}$ instead of k/\sqrt{n} .

2.3. DESCRIPTION OF THE SIGNAL FRAGMENTATION ALGORITHM

To start with, we will devote to each image $X_i \in \mathbb{R}^d$ of an i.i.d. training set X_i , $1 \leq i \leq n$ a single fragment equal to the whole X_i .

This means that we will start with $k = n$ and a classification function ℓ_n defined by

$$\ell_n^{-1}(j) = A_{n,j} \times B_{n,j} = \{j\} \times \llbracket 1, d \rrbracket.$$

From there, we will iteratively define ℓ_k for larger values of k setting for some pair $J_k \subset \llbracket 1, k \rrbracket$ of labels

$$\begin{aligned} A_{k+1,k+1} &= \bigsqcup_{j \in J_k} A_{k,j}, & A_{k+1,j} &= A_{k,j}, \quad j \in \llbracket 1, k \rrbracket, \\ B_{k+1,k+1} &\subset \bigcap_{j \in J_k} B_{k,j}, & B_{k+1,j} &= \begin{cases} B_{k,j} \setminus B_{k+1,k+1}, & j \in J_k, \\ B_{k,j}, & j \in \llbracket 1, k \rrbracket \setminus J_k. \end{cases} \end{aligned}$$

These equations define ℓ_{k+1} from ℓ_k . We will explain later on how to choose J_k and $B_{k+1,k+1}$. In all cases, however, we can readily remark that for any $k \geq n$,

$$\{A_{k,j} \times B_{k,j} : 1 \leq j \leq k, B_{k,j} \neq \emptyset\}$$

is a partition of $\llbracket 1, n \rrbracket \times \llbracket 1, d \rrbracket$. Note that this partition may contain less than k components (and even less than n components) due to the fact that $B_{k,j}$ is decreasing with $k \geq j$ for a fixed value of j and may become empty. Consider

$$Q_{I,S,V}^{(k)} = \overline{\mathbb{P}}_{I,S} \mathcal{N}(C_{k,\ell_k(I,S),S}, \sigma^2).$$

where

$$C_{k,j,s} = \arg \min \left\{ |c - \overline{\mathbb{P}}_{I|I \in A_{k,j}}(X_{I,s})| : c \in \mathcal{C} \right\}, \quad s \in B_j,$$

and choose

$$\begin{aligned}
B_{k+1,k+1} = & \left\{ s \in \bigcap_{j \in J_k} B_{k,j} : \mathbf{Var}(X_{I,s} | \bar{\mathbb{P}}_{I|I \in A_{k+1,k+1}}) \right. \\
& \left. + \min_{c \in \mathcal{C}} \left(\bar{\mathbb{P}}_{I|I \in A_{k+1,k+1}}(X_{I,s}) - c \right)^2 \leq \alpha \right\}. \quad (2.9)
\end{aligned}$$

Doing so, we are sure at each step that

$$D(A_k, B_k, C_k) \leq \mathbf{K}(Q_{X,S,V}^{(k)}, \bar{\mathbb{P}}_{X,S,V}) \leq \frac{\alpha}{2\sigma^2}.$$

Indeed, from the decomposition of the Kullback divergence (see Lemma 1 on page 5) and the law of iterated expectations, we see that

$$\begin{aligned}
\mathbf{K}(Q_{I,S,V}^{(k)}, \bar{\mathbb{P}}_{I,S,V}) &= \frac{1}{2\sigma^2} \bar{\mathbb{P}}_{I,S} \left[(C_{k,\ell_k(I,S),S} - X_S)^2 \right] \\
&= \frac{1}{2\sigma^2} \sum_{j=1}^k \bar{\mathbb{P}}_{I,S}(A_{k,j} \times B_{k,j}) \bar{\mathbb{P}}_{S|S \in B_{k,j}} \left\{ \bar{\mathbb{P}}_{I|I \in A_{k,j}} \left[(C_{k,j,S} - X_{I,S})^2 \right] \right\} \\
&= \frac{1}{2\sigma^2} \sum_{j=1}^k \bar{\mathbb{P}}_{I,S}(A_{k,j} \times B_{k,j}) \bar{\mathbb{P}}_{S|S \in B_{k,j}} \left[\mathbf{Var}(X_{I,S} | \bar{\mathbb{P}}_{I|I \in A_{k,j}}) + (\bar{\mathbb{P}}_{I|I \in A_{k,j}}(X_{I,S}) - C_{k,j,S})^2 \right].
\end{aligned}$$

We have now to discuss the choice of $J_k = \{i_k, j_k\}$. Assume that $\max\{n, k, |\mathcal{C}|\} < 2^L$, so that all coordinates in (A_k, B_k, C_k) can be coded with L bits. In this case (which is an obvious case of the Kraft inequality),

$$q(A_k, B_k, C_k) = 2^{-L(|A_k|+2|B_k|)},$$

where $|A_k| = \sum_{j=1}^k |A_{k,j}|$, and $|B_k| = \sum_{j=1}^k |B_{k,j}|$, is a sub-probability measure. Remark now that the code length decrease is

$$\log_2(q(A_{k+1}, B_{k+1}, C_{k+1})) - \log_2(q(A_k, B_k, C_k)) = 2L|B_{k+1,k+1}| - L|A_{k+1,k+1}|.$$

Maximizing the code length decrease would lead to choose

$$J_k = \{j_{k,1}, j_{k,2}\} \in \arg \max_{J_k} (2|B_{k+1,k+1}| - |A_{k+1,k+1}|).$$

This requires to compute $A_{k+1,k+1}$ and $B_{k+1,k+1}$ for all possible choices of the pair J_k . We have tested a faster approximation to this minimization, consisting in choosing

$$j_{k,1} \in \arg \max_j (2|B_{k,j}| - |A_{k,j}|)$$

and then

$$j_{k,2} \in \arg \max_{j_{k,2}} (2|B_{k+1,k+1}| - |A_{k+1,k+1}|).$$

This requires to compute $A_{k+1,k+1}$ and $B_{k+1,k+1}$ only for $k-1$ possible values of J_k , namely for

$$J_k \in \left\{ \{j_{k,1}, j\} : j \in \llbracket 1, k \rrbracket \setminus \{j_{k,1}\} \right\},$$

instead of $k(k-1)/2$ possible values if we opt for a full optimization. This heuristic simplification is based on the fact that

$$2|B_{k+1,k+1}| - |A_{k+1,k+1}| \leq 2|B_{k,j_{k,1}}| - |A_{k,j_{k,1}}|,$$

so that the left-hand side cannot be big if the right-hand side is already small, justifying the idea of maximizing the right-hand side to choose $j_{k,1}$ and then the left-hand side to choose $j_{k,2}$.

A natural stopping rule in this framework is to continue as long as we can decrease the code length, that is as long as

$$2|B_{k+1,k+1}| - |A_{k+1,k+1}| > 0.$$

2.4. SYNTAX ANALYSIS

Now that we have a mean to represent an i.i.d. sample of images (X_1, \dots, X_n) by sets of fragments $(\bar{A}_1, \dots, \bar{A}_n)$, drawn from a fragment codebook (B, C) , we will carry further the change of representation by defining a fragment grammar.

Our approach will still be based on compression theory as we have mentioned in the introduction.

The algorithm we will now describe consists in a sequence of lossless codes for the sample $(\bar{A}_1, \dots, \bar{A}_n)$ made of n random sets of labels ranging in $\llbracket 1, k \rrbracket$. As in the previous section, our guide will be to decrease the code length at each step. This can be coded to start with by listing the content of each set, with separators, resulting in the sequence

$$w_{1,1} \dots w_{1,r_1} \wedge w_{2,1} \dots w_{2,r_2} \wedge \dots \wedge w_{n,1} \dots w_{n,r_n},$$

where \wedge is a supplementary label used as a separator and where the labels $w_{i,j} \in \llbracket 1, k \rrbracket$ and \wedge are coded with a prefix binary code (so that they can for instance either be coded with a fixed length code or with a prefix binary code for the integers). The syntax of this initial representation is

$$\{\{w\}\wedge\}\{w\}$$

where we have used $\{\}$ to denote repetition, as in Extended Backus Naur specifications and where $w \leq k$ are initial labels. From there we will put $\bar{A}_i = A_{0,i}$ and choosing a pair of labels $J_1 \subset \llbracket 1, k \rrbracket$ and a new label numbered $k+1$, we will define the new sequence of sets

$$A_{1,i} = \begin{cases} (A_{0,i} \setminus J_1) \cup \{k+1\}, & \text{when } J_1 \subset A_{0,i}, \\ A_{0,i}, & \text{otherwise.} \end{cases}$$

To recover $A_{0,i}, 1 \leq i \leq n$ from $A_{1,i}, 1 \leq i \leq n$, we need to code also the value of $J_1 = \{a, b\}$. We obtain in this way a new code. The syntax for this new code is of the form

$$\{\{w|p\}\wedge\}\{pab\}$$

where $p = k + 1 > k$ is a pair label and where $|$ means *or* as in Extended Backus Naur specifications. In order to shrink as much as possible the length of the representation, we have to choose J_1 to maximize

$$\sum_{i=1}^n \mathbb{1}(J_1 \subset A_{0,i}),$$

the number of times it appears in the random sets $A_{0,i}$, $1 \leq i \leq n$ (at least if we code labels with a fixed length code, as we will assume for simplicity in all this discussion). We can repeat this process choosing at each step a most frequent pair, to form new sequences of random sets

$$A_{0,i}, \dots, A_{m,i}, 1 \leq i \leq n.$$

We obtain a new shorter code at each time by concatenating the code for $A_{m,i}$, $1 \leq i \leq n$ and the list of rewriting rules defining the pairs. The syntax of such a code is thus of the form

$$\{\{w|p\} \wedge\} \{pab\}$$

(We do not need separators in the description of the pairs, since we know they are always triplets.) Note that $w \in \llbracket 1, k \rrbracket$, $p \in \llbracket k + 1, k + m \rrbracket$ and that $a, b \in \llbracket 1, k + m \rrbracket$ may be so to speak either terminal or non terminal indices (or symbols). Note also that we can forget to represent p explicitly in the $\{pab\}$ section of the representation, since we can assume that

$$\{pab\} = p_1 a_1 b_1, \dots, p_m a_m b_m,$$

where $p_i = k + i$. Doing this, we obtain a code of the type

$$\{\{w|p\} \wedge\} \{ab\}$$

Remark that, as long as the chosen pair appears at least three times, we decrease the length of the representation at each step. This can be taken as a stopping rule. We can also use a higher threshold to stop earlier (given that the frequency of the most frequent pair is decreasing).

From an image perspective, this compression step consists in merging two fragments a and b that are frequently seen together in the image sample (X_1, \dots, X_n) . Inside a fixed pair, elements can be viewed as a context of each other. It is rather a "main" context as it is in a way the most frequent one. Therefore, this merging operation is twofold : it simplifies the representation of the set $A_{m,i}$ and determines in the meantime the context of each fragment. Notice that this procedure is very similar to the fragmentation step and can be seen as its dual version in the sense that we exchange the role played by the sets A and B . Indeed, in contrast with fragmentation, here we make $A_{m,i}$ smaller and create a bigger fragment B_p .

Prepare now to push the compression of the representation further by performing permutations on the list of pairs

$$a_1 b_1 \dots a_m b_m$$

Put $a_i = c_{i,1}$ and $b_i = c_{i,2}$. We represent the pairs by the permutation

$$f(c_{\sigma(1),\psi(1,1)})f(c_{\sigma(1),\psi(1,2)}) \dots f(c_{\sigma(m),\psi(m,1)})f(c_{\sigma(m),\psi(m,2)}), \quad (2.10)$$

where the transformation

$$f(c) = \begin{cases} c, & \text{when } c \in \llbracket 1, k \rrbracket, \\ k + \sigma^{-1}(c - k), & \text{when } c \in \llbracket k + 1, k + m \rrbracket. \end{cases}$$

of the pair indices should also be applied to the first part of the representation, that is to $\{\{w|p\} \wedge\}$, where p becomes $f(p)$. Introduce the counter

$$\xi(c) = \sum_{(i,j) \in \llbracket 1, m \rrbracket \times \{1,2\}} \mathbb{1}(c_{i,j} = c)$$

choose

$$c_* \in \arg \max_{c \in \llbracket 1, k+m \rrbracket} \xi(c),$$

and choose

$$\sigma(1), \psi(1, 1), \psi(1, 2), \dots, \sigma(\xi(c_*)), \psi(\xi(c_*), 1), \psi(\xi(c_*), 2)$$

in such a way that

$$c_{\sigma(i),\psi(i,1)} = c_*, \quad 1 \leq i \leq \xi(c_*).$$

Repeat this process on the remaining pairs as many times as needed, until we have chosen the whole permutation $\sigma(i), \psi(i, j)$. After applying transformation (2.10) to the pairs $a_i b_i$, we see that each possible value of a_i comes in succession. Thus we can factorize (2.10) in the form

$$a_1 b_{1,1} \dots b_{1,q_1} \wedge a_2 b_{2,1} \dots b_{2,q_2} \wedge \dots \wedge a_{m'} b_{m',1} \dots b_{m',q_{m'}},$$

where the notation is changed and we need some separators but where $m' \leq m$. (With the new notation $a_1, \dots, a_{m'}$ is a sequence without repetition, whereas in the previous representation (a_1, \dots, a_m) possibly contained repeated values.) Note that the permutation in (2.10) was chosen to maximize sequentially $q_1, q_2, \dots, q_{m'}$. We can then put the sequence $a_1 \dots a_{m'}$ in front to get

$$a_1 \dots a_{m'} \wedge b_{1,1} \dots b_{1,q_1} \wedge b_{2,1} \dots b_{2,q_2} \wedge \dots \wedge b_{m',1} \dots b_{m',q_{m'}}$$

We don't lose information here, since we can recover the previous representation by redistributing each a_i in front of the sequence of contexts $b_{i,1} \dots b_{i,q_i}$. The new syntax for the pairs description is the grammar

$$G ::= \{a\} \wedge \{\{b\} \wedge\} \{b\}$$

instead of $\{ab\}$, so that the syntax for the new sample code is

$$\{\{w|p\} \wedge\} \{a\} \wedge \{\{b\} \wedge\} \{b\} \quad (2.11)$$

instead of

$$\{\{w\} \wedge\} \{w\} \quad (2.12)$$

(We know in the new code that the description of the random sets ends before separator \wedge number n and is followed by the description of the pairs.) Let us note that G is represented as parameter α in section (2) and that $\mu(\alpha) = 2^{-|G|}$, where $|G|$ represents the number of bits used to represent G . On the other hand $q_\alpha(\bar{A}) = 2^{-|D|}$, where $D := \{\{w|p\} \wedge\}$ is the description of \bar{A} . We now remark that the end $\{\{b\} \wedge\} \{b\}$ of the new code (2.11) has the same syntax as the initial code (2.12). We can thus compress it in the same way by indexing iteratively the most frequent syntax pair $J_s = \{c, d\}$ as long as it appears at least three times. Here the frequency is computed within the grammar G , or in other words the frequency is the number of different contexts a_i in which fragment c appears along with d within the list $b_{i,1} \dots b_{i,q_i}$.

This analysis is different from what is usually done in statistics, where frequencies are computed within the data set (X_1, \dots, X_n) . Using syntax analysis, we produce a second level of classification based on a kind of context analysis. This step is necessary if we want to put a common (syntax) label on fragments that appear in different images at different locations. To give an example, let us assume that we have some cats (not necessarily the same) appearing in different images, not in the same place either, but lying on a sofa or on a bed. In that case, we can say that we have a fixed context *the sofa* or *the bed* and that the cats appear frequently in those two contexts. Then, our syntax analysis will tend to put a common syntax label *cat* (or more accurately *what appears on beds and sofas*) on those lazy cats sharing the same context.

As a matter of fact, our approach is very close to the modeling of language in which the syntactic category of a word is usually mainly determined by the type of contexts in which it occurs. Indeed, it is usually impossible to guess syntactic categories from words morphology. To draw a parallel with language modeling, one can think of our images (X_1, \dots, X_n) as sentences $(\bar{A}_1, \dots, \bar{A}_n)$ containing sequences of words $\{\{w\} \wedge\} \{w\}$. However, in our setting the words w belonging to the set \bar{A}_j are not ordered as in language modeling, which makes a major difference. This contextual approach in the case of modeling image patches is not completely new and has been applied to contextual bags of visual words in [Li+11]. However, we use compression both to determine the context and the classification with respect to the context. This differs from classical context modeling that requires to estimate conditional probability distributions in order to determine the context and to perform a contextual classification.

The syntax for the new shorter code for $\{\{b\} \wedge\} \{b\}$ becomes

$$\{\{b|s\} \wedge\} \{cd\}$$

so that the syntax of the new shorter code for the sample becomes

$$\{\{w|p\} \wedge\} \{a\} \wedge \{\{b|s\} \wedge\} \{cd\}$$

This code is made of two parts. The description $D ::= \{\{w|p\} \wedge\}$ of the random sets $A_{0,i}$, $1 \leq i \leq n$ using a mixture of terminal symbols of type $w \in \llbracket 1, k \rrbracket$ and of non terminal symbols of type $p \in \llbracket k+1, k+m \rrbracket$ and the new grammar

$$G ::= \{a\} \wedge \{\{b|s\} \wedge\} \{cd\} \quad (2.13)$$

specifying how to rewrite each p into a (unique) sequence of terminal symbols. This grammar itself has been compressed using non terminal symbols of type s .

Remark now that G defines a classification function f of its symbols. Let us start with the definition of $f(p)$, where $p \in \llbracket k+1, k+m \rrbracket$ is a non terminal symbol. The grammar G contains a (unique) rule $p ::= ab$ and in the compressed representation of the context of a

$$C_a = \{b' : p' ::= ab' \in G\}$$

either b has been left unchanged, in which case we set $f(p) = b$, or b is produced by a unique non terminal syntax label s , in which case we define $f(p) = s$. When j is a terminal symbol, that is when $j \in \llbracket 1, k \rrbracket$, we set $f(j) = j$. We obtain a classification function

$$f : \llbracket 1, k+m \rrbracket \longrightarrow \llbracket 1, k+m+t \rrbracket$$

where t is the number of syntax pairs cd .

We can now recode the description D using the classification function f . For any $j \in \llbracket 1, k+m \rrbracket$ we can write

$$f^{-1}(f(j)) = \{i_1 < i_2 < \dots < i_\ell\}$$

and define $h(j)$ as the solution of the equation $j = i_{h(j)}$. It is easy to see that

$$j \mapsto (f(j), h(j)), \quad j \in \llbracket 1, k+m \rrbracket$$

is one to one. Therefore we can recode the description D in the form

$$\{\{fh\} \wedge\}$$

More precisely we get

$$D = f_{1,1}h_{1,1} \dots f_{1,r_1}h_{1,r_1} \wedge \dots \wedge f_{n,1}h_{n,1} \dots f_{n,r_n}h_{n,r_n}.$$

Gathering the values of h following the same value of f in each element of the sample we can factorize the representation of D to obtain a code of type

$$\{\{f\{h\} \wedge\} \wedge\}$$

where sample elements are separated by a double sign $\wedge\wedge$. Namely

$$\begin{aligned} D = f_{1,1}h_{1,1,1} \dots h_{1,1,q_{1,1}} \wedge \dots \wedge f_{1,r_1}h_{1,r_1,1} \dots h_{1,r_1,q_{1,r_1}} \wedge \wedge \\ \dots \wedge \wedge f_{n,1}h_{n,1,1} \dots h_{n,1,q_{n,1}} \wedge \dots \wedge f_{n,r_n}h_{n,r_n,1} \dots h_{n,r_n,q_{n,r_n}}, \end{aligned}$$

with a change of indexation. This representation can be split into

$$\{\{f\} \wedge\} \{\{h\} \wedge\}$$

where the values of h corresponding to each value of f have been gathered on the right. Namely, we get now


$$D = f_{1,1} \dots f_{1,r_1} \wedge \dots \wedge f_{n,1} \dots f_{n,r_n} \wedge h_{1,1,1} \dots h_{1,1,q_{1,1}} \wedge \dots \wedge h_{n,r_n,1} \dots h_{n,r_n,q_{n,r_n}}.$$

In this way, we obtain a new sample code of type

$$\{\{f\} \wedge\} \{\{h\} \wedge\} G,$$

where G is as in (2.13). This code is made of a new description D made of a family $R := \{\{f\} \wedge\}$ of random sets with syntax labels and of a specification $\{\{h\} \wedge\}$ describing the way to rewrite each syntax label. This is followed by a grammar G describing the meaning of non terminal labels with the help of rewriting rules.

We can now repeat the whole compression process on R to create multiple levels of syntax analysis, resulting in the estimation of a syntax tree for each initial random set \bar{A}_i , $1 \leq i \leq n$.



In the above figure we have represented the set of fragment labels indirectly indexed by a pair label. This is why there may be more than two fragment labels indexed by a single pair label. For instance, if the grammar contains $p_1 ::= w_1 p_2$ and $p_2 ::= w_2 w_3$, then we see that p_1 rewrites to $w_1 w_2 w_3$.

At the end of the syntax analysis stage, we have computed a new shorter binary code T for $(\bar{A}_1, \dots, \bar{A}_n)$. This results in a new lossy code (T, B, C) for the image sample (X_1, \dots, X_n) . The code (T, B, C) defines an approximation (Y_1, \dots, Y_n) of the sample (X_1, \dots, X_n) according to the formula

$$Y_i = \sum_{j \in \bar{A}_i} C_j, \quad 1 \leq i \leq n,$$

where \bar{A}_i can be computed from T . We use an algorithm that ensures that the quadratic distortion is under a given level, namely

$$\bar{\mathbb{P}}_{I,S} [(X_{I,S} - Y_{I,S})^2] = n^{-1} d^{-1} \|X - Y\|^2 \leq \alpha.$$

This is more precisely a property of the fragmentation algorithm, that is maintained afterwards, since the following steps are lossless compression operations.

In this coding approach we do not estimate nor select from data a statistical model, and this is precisely what we wanted to avoid, in the hope of obtaining an algorithm that can lead to meaningful results on smaller samples. This algorithm is very simple as it combines mainly two actions: grouping and context analysis. For this reason, it could inspire new biological neuron models describing the way the mammals' brain is processing perceptive inputs. Indeed, using short codes for frequently concommitent events is compatible with the modeling of a Pavlovian or respondent conditioning. Typically we can have in mind the cat running to eat, as soon as she hears the sound of kibbles falling into her dish.

2.5. RELATION WITH STATISTICAL ESTIMATION

Nonetheless, a link can be made with statistical estimation through the following computations.

The algorithm we described in the previous sections produces a code $\theta(X_1, \dots, X_n) = (T, B, C)$ corresponding to an approximation $(Y_1, \dots, Y_n) = f \circ \theta(X_1, \dots, X_n) = g(X_1, \dots, X_n)$ of (X_1, \dots, X_n) . If we assume that T , B and C are coded by binary prefix codes, we get by concatenation a binary prefix code for θ and we can define $|\theta|$ as the code length of θ in bits. We can then define according to the Kraft inequality a sub-probability measure by the formula $Q_\theta = 2^{-|\theta|}$. Since $Y = f(\theta)$, we get also a sub-probability measure on Y , putting $Q_Y = Q_\theta \circ f^{-1}$.

Consider then the measures $\mathbb{P}_{X, Y, V}$ and $Q_{Y, V}$ defined by

$$\begin{aligned} \mathbb{P}_X &= \mathbb{P}_{X_1}^{\otimes n}, & \mathbb{P}_{Y|X} &= \delta_{g(X)}, \\ \mathbb{P}_{V|X, Y} &= \mathcal{N}(X, \sigma^2 I_{nd}), & Q_{V|Y} &= \mathcal{N}(Y, \sigma^2 I_{nd}). \end{aligned}$$

We see that as previously, $V = (V_1, \dots, V_n)$ is a noisy version of X and we can raise the question of the prediction of \mathbb{P}_{V_n} from the sample (V_1, \dots, V_{n-1}) . Note that $\mathbb{P}_{V_1, \dots, V_n} = \mathbb{P}_{V_1}^{\otimes n}$ is an i.i.d. sequence of vector valued random variables. Consider the progressive estimator based on $Q_{Y, V}$ and defined as

$$\bar{Q}_{V_n | V_1, \dots, V_{n-1}} = \frac{1}{n} \sum_{i=1}^n Q_{V_n | V_1, \dots, V_{i-1}}.$$

Using the convexity of the divergence, the fact that Q_{V_1, \dots, V_n} is exchangeable, and the fact that $\mathbb{P}_{V_1, \dots, V_n}$ is i.i.d, we can prove

LEMMA 4

$$\begin{aligned} \frac{1}{d} \mathbb{P}_{V_1, \dots, V_{n-1}} [\mathcal{K}(\mathbb{P}_{V_n}, \bar{Q}_{V_n | V_1, \dots, V_{n-1}})] &\leq \frac{1}{nd} \mathcal{K}(\mathbb{P}_{V_1, \dots, V_n}, Q_{V_1, \dots, V_n}) \\ &\leq \frac{1}{nd} \mathcal{K}(\mathbb{P}_{Y_1, \dots, Y_n}, Q_{Y_1, \dots, Y_n}) + \frac{\alpha}{2\sigma^2} \leq \frac{1}{nd} \mathcal{K}(\mathbb{P}_\theta, Q_\theta) + \frac{\alpha}{2\sigma^2}. \end{aligned}$$

Therefore, if we can find a coding distribution Q_θ and a distortion level α depending on the sample size n such that the right-hand side goes to zero when n goes to infinity, we obtain a consistent estimator of \mathbb{P}_{V_n} , the distribution of the random image we sampled from convoluted with a Gaussian noise.

The interest of this lemma is of course mainly theoretical. The idea is not to compute $\overline{Q}_{V_n|V_1,\dots,V_{n-1}}$ in practice, but only to show that in the case where the coding distribution Q_θ is sufficiently efficient to make $\frac{1}{nd}\mathcal{K}(\mathbb{P}_\theta, Q_\theta)$ small, then Q_θ contains enough information to estimate \mathbb{P}_V accurately, since $\overline{Q}_{V_n|V_1,\dots,V_{n-1}}$ is a function of Q_θ . This is an indication that θ is a relevant representation of the information contained in \mathbb{P}_{V_n} and therefore of at least some part of the information contained in \mathbb{P}_{X_n} (since V_n is obtained by adding some noise to X_n).

PROOF. By convexity,

$$\mathbb{P}_{V_1,\dots,V_{n-1}}\left[\mathcal{K}(\mathbb{P}_{V_n}, \overline{Q}_{V_n|V_1,\dots,V_{n-1}})\right] \leq \frac{1}{n}\mathbb{P}_{V_1,\dots,V_{n-1}}\left(\sum_{i=1}^n \mathcal{K}(\mathbb{P}_{V_n}, Q_{V_n|V_1,\dots,V_{i-1}})\right).$$

Remark now that

$$Q_{V_n|V_1,\dots,V_{i-1}} = Q_{V_i|V_1,\dots,V_{i-1}},$$

because, Q being exchangeable,

$$Q_{V_1,\dots,V_{i-1},V_n} = Q_{V_1,\dots,V_i}.$$

Moreover, $\mathbb{P}_{V_n} = \mathbb{P}_{V_i}$, so that

$$\begin{aligned} \mathbb{P}_{V_1,\dots,V_{n-1}}\left(\sum_{i=1}^n \mathcal{K}(\mathbb{P}_{V_n}, Q_{V_n|V_1,\dots,V_{i-1}})\right) &= \mathbb{P}_{V_1,\dots,V_{n-1}}\left(\sum_{i=1}^n \mathcal{K}(\mathbb{P}_{V_i}, Q_{V_i|V_1,\dots,V_{i-1}})\right) \\ &= \sum_{i=1}^n \mathbb{P}_{V_1,\dots,V_{i-1}}\left[\mathcal{K}(\mathbb{P}_{V_i|V_1,\dots,V_{i-1}}, Q_{V_i|V_1,\dots,V_{i-1}})\right] = \mathcal{K}(\mathbb{P}_{V_1,\dots,V_n}, Q_{V_1,\dots,V_n}). \end{aligned}$$

This proves the first inequality of the lemma. To prove the second inequality, remark that

$$\mathcal{K}(\mathbb{P}_V, Q_V) \leq \mathcal{K}(\mathbb{P}_{Y,V}, Q_{Y,V}) = \mathcal{K}(\mathbb{P}_Y, Q_Y) + \mathbb{P}_Y\left[\mathcal{K}(\mathbb{P}_{V|Y}, Q_{V|Y})\right].$$

But $\mathbb{P}_{V|Y} = \mathbb{P}_{X|Y}(\mathbb{P}_{V|X,Y}) = \mathbb{P}_{X|Y}(\mathbb{P}_{V|X})$. Accordingly,

$$\mathbb{P}_Y\left[\mathcal{K}(\mathbb{P}_{V|Y}, Q_{V|Y})\right] \leq \mathbb{P}_Y\mathbb{P}_{X|Y}\left[\mathcal{K}(\mathbb{P}_{V|X}, Q_{V|Y})\right] = \frac{1}{2\sigma^2}\mathbb{P}_{X,Y}(\|X - Y\|^2) \leq \frac{nd\alpha}{2\sigma^2}.$$

This proves the second inequality of the lemma. To prove the third, it is enough to note that, since $Y = f(\theta)$,

$$\mathcal{K}(\mathbb{P}_Y, Q_Y) \leq \mathcal{K}(\mathbb{P}_{\theta,Y}, Q_{\theta,Y}) = \mathcal{K}(\mathbb{P}_\theta, Q_\theta).$$

□

CHAPTER 3

Information k -means and information fragmentation algorithms

3.1. INFORMATION k -MEANS ALGORITHMS

In this part of our work, we will present and study various extensions of the k -means algorithm for their own sake. All these variants offer a mean of estimating the distribution $\mathbb{P}_{\mathbb{P}_{Y|X}}$ of a conditional probability measure $\mathbb{P}_{Y|X}$ from the observation of an i.i.d. sample $\mathbb{P}_{Y|X=X_i}$. These variants include the information fragmentation algorithm forming the first step of the syntax analysis scheme described in the overview.

Consider a couple of random variables $(X, Y) \in \mathfrak{X} \times \mathfrak{Y}$, where \mathfrak{X} and \mathfrak{Y} are complete separable metric spaces, so that we can define regular conditional probability measures. Suppose there exists a reference measure $\nu \in \mathcal{M}_+^1(\mathfrak{Y})$ such that $\mathbb{P}(\mathbb{P}_{Y|X} \ll \nu) = 1$. Define $p_X = \frac{d\mathbb{P}_{Y|X}}{d\nu}$. We are interested in the case where $\mathbb{P}_{Y|X}$ is known and represents a bag of words model. This means that each random sample X is described by a random probability measure $\mathbb{P}_{Y|X}$. In the original bag of words model, \mathfrak{Y} is a set of words, and $\mathbb{P}_{Y|X}$ is the distribution of words in a text X drawn at random from some corpus of texts. Here we include the case where \mathfrak{X} and \mathfrak{Y} can be more general measurable spaces.

We consider the following minimization problem also called information k -means problem

$$\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right),$$

where $\llbracket 1, k \rrbracket = \{1, \dots, k\}$, $\mathbb{L}_{+,1}^1(\nu) = \left\{ q \in \mathbb{L}^1(\nu) : q \geq 0, \int q \, d\nu = 1 \right\}$ and

$$\mathcal{K}(q_j, p_X) = \begin{cases} \int q_j \log(q_j/p_X) \, d\nu, & \int q_j \mathbb{1}(p_X = 0) \, d\nu = 0, \\ +\infty, & \text{otherwise} \end{cases}$$

is the Kullback divergence between densities. The purpose of this section is to discuss the general properties of the information k -means problem and to build a mathematical

framework and algorithms to perform the minimization. As we have seen in the overview, we chose to study this algorithm rather than the better known k -means divergence algorithm

$$\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(p_X, q_j) \right)$$

because of Proposition 2 on page 13, showing that our proposal contains the classical Euclidean k -means as a special case.

Let us state some version of the Bayes rule that will be useful in the following discussion.

LEMMA 5 *Let $\mathbb{P}_{X,Y}$ be a joint distribution defined on the product of two Polish spaces. The following statements are equivalent:*

1. *There exists a measure μ such that $\mathbb{P}_{Y|X} \ll \mu$, \mathbb{P}_X almost surely;*
2. *$\mathbb{P}_{Y|X} \ll \mathbb{P}_Y$, \mathbb{P}_X almost surely;*
3. *$\mathbb{P}_{X,Y} \ll \mathbb{P}_X \otimes \mathbb{P}_Y$;*
4. *$\mathbb{P}_{X|Y} \ll \mathbb{P}_X$, \mathbb{P}_Y almost surely.*

Moreover, they imply the following identities between Radon–Nikodym derivatives:

$$\frac{d\mathbb{P}_{X,Y}}{d(\mathbb{P}_X \otimes \mathbb{P}_Y)} = \frac{d\mathbb{P}_{Y|X}}{d\mathbb{P}_Y} = \frac{d\mathbb{P}_{X|Y}}{d\mathbb{P}_X}.$$

PROOF. To prove that 1. implies 2., it is sufficient to show that $\mathbb{P}_{Y|X} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) = 0$, \mathbb{P}_X almost surely. But when 1. is true

$$\mathbb{P}_{Y|X} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) = \int \mathbf{1} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) \frac{d\mathbb{P}_{Y|X}}{d\mu} d\mu.$$

Thus by the Tonelli-Fubini theorem

$$\begin{aligned} \mathbb{P}_X \left(\mathbb{P}_{Y|X} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) \right) &= \mathbb{P}_X \left(\int \mathbf{1} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) \frac{d\mathbb{P}_{Y|X}}{d\mu} d\mu \right) \\ &= \int \mathbf{1} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) \mathbb{P}_X \left[\frac{d\mathbb{P}_{Y|X}}{d\mu} \right] d\mu \\ &= \int \mathbf{1} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) \frac{d\mathbb{P}_Y}{d\mu} d\mu = 0. \end{aligned}$$

Therefore $\mathbb{P}_{Y|X} \left(\frac{d\mathbb{P}_Y}{d\mu} = 0 \right) = 0$, \mathbb{P}_X almost surely. Obviously 2. implies 1. Now let us show that 2. implies 3. Let f be a bounded measurable function, we have by Fubini's theorem

$$\begin{aligned} \int f d\mathbb{P}_{X,Y} &= \int \left(\int f d\mathbb{P}_{Y|X} \right) d\mathbb{P}_X = \int \left(\int f \frac{d\mathbb{P}_{Y|X}}{d\mathbb{P}_Y} d\mathbb{P}_Y \right) d\mathbb{P}_X \\ &= \int f \frac{d\mathbb{P}_{Y|X}}{d\mathbb{P}_Y} d(\mathbb{P}_Y \otimes d\mathbb{P}_X), \end{aligned}$$

implying 3. and that \mathbb{P}_X almost surely

$$\frac{d\mathbb{P}_{Y|X}}{d\mathbb{P}_Y} = \frac{d\mathbb{P}_{X,Y}}{d(\mathbb{P}_X \otimes \mathbb{P}_Y)}.$$

We will show now that 3. implies 2. Let f be a bounded measurable function, we have by Fubini's theorem

$$\begin{aligned} \int f d\mathbb{P}_{X,Y} &= \int f \frac{d\mathbb{P}_{X,Y}}{d(\mathbb{P}_X \otimes \mathbb{P}_Y)} d(\mathbb{P}_X \otimes d\mathbb{P}_Y) \\ &= \int \left(\int f \frac{d\mathbb{P}_{X,Y}}{d(\mathbb{P}_X \otimes \mathbb{P}_Y)} d\mathbb{P}_Y \right) d\mathbb{P}_X \\ &= \int \left(\int f d\mathbb{P}_{Y|X} \right) d\mathbb{P}_X, \end{aligned}$$

showing that \mathbb{P}_X almost surely $\mathbb{P}_{Y|X} \ll \mathbb{P}_Y$ and

$$\frac{d\mathbb{P}_{Y|X}}{d\mathbb{P}_Y} = \frac{d\mathbb{P}_{X,Y}}{d(\mathbb{P}_X \otimes \mathbb{P}_Y)}.$$

The equivalence between 3. and 4. is immediate by interchanging the roles of X and Y . \square

The following lemma will be useful to optimize the information k -means criterion.

LEMMA 6 *Let $\pi \in \mathcal{M}_+^1(\Omega)$ be a probability measure on the measurable space Ω . Let $h : \Omega \rightarrow \mathbb{R} \cup \{+\infty\}$ be a measurable function such that*

$$Z = \int \exp(-h) d\pi < \infty.$$

Let $\pi_{\exp(-h)}$ be the probability measure whose density with respect to π is proportional to $\exp(-h)$ so that

$$\frac{d\pi_{\exp(-h)}}{d\pi} = \frac{\exp(-h)}{Z}.$$

The identity

$$\inf_{\eta \in \mathbb{Z}} \left(\mathcal{K}(\rho, \pi) + \int \max\{h, \eta\} d\rho \right) = -\log \left(\int \exp(-h) d\pi \right) + \mathcal{K}(\rho, \pi_{\exp(-h)}) \in \mathbb{R} \cup \{+\infty\}$$

is satisfied for any $\rho \in \mathcal{M}_+^1(\Omega)$ and implies that

$$\inf_{\rho \in \mathcal{M}_+^1(\Omega)} \inf_{\eta \in \mathbb{Z}} \left(\mathcal{K}(\rho, \pi) + \int \max\{h, \eta\} d\rho \right) = -\log \left(\int \exp(-h) d\pi \right),$$

the minimum being reached when $\rho = \pi_{\exp(-h)}$.

Note that the lemma could also be written as

$$\mathcal{K}(\rho, \pi) + \int h d\rho = -\log \left(\int \exp(-h) d\pi \right) + \mathcal{K}(\rho, \pi_{\exp(-h)})$$

if we are willing to follow the convention that

$$\int h \, d\rho = \inf_{\eta \in \mathbb{Z}} \int \max\{h, \eta\} \, d\rho$$

and that $+\infty - \infty = +\infty$.

PROOF. See [Cat04, page 159]. Note that the role of $\eta \in \mathbb{Z}$ in this lemma is only to make sure that the integrals are always well defined in $\mathbb{R} \cup \{+\infty\}$ in the sense that the negative part of the integrand is integrable. When ρ is not absolutely continuous with respect to π , it is also not absolutely continuous with respect to $\pi_{\exp(-h)}$ since $\pi(A) = 0$ if and only if $\pi_{\exp(-h)}(A) = 0$. In this case $\mathcal{K}(\rho, \pi) = \mathcal{K}(\rho, \pi_{\exp(-h)}) = +\infty$ and the identity is true, both sides being equal to $+\infty$. When $\rho \ll \pi$, then $\rho \ll \pi_{\exp(-\max\{h, \eta\})}$ and

$$\frac{d\rho}{d\pi_{\exp(-\max\{h, \eta\})}} = Z_\eta \exp(\max\{h, \eta\}) \frac{d\rho}{d\pi},$$

where

$$Z_\eta = \int \exp(-\max\{h, \eta\}) \, d\pi < +\infty.$$

Therefore

$$\mathcal{K}(\rho, \pi_{\exp(-\max\{h, \eta\})}) = \log(Z_\eta) + \int \left[\max\{h, \eta\} + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho.$$

By the monotone convergence theorem

$$\lim_{\eta \rightarrow -\infty} Z_\eta = Z \text{ and } \lim_{\eta \rightarrow -\infty} \int \left[\max\{h, \eta\} + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho = \int \left[h + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho,$$

since we know that

$$\int \left[\log(Z) + h + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho = \int \log\left(\frac{d\rho}{d\pi_{\exp(-h)}}\right) d\rho - \frac{d\rho}{d\pi_{\exp(-h)}} d\pi_{\exp(-h)} \leq \exp(-1) < +\infty$$

and therefore that

$$\int \left[h + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho < +\infty.$$

This proves that

$$\begin{aligned} \lim_{\eta \rightarrow -\infty} \mathcal{K}(\rho, \pi_{\exp(-\max\{h, \eta\})}) &= \log(Z) + \int \left[h + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho = \mathcal{K}(\rho, \pi_{\exp(-h)}) \\ &= \log(Z) + \inf_{\eta \in \mathbb{Z}} \int \left[\max\{h, \eta\} + \log\left(\frac{d\rho}{d\pi}\right) \right] d\rho \\ &= \log(Z) + \inf_{\eta \in \mathbb{Z}} \left(\int \max\{h, \eta\} d\rho + \int \log\left(\frac{d\rho}{d\pi}\right) d\rho \right) \\ &= \log(Z) + \inf_{\eta \in \mathbb{Z}} \left(\int \max\{h, \eta\} d\rho + \mathcal{K}(\rho, \pi) \right), \end{aligned}$$

and therefore that

$$\mathcal{K}(\rho, \pi_{\exp(-h)}) - \log(Z) = \inf_{\eta \in \mathbb{Z}} \left(\mathcal{K}(\rho, \pi) + \int \max\{h, \eta\} d\rho \right)$$

as stated in the lemma. The second statement of the lemma is a consequence of the fact that the Kullback divergence is non negative. \square

LEMMA 7 Let $\mathbb{P}_{X,Y}$ be a joint distribution defined on the product of two Polish spaces. Assume that $\mathbb{P}_X(\mathbb{P}_{Y|X} \ll \mathbb{P}_Y) = 1$. Consider the normalizing constant

$$Z = \mathbb{P}_Y \left(\exp[-\mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y})] \right).$$

Obviously, $Z \in [0, 1]$. If $Z = 0$, then

$$\inf_{Q_Y \in \mathcal{M}_+^1(\mathcal{Y})} \mathbb{P}_X[\mathcal{K}(Q_Y, \mathbb{P}_{Y|X})] = +\infty.$$

Otherwise, $Z > 0$ and for any $Q_Y \in \mathcal{M}_+^1(\mathcal{Y})$,

$$\mathbb{P}_X[\mathcal{K}(Q_Y, \mathbb{P}_{Y|X})] = \mathcal{K}(Q_Y, Q_Y^*) + \mathbb{P}_X[\mathcal{K}(Q_Y^*, \mathbb{P}_{Y|X})] = \mathcal{K}(Q_Y, Q_Y^*) + \log(Z^{-1}),$$

where $Q_Y^* \ll \mathbb{P}_Y$ is defined by the relation

$$\begin{aligned} \frac{dQ_Y^*}{d\mathbb{P}_Y} &= Z^{-1} \exp[-\mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y})] \\ &= Z^{-1} \exp \left\{ \mathbb{P}_X \left[\log \left(\frac{d\mathbb{P}_{Y|X}}{d\mathbb{P}_Y} \right) \right] \right\}. \end{aligned} \quad (3.1)$$

Consequently

$$\inf_{Q_Y \in \mathcal{M}_+^1(\mathcal{Y})} \mathbb{P}_X[\mathcal{K}(Q_Y, \mathbb{P}_{Y|X})] = \mathbb{P}_X[\mathcal{K}(Q_Y^*, \mathbb{P}_{Y|X})] = \log(Z^{-1}) < \infty,$$

The probability measure Q_Y^* represents the geometric mean of $\mathbb{P}_{Y|X}$ with respect to \mathbb{P}_X .

PROOF. By Lemma 1 on page 5,

$$\mathbb{P}_X[\mathcal{K}(Q_Y, \mathbb{P}_{Y|X})] = \mathcal{K}(\mathbb{P}_X \otimes Q_Y, \mathbb{P}_{X,Y}) = \mathcal{K}(Q_Y, \mathbb{P}_Y) + Q_Y[\mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y})]. \quad (3.2)$$

Thus, when (3.2) is finite, $Q_Y \ll \mathbb{P}_Y$ and

$$Q_Y[\mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y}) < +\infty] = 1,$$

so that

$$\mathbb{P}_Y[\mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y}) < +\infty] > 0,$$

implying that $Z > 0$. Assuming from now on that (3.2) is finite, introduce

$$\mathcal{A} = \left\{ y : \mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y=y}) < +\infty \right\}.$$

From Lemma 6 on page 37 and (3.2), for any $Q_Y \in \mathcal{M}_+^1(\mathcal{A})$,

$$\begin{aligned} \mathbb{P}_X[\mathcal{K}(Q_Y, \mathbb{P}_{Y|X})] &= -\underbrace{\log[\mathbb{P}_Y(\mathcal{A})] - \log \mathbb{P}_{Y|Y \in \mathcal{A}} \left\{ \exp[-\mathcal{K}(\mathbb{P}_X, \mathbb{P}_{X|Y})] \right\}}_{=\log(Z^{-1})} + \mathcal{K}(Q_Y, Q_Y^*) \\ &= \mathbb{P}_X[\mathcal{K}(Q_Y^*, \mathbb{P}_{Y|X})] + \mathcal{K}(Q_Y, Q_Y^*). \end{aligned}$$

Moreover, when $Q_Y(\mathcal{A}) < 1$, $Q_Y \not\ll Q_Y^*$, so that both members are equal to $+\infty$. The identity (3.1) is a consequence of Lemma 5 on page 36. \square

PROPOSITION 8 *The information k -means problem can be expressed as*

$$\begin{aligned}
\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \inf_{(q_1, \dots, q_k) \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}, p_X) \right) \\
&= \inf_{(q_1, \dots, q_k) \in (\mathbb{L}_{+,1}^1(\nu))^k} \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}, p_X) \right) \\
&= \inf_{(q_1, \dots, q_k) \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\mathcal{K}(q_{\ell_q^*(X)}, p_X) \right) \\
&= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}^{\star,\ell}, p_X) \right) \\
&= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \mathbb{P}_X \left(\log(Z_{\ell(X)}^{-1}) \right),
\end{aligned}$$

where $\ell_q^*: \mathcal{X} \mapsto \llbracket 1, k \rrbracket$ is the best classification function for a fixed $q = (q_1, \dots, q_k)$ defined as

$$\ell_q^*(x) = \arg \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_x), \quad x \in \mathcal{X},$$

whereas $q_1^{\star,\ell}, \dots, q_k^{\star,\ell}$ are the best information k -means centers with respect to $\ell(X)$ defined as

$$q_j^{\star,\ell} = Z_j^{-1} \exp \left\{ \mathbb{P}_{X \mid \ell(X)=j} [\log(p_X)] \right\}, \quad j \in \llbracket 1, k \rrbracket,$$

where

$$Z_j = \int \exp \left\{ \mathbb{P}_{X \mid \ell(X)=j} [\log(p_X)] \right\} d\nu,$$

with the convention that $q_j^{\star,\ell}$ can be given any arbitrary value in the case when $Z_j = 0$, the corresponding criterion being in this case infinite. Besides, we have the following Pythagorean identity

$$\mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}, p_X) \right) = \mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}^{\star,\ell}, p_X) \right) + \mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}^{\star,\ell}, q_{\ell(X)}^{\star,\ell}) \right).$$

PROOF. This proposition is a straightforward consequence of Lemma 7 applied to $\mathbb{P}_{X, Y \mid \ell(X)=j}$. \square

Let us state the empirical counterpart of the previous proposition.

COROLLARY 9 *Let X_1, \dots, X_n be an i.i.d sample drawn from \mathbb{P}_X . Then, the empirical version of the information k -means problem tries to partition the observations p_{X_1}, \dots, p_{X_n} into k -clusters, what is expressed here by*

$$\begin{aligned}
\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_{X_i}) &= \inf_{\ell: \llbracket 1, n \rrbracket \rightarrow \llbracket 1, k \rrbracket} \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \frac{1}{n} \sum_{i=1}^n \mathcal{K}(q_{\ell(i)}, p_{X_i}) \\
&= \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \inf_{\ell: \llbracket 1, n \rrbracket \rightarrow \llbracket 1, k \rrbracket} \frac{1}{n} \sum_{i=1}^n \mathcal{K}(q_{\ell(i)}, p_{X_i}) = \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \frac{1}{n} \sum_{i=1}^n \mathcal{K}(q_{\ell_q^*(i)}, p_{X_i}) \\
&= \inf_{\ell: \llbracket 1, n \rrbracket \rightarrow \llbracket 1, k \rrbracket} \frac{1}{n} \sum_{i=1}^n \mathcal{K}(q_{\ell(i)}^{\star,\ell}, p_{X_i}) = \inf_{\ell: \llbracket 1, n \rrbracket \rightarrow \llbracket 1, k \rrbracket} \sum_{j=1}^k \frac{|\ell^{-1}(j)|}{n} \log(Z_j^{-1}),
\end{aligned}$$

where $\ell_q^* : \mathfrak{X} \mapsto \llbracket 1, k \rrbracket$ is the best classification function for a fixed $q = (q_1, \dots, q_k)$ defined as

$$\ell_q^*(i) = \arg \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_{X_i})$$

whereas $q_j^{*,\ell}, j \in \llbracket 1, k \rrbracket$ are the information k -means centers defined as

$$q_j^{*,\ell} = Z_j^{-1} \left(\prod_{i \in \ell^{-1}(j)} p_{X_i} \right)^{1/|\ell^{-1}(j)|},$$

where

$$Z_j = \int \left(\prod_{i \in \ell^{-1}(j)} p_{X_i} \right)^{1/|\ell^{-1}(j)|} d\nu.$$

PROOF. Apply the previous proposition to the empirical measure $\overline{\mathbb{P}}_X = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$ of the sample X_1, \dots, X_n . \square

LEMMA 10 *Let us assume that $\mathbb{P}_X \left(\int p_X^2 d\nu \right) < \infty$. Then, the optimal centers $q_j^{*,\ell}$ defined in the previous lemma verify $q_j^{*,\ell} \in \mathbb{L}^2(\nu)$. Furthermore, in this case*

$$\begin{aligned} \inf \left\{ \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) : q \in \left(\mathbb{L}_{+,1}^1(\nu) \right)^k \right\} \\ = \inf \left\{ \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) : q \in \left(\mathbb{L}_{+,1}^1(\nu) \cap \mathbb{L}^2(\nu) \right)^k \right\}. \end{aligned}$$

PROOF. Apply Jensen's inequality and the Fubini-Tonelli theorem to obtain that $q_j^{*,\ell} \in \mathbb{L}^2(\nu)$. Indeed, for any $j \in \llbracket 1, k \rrbracket$, if $Z_j = 0$, we can pick up any value for $q_j^{*,\ell}$, and in particular a value in $\mathbb{L}^2(\nu)$, in the same way if $\mathbb{P}_X(\ell(X) = j) = 0$, we can make an arbitrary choice for $q_j^{*,\ell}$, otherwise, $Z_j > 0$, and

$$\begin{aligned} \int (q_j^{*,\ell})^2 d\nu &= Z_j^{-2} \int \exp \left\{ 2\mathbb{P}_{X|\ell(X)=j} [\log(p_X)] \right\} d\nu \\ &\leq Z_j^{-2} \mathbb{P}_{X|\ell(X)=j} \left(\int p_X^2 d\nu \right) \leq Z_j^{-2} \mathbb{P}_X(\ell(X) = j)^{-1} \mathbb{P}_X \left(\int p_X^2 d\nu \right) < \infty \end{aligned}$$

Then according to Proposition 8 on the facing page

$$\begin{aligned} \mathbb{P}_X \left[\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right] &= \inf_{\ell: \mathfrak{X} \mapsto \llbracket 1, k \rrbracket} \mathbb{P}_X \left[\mathcal{K}(q_{\ell(X)}, p_X) \right] \\ &\geq \inf_{\ell: \mathfrak{X} \mapsto \llbracket 1, k \rrbracket} \mathbb{P}_X \left[\mathcal{K}(q_{\ell(X)}^{*,\ell}, p_X) \right] \geq \inf_{\ell: \mathfrak{X} \mapsto \llbracket 1, k \rrbracket} \mathbb{P}_X \left[\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j^{*,\ell}, p_X) \right], \end{aligned}$$

showing that we can restrict the optimization to $q_j \in \mathbb{L}^2(\nu)$. \square

PROPOSITION 11 *Consider the particular case when $\mathfrak{X} = \mathfrak{Y} = \mathbb{R}^p$ and choose as reference measure the Gaussian measure $\nu = \mathcal{N}_p(0_{\mathbb{R}^p}, \sigma^2 I_p)$. Choose also $\mathbb{P}_{Y|X} = \mathcal{N}_p(X, \sigma^2 I_p)$ where*

$\sigma > 0$. In this particular setting, the information k -means problem is identical to the classical euclidian k -means problem. Namely

$$\begin{aligned} \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \frac{1}{2\sigma^2} \mathbb{P}_X \left(\|X - \mathbb{P}_{X|\ell(X)}(X)\|_2^2 \right) \\ &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \inf_{\mu_1, \dots, \mu_k \in \mathbb{R}^{p \times k}} \frac{1}{2\sigma^2} \mathbb{P}_X \left(\|X - \mu_{\ell(X)}\|_2^2 \right) \\ &= \inf_{\mu_1, \dots, \mu_k \in \mathbb{R}^{p \times k}} \frac{1}{2\sigma^2} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|_2^2 \right). \end{aligned}$$

PROOF. In this situation, p_X is equal to

$$\begin{aligned} p_X(y) &= \frac{d\mathcal{N}_p(X, \sigma^2 I_p)}{d\nu}(y) = \exp \left\{ -\frac{1}{2\sigma^2} \left(\|y - X\|_2^2 - \|y\|_2^2 \right) \right\} \\ &= \exp \left\{ -\frac{1}{2\sigma^2} \left(\|X\|_2^2 - 2y^\top X \right) \right\}. \end{aligned}$$

According to Proposition 8 on page 40 for a given classification function ℓ , the optimal centers in the information k -means problem are given by

$$q_j^{\star,\ell} = Z_j^{-1} \exp \left\{ \mathbb{P}_{X|\ell(X)=j} [\log(p_X)] \right\}, \quad j \in \llbracket 1, k \rrbracket.$$

Accordingly

$$q_j^{\star,\ell}(y) = Z_j^{-1} \exp \left\{ -\frac{1}{2\sigma^2} \mathbb{P}_{X|\ell(X)=j} \left(\|X\|_2^2 - 2y^\top X \right) \right\} = \frac{d\mathcal{N}_p(\mathbb{P}_{X|\ell(X)=j}(X), \sigma^2 I_p)}{d\nu}.$$

Then, computing the Kullback divergence between two multivariate normal distributions, we obtain

$$\mathcal{K}(q_j^{\star,\ell}, p_X) = \mathcal{K} \left(\mathcal{N}_p(\mathbb{P}_{X|\ell(X)=j}(X), \sigma^2 I_p), \mathcal{N}_p(X, \sigma^2 I_p) \right) = \frac{1}{2\sigma^2} \|X - \mathbb{P}_{X|\ell(X)=j}(X)\|_2^2.$$

Thus from Proposition 8 on page 40, the information k -means loss is such that

$$\begin{aligned} \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \mathbb{P}_X \left(\mathcal{K}(q_{\ell(X)}^{\star,\ell}, p_X) \right) \\ &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \frac{1}{2\sigma^2} \mathbb{P}_X \left(\|X - \mathbb{P}_{X|\ell(X)}(X)\|_2^2 \right). \end{aligned}$$

Besides,

$$\begin{aligned} \inf_{\mu_1, \dots, \mu_k \in \mathbb{R}^{p \times k}} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|_2^2 \right) &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \inf_{\mu_1, \dots, \mu_k \in \mathbb{R}^{p \times k}} \mathbb{P}_X \left(\|X - \mu_{\ell(X)}\|_2^2 \right) \\ &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \inf_{\mu_1, \dots, \mu_k \in \mathbb{R}^{p \times k}} \left\{ \mathbb{P}_X \left(\|X - \mathbb{P}_{X|\ell(X)}(X)\|_2^2 \right) + \mathbb{P}_X \left(\|\mathbb{P}_{X|\ell(X)}(X) - \mu_{\ell(X)}\|_2^2 \right) \right\} \\ &= \inf_{\ell: \mathcal{X} \rightarrow \llbracket 1, k \rrbracket} \mathbb{P}_X \left(\|X - \mathbb{P}_{X|\ell(X)}(X)\|_2^2 \right). \end{aligned}$$

□

We are now going to linearize the information k -means algorithm, using the kernel trick. Let us introduce the separable Hilbert space $H = \{(f, x) \in \mathbb{L}^2(\nu) \times \mathbb{R}\}$ equipped with the inner-product

$$\langle h, h' \rangle = \langle h_1, h'_1 \rangle_{\mathbb{L}^2(\nu)} + \mu h_2 h'_2, \quad h = (h_1, h_2), h' = (h'_1, h'_2) \in H,$$

where $\mu > 0$ is a positive real parameter to be chosen afterwards. The associated norm is

$$\|(h_1, h_2)\| = \sqrt{\langle h_1, h_1 \rangle_{\mathbb{L}^2(\nu)} + \mu h_2^2} = \sqrt{\int h_1^2 d\nu + \mu h_2^2}, \quad h = (h_1, h_2) \in H.$$

Define for any constant $B \in \mathbb{R}_+$

$$\Theta_B = \left\{ (q, \mathcal{K}(q, 1)) : q \in \mathbb{L}_{+,1}^1(\nu) \cap \mathbb{L}^2(\nu), \int q^2 d\nu \leq B^2 \right\} \subset H,$$

this definition being justified by the fact that

$$\mathcal{K}(q, 1) = \int q \log(q) d\nu \leq \log \left(\int q^2 d\nu \right) < +\infty$$

whenever $\int q^2 d\nu < +\infty$.

LEMMA 12 *Assume that $\text{ess sup}_X \int \log(p_X)^2 d\nu < \infty$ and $\text{ess sup}_X \int p_X^2 d\nu < \infty$. Remark first that the smallest information ball containing the support of \mathbb{P}_{p_X} satisfies*

$$\begin{aligned} \inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X) &\leq \text{ess sup}_X \mathcal{K}(1, p_X) \\ &= \text{ess sup}_X \int \log(p_X^{-1}) d\nu \leq \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right)^{1/2} < \infty. \end{aligned}$$

Define $B = \text{ess sup}_X \left(\int p_X^2 d\nu \right)^{1/2} \exp \left[\inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X) \right] < \infty$ and consider the random variable

$$W = (-\log(p_X), \mu^{-1}) \in H.$$

The following two minimization problems are equivalent

$$\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) = \inf_{\theta \in \Theta_B^k} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle_H \right).$$

PROOF. Let $B' = \text{ess sup}_X \left(\int p_X^2 d\nu \right)^{1/2}$ and $C = \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right)^{1/2}$. First let us remark that under the hypothesis of the lemma, the information k -means criterion is finite. Indeed,

$$\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) \leq \mathbb{P}_X \left(\mathcal{K}(1, p_X) \right) = \mathbb{P}_X \left(\int \log(p_X^{-1}) d\nu \right) \leq C < \infty.$$

Now, for any classification function $\ell : \mathcal{X} \mapsto \llbracket 1, k \rrbracket$ for which the criterion is finite, we know from Lemma 10 on page 41 that $q_j^{\star, \ell} \in \mathbb{L}^2(\nu)$ and we can remark that

$$\mathcal{K}(q_j^{\star, \ell}, p_X) = \langle \theta_j^{\star, \ell}, W \rangle_H$$

where $\theta_j^{\star, \ell} = (q_j^{\star, \ell}, \mathcal{K}(q_j^{\star, \ell}, 1))$. So, it is sufficient to conclude the proof to show that $\theta_j^{\star, \ell} \in \Theta_B$. As in the proof of Lemma 10 on page 41,

$$\int (q_j^{\star, \ell})^2 d\nu \leq Z_j^{-2} \mathbb{P}_{X \mid \ell(X)=j} \left(\underbrace{\int p_X^2 d\nu}_{\leq B'^2} \right) \leq Z_j^{-2} B'^2, \quad j \in \llbracket 1, k \rrbracket.$$

By Jensen's inequality, for any $j \in \llbracket 1, k \rrbracket$,

$$\begin{aligned} Z_j &= \sup_{q \in \mathbb{L}_{+,1}^1(\nu)} \int q \exp \left\{ \mathbb{P}_{X \mid \ell(X)=j} [\log(p_X/q)] \right\} d\nu \\ &\geq \sup_{q \in \mathbb{L}_{+,1}^1(\nu)} \exp \left\{ \mathbb{P}_{X \mid \ell(X)=j} \left[\int q \log(p_X/q) \right] \right\} \\ &= \exp \left\{ - \inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \mathbb{P}_{X \mid \ell(X)=j} [\mathcal{K}(q, p_X)] \right\}. \end{aligned}$$

Hence

$$Z_j^{-1} \leq \exp \left\{ \inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X) \right\} \leq \exp(C).$$

Therefore

$$\left(\int (q_j^{\star, \ell})^2 d\nu \right)^{1/2} \leq B' \exp \left[\inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X) \right] = B \leq B' \exp(C) < \infty,$$

proving that $B < \infty$ and that $\theta_j^{\star, \ell} = (q_j^{\star, \ell}, \mathcal{K}(q_j^{\star, \ell}, 1)) \in \Theta_B$, which concludes the proof. \square

PROPOSITION 13 *Under the hypotheses of the previous lemma there exists an optimal quantizer $\theta^* \in \Theta_B^k$ minimizing the k -means risk, that is such that*

$$\mathbb{E} \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right) = \inf_{\theta \in \Theta_B^k} \mathbb{E} \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right).$$

PROOF. This closely follows the proof of Theorem 3.2 in [Fis10]. Indeed, let us first remark that the linear form $\theta_j \mapsto \langle \theta_j, W \rangle$ is weakly continuous, considering H equipped with its weak topology. It implies that $\theta = (\theta_1, \dots, \theta_k) \mapsto \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle$ is weakly continuous since a finite minimum of continuous function is a continuous function. Moreover, for any $\theta' \in \Theta_B \subset H$, by the Cauchy-Schwarz inequality

$$\begin{aligned} \langle \theta', W \rangle &\leq \|\theta'\| \|W\| \\ &\leq \|\Theta_B\| \|W\|_\infty < \infty, \end{aligned}$$

where $\|W\|_\infty = \text{ess sup}_X \|W\|$ and $\|\Theta_B\| = \sup\{\|\theta\| : \theta \in \Theta_B\}$. Let $\{\theta_n\}_{n \in \mathbb{N}}$ be a sequence in Θ_B^k converging weakly to $\theta \in \Theta_B^k$. Then, by the dominated convergence theorem

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_n^j, W \rangle \right) &= \mathbb{P}_W \left(\lim_{n \rightarrow \infty} \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_n^j, W \rangle \right) \\ &= \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right). \end{aligned}$$

Therefore, $\theta = (\theta_1, \dots, \theta_k) \mapsto \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right)$ is weakly continuous on the weakly compact set Θ_B^k , which implies that it reaches its minimum. Note that we use the weak topology since for the strong topology (associated with the Hilbert norm of H) the closed ball Θ_B is not compact in the case when H is of infinite dimension. \square

3.2. INFORMATION FRAGMENTATION

We are now going to set information k -means into a broader context. This will lead us to propose modified criteria and more general representations, for bag of words, and more generally for random signals, observed through a statistical sample.

We consider as in the previous sections a couple of random variables $(X, Y) \in \mathfrak{X} \times \mathfrak{Y}$ such that $\mathbb{P}(\mathbb{P}_{Y|X} \ll \nu) = 1$ for some dominating measure $\nu \in \mathcal{M}_+^1(\mathfrak{Y})$. We assume that \mathfrak{X} and \mathfrak{Y} are Polish spaces, so that regular conditional probability measures exist.

3.2.1. RECALL OF THE INFORMATION k -MEANS SETTING. In information k -means, we were trying to find a random variable $W = \ell(X) \in \llbracket 1, k \rrbracket$ and a conditional probability measure $Q_{Y|W}$ minimizing

$$\mathbb{P}_{X,W} \left[\mathfrak{K}(Q_{Y|W}, \mathbb{P}_{Y|X}) \right].$$

3.2.2. FIRST GENERALIZATION: ESTIMATING A JOINT DISTRIBUTION. Instead of estimating the conditional probability measure $\mathbb{P}_{Y|X}$, we may be willing to estimate the joint distribution $\mathbb{P}_{X,Y}$ by some simpler measure $Q_{X,Y}$. If we are willing to use the same simplification as in information k -means, we are led to assume that

$$Q_{Y|X} = Q_{Y|W}, \tag{3.3}$$

where $W = \ell(X) \in \llbracket 1, k \rrbracket$ is a measurable function of X . Meanwhile, the first marginal Q_X can be left unconstrained. Note that condition (3.3) implies that

$$Q_{X,Y|W} = Q_{X|W} Q_{Y|W},$$

since $Q_{X,Y|W} = Q_{X|W} Q_{Y|X,W}$ and $Q_{Y|X,W} = Q_{Y|X} = Q_{Y|W}$ from (3.3).

If we still want to use an information criterion, we are led to consider the minimization of

$$\mathfrak{K}(Q_{X,Y}, \mathbb{P}_{X,Y}).$$

From Lemma 1 on page 5, it decomposes into

$$\begin{aligned}\mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) &= \mathcal{K}(Q_X, \mathbb{P}_X) + Q_X[\mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X})] \\ &= \mathcal{K}(Q_X, \mathbb{P}_X) + Q_{X,W}[\mathcal{K}(Q_{Y|W}, \mathbb{P}_{Y|X})].\end{aligned}$$

Remark first that if, instead of leaving Q_X unconstrained, we let $Q_X = \mathbb{P}_X$, we get that $Q_{X,W} = \mathbb{P}_{X,W}$ (since $W = \ell(X)$), and we fall back on the information k -means problem.

Let us see now what happens if we let Q_X be free and use the optimal value of this distribution. From Lemma 6 on page 37, we obtain

$$\begin{aligned}\inf_{Q_X} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) &= \inf_{Q_X} \left\{ \mathcal{K}(Q_X, \mathbb{P}_X) + Q_X[\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X})] \right\} \\ &= -\log \left\{ \mathbb{P}_X \left[\exp(-\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X})) \right] \right\},\end{aligned}$$

where the optimal value of Q_X is absolutely continuous with respect to \mathbb{P}_X , with density

$$\frac{dQ_X}{d\mathbb{P}_X} = Z^{-1} \exp \left[-\mathcal{K}(Q_{Y|\ell(X)}, \mathbb{P}_{Y|X}) \right]. \quad (3.4)$$

PROPOSITION 14 *Consider the modified k -means criterion*

$$\mathcal{C}(\ell : \mathcal{X} \rightarrow \llbracket 1, k \rrbracket, \mu_j \in \mathcal{M}_+^1(\mathcal{Y}), 1 \leq j \leq k) = -\log \left\{ \mathbb{P}_X \left[\exp(-\mathcal{K}(\mu_{\ell(X)}, \mathbb{P}_{Y|X})) \right] \right\}.$$

For a given set of k probability measures $\mu_j \in \mathcal{M}_+^1(\mathcal{Y})$, $1 \leq j \leq k$, the optimal classification function ℓ is

$$\ell^*(X) = \arg \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(\mu_j, \mathbb{P}_{Y|X}).$$

For a given classification function $\ell : \mathcal{X} \rightarrow \llbracket 1, k \rrbracket$, we can lower the criterion introducing Q_X defined by its density

$$\frac{dQ_X}{d\mathbb{P}_X} = Z^{-1} \exp \left[-\mathcal{K}(\mu_{\ell(X)}, \mathbb{P}_{Y|X}) \right]$$

and

$$\frac{d\mu'_j}{d\nu} = Z^{-1} \exp \left\{ Q_{X|\ell(X)=j} \left[\log \left(\frac{d\mathbb{P}_{Y|X}}{d\nu} \right) \right] \right\}, \quad 1 \leq j \leq k.$$

We obtain that

$$\mathcal{C}(\ell, \mu'_j, 1 \leq j \leq k) = \mathcal{C}(\ell, \mu_j, 1 \leq j \leq k) - Q_X \left[\mathcal{K}(\mu_{\ell(X)}, \mu'_{\ell(X)}) \right].$$

Thus we have a descent algorithm that can reach a local minimum of the criterion and can play the role that Lloyd's algorithm plays for the previous information k -means criterion.

3.2.3. INFORMATION FRAGMENTATION. So far, we have made the hypotheses that $W = \ell(X)$ and that

$$Q_{Y|X} = Q_{Y|W}, \quad (3.5)$$

meaning that under the probability Q the conditional probability measures $Q_{Y|X=x}$ are equal for all $x \in \ell^{-1}(j)$. In this setting, Q is a good approximation of \mathbb{P} when $Q_{Y|W=j}$ is a good approximation of $\mathbb{P}_{Y|X=x}$ for each $x \in \ell^{-1}(j)$.

Moreover, hypothesis (3.5) can equivalently be written in the two following forms:

$$\begin{aligned} Q_{Y|X,W} &= Q_{Y|W}, \\ \text{or equivalently} \quad Q_{X,Y|W} &= Q_{X|W} Q_{Y|W}. \end{aligned} \quad (3.6)$$

These two last formulations are equivalent without any hypothesis on W , since it is always the case that $Q_{X,Y|W} = Q_{X|W} Q_{Y|X,W}$. On the other hand they may not be equivalent to (3.5) when W is not assumed to be a measurable function of X .

In the current section, we will relax the constraint on W to $W = \ell(X, Y) \in \llbracket 1, k \rrbracket$, in other words we will assume that W is a measurable function of (X, Y) , or that $\sigma(W) \subset \sigma(X, Y)$. This means that, instead of giving a label W to each bag of words X , we now give labels to each word Y of each bag of words X . In the mean time, we will keep hypothesis (3.6) on Q .

In this new setting,

$$Q_{Y|X} = Q_{W|X}(Q_{Y|X,W}) = Q_{W|X}(Q_{Y|W}).$$

This means that under Q the conditional probability measure $Q_{Y|X}$ describing the content of the signal X is a mixture of k *fragments* $Q_{Y|W=j}$, $1 \leq j \leq k$. We see therefore that the power of approximation of this new model is greater than the previous one. While in the previous model each bag of words $\mathbb{P}_{Y|X}$ had to be close to a model $Q_{Y|W}$ taking only k possible values, in the new model each bag of words $\mathbb{P}_{Y|X}$ has to be close to a linear combination of k fragments $Q_{Y|W}$.

Another important interpretation of the new model comes from the identity

$$Q_{X,Y} = Q_W(Q_{X|W} \otimes Q_{Y|W}).$$

It shows that we are looking for an approximation of $\mathbb{P}_{X,Y}$ by a mixture of k product distributions.

If we were to achieve this with the usual EM algorithm for mixture estimation, we would try to minimize

$$\inf_{W,Q} \mathcal{K}(\mathbb{P}_{X,Y}, Q_{X,Y})$$

(or rather to decrease this criterion iteratively). This would fit into the usual framework of statistical inference that is to find a model that can predict with the highest possible probability all observed data configurations.

Here, in connection with the information k -means algorithm, we raise the question of minimizing the reverse divergence

$$\inf_{W, Q} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}).$$

In this framework, instead of looking for a model that can predict all the observations, we look for a model whose predictions can all be observed with the highest possible probability. This is closer to what is held for a valid theory in experimental sciences. In physics, for instance, a model is considered to be valid if all its predictions can be observed. On the other hand, a model that can predict all observations at the price of also predicting events that cannot be observed would be considered as false or not relevant.

Let us now state the equivalent of Proposition 14 on page 46, that is let us describe the generalization of Lloyd's algorithm to information fragmentation.

PROPOSITION 15 *Consider k centers $\mu_j \in \mathcal{M}_+^1(\mathcal{Y})$, $1 \leq j \leq k$. Let*

$$\mathcal{T} = \left\{ A \subset \llbracket 1, k \rrbracket : \mu_i \perp \mu_j, i \neq j \in A \right\}$$

be the set of tilings by mutually singular probability measures μ_j . The partial minimum

$$\inf_{Q_{X,Y}} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y})$$

taken on all probability measures $Q_{X,Y}$, such that for some measurable function $\ell : \mathcal{X} \times \mathcal{Y} \rightarrow \llbracket 1, k \rrbracket$,

$$Q_{X,Y} \left[Q_{Y|X, \ell(X, Y)} = \mu_{\ell(X, Y)} \right] = 1 \quad (3.7)$$

is equal to

$$\begin{aligned} & - \sup_{\ell} \log \left\{ \mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_j \left(\ell_X^{-1}(j) \right) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right\} \\ & = - \log \mathbb{P}_X \left(\sup_{A \in \mathcal{T}} \left\{ \sum_{j \in A} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right\} \right), \end{aligned}$$

where

$$\begin{aligned} \ell_x : \mathcal{Y} & \rightarrow \llbracket 1, k \rrbracket \\ y & \mapsto \ell(x, y). \end{aligned}$$

For any given choice of ℓ , putting $W = \ell(X, Y)$, the optimum in $Q_{W,X}$ is reached when

$$\frac{dQ_{W,X}}{d\mathbb{P}_{W,X}} = Z^{-1} \exp \left[-\mathcal{K}(\mu_W, \mathbb{P}_{Y|W,X}) \right]. \quad (3.8)$$

Symmetric formulas apply when we exchange the role of X and Y . They describe the optimization on all probability measures $Q_{X,Y}$ such that

$$Q_{X,Y} \left[Q_{X|Y, \ell(X, Y)} = \rho_{\ell(X, Y)} \right] = 1,$$

where $\rho_j \in \mathcal{M}_+^1(\mathcal{X})$, $1 \leq j \leq k$ are given. Note that in this case, the optimal value of $Q_{W,Y}$ is given by

$$\frac{dQ_{W,Y}}{d\mathbb{P}_{W,Y}} = Z^{-1} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,Y}) \right],$$

and that we can use the identity

$$\mathcal{K}(\rho_W, \mathbb{P}_{X|W,Y}) = \mathcal{K}(\rho_W, \mathbb{P}_{X|W}) - \rho_W \left[\log \left(\frac{d\mathbb{P}_{Y|W,X}}{d\mathbb{P}_{Y|W}} \right) \right], \quad (3.9)$$

to avoid computing $\mathbb{P}_{X|Y,W}$ explicitly.

Iterating these two optimization steps, we reach a local minimum for the optimization on all probability measures $Q_{X,Y}$, such that for some measurable function $\ell : \mathcal{X} \times \mathcal{Y} \rightarrow \llbracket 1, k \rrbracket$

$$Q_{X,Y} \left[Q_{X,Y|\ell(X,Y)} = Q_{X|\ell(X,Y)} \otimes Q_{Y|\ell(X,Y)} \right] = 1.$$

PROOF. Assume for the moment that the classification function ℓ is fixed. Note that

$$Q_{Y|X, \ell(X,Y)=j} = Q_{Y|X, Y \in \ell_X^{-1}(j)}.$$

Let us set

$$Q_{Y|X, Y \in \ell_X^{-1}(j)} = \mu_j,$$

whenever $\mu_j[\ell_X^{-1}(j)] = 1$, and give it any arbitrary value otherwise. Let us set $Q_{W,X}$ as described in equation (3.8). This defines $Q_{X,Y}$. Remark that condition (3.7) is satisfied. Indeed

$$\frac{dQ_{W|X}}{d\mathbb{P}_{W|X}}(j) = Z_X^{-1} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X, W=j}) \right],$$

so that when $\mu_j[\ell_X^{-1}(j)] < 1$, $\mathcal{K}(\mu_j, \mathbb{P}_{Y|X, W=j}) = \mathcal{K}(\mu_j, \mathbb{P}_{Y|X, Y \in \ell_X^{-1}(j)}) = +\infty$ and therefore $Q_{W|X}(j) = Q(\ell(X,Y) = j | X) = 0$. Thus

$$Q_{Y|X} \left[\ell(X,Y) \in \left\{ j : \mu_j[\ell_X^{-1}(j)] = 1 \right\} \right] = 1,$$

implying condition (3.7) according to the construction of $Q_{Y|X, \ell(X,Y)=j}$.

From the decomposition formula for the Kullback divergence (Lemma 1 on page 5), any probability measure $Q'_{X,Y}$ satisfying condition (3.7) is such that

$$\mathcal{K}(Q'_{X,Y}, \mathbb{P}_{X,Y}) = \mathcal{K}(Q'_{X,Y,W}, \mathbb{P}_{X,Y,W}) = \mathcal{K}(Q'_{X,W}, \mathbb{P}_{X,W}) + Q'_{X,W} \left[\mathcal{K}(\mu_W, \mathbb{P}_{Y|X,W}) \right].$$

Therefore, according to Lemma 6,

$$\inf_{Q'_{X,W}} \mathcal{K}(Q'_{X,Y}, \mathbb{P}_{X,Y}) = \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) = -\log \left\{ \mathbb{P}_{X,W} \left[\exp \left(-\mathcal{K}(\mu_W, \mathbb{P}_{Y|X,W}) \right) \right] \right\}.$$

Remark that

$$\frac{d\mathbb{P}_{Y|X,W}}{d\mathbb{P}_{Y|X}} = \frac{d\mathbb{P}_{W|X,Y}}{d\mathbb{P}_{W|X}},$$

so that

$$\begin{aligned}\mathcal{K}(\mu_W, \mathbb{P}_{Y|X,W}) &= \mathcal{K}(\mu_W, \mathbb{P}_{Y|X}) - \mu_W \left[\log \left(\frac{d\mathbb{P}_{Y|X,W}}{d\mathbb{P}_{Y|X}} \right) \right] \\ &= \mathcal{K}(\mu_W, \mathbb{P}_{Y|X}) - \mu_W \left[\log \left(\frac{d\mathbb{P}_{W|X,Y}}{d\mathbb{P}_{W|X}} \right) \right].\end{aligned}$$

Accordingly,

$$\begin{aligned}\mathbb{P}_{W|X} \left[\exp \left(-\mathcal{K}(\mu_W, \mathbb{P}_{Y|X,W}) \right) \right] &= \mathbb{P}_{W|X} \left\{ \exp \left\{ -\mathcal{K}(\mu_W, \mathbb{P}_{Y|X}) + \mu_W \left[\log \left(\frac{d\mathbb{P}_{W|X,Y}}{d\mathbb{P}_{W|X}} \right) \right] \right\} \right\} \\ &= \sum_{j=1}^k \mathbb{1} \left[\mathbb{P}_{Y|X}(\ell_X^{-1}(j)) > 0 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) + \mu_j [\log(\mathbb{1}(\ell(X, Y) = j))] \right] \\ &= \sum_{j=1}^k \mathbb{1} \left[\mathbb{P}_{Y|X}(\ell_X^{-1}(j)) > 0 \right] \mathbb{1} \left[\mu_j(\ell_X^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \\ &= \sum_{j=1}^k \mathbb{1} \left[\mu_j(\ell_X^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right],\end{aligned}$$

because when $\mu_j(\ell_X^{-1}(j)) = 1$ and $\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) < +\infty$, necessarily $\mu_j \ll \mathbb{P}_{Y|X}$, so that $\mathbb{P}_{Y|X}(\ell_X^{-1}(j)) > 0$. Therefore

$$\begin{aligned}\inf \left\{ \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) : Q_{X,Y} \text{ satisfies (3.7) for some } \ell \right\} &= - \sup_{\ell} \log \left[\mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_j(\ell_X^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right].\end{aligned}$$

For a given classification function ℓ , consider

$$A_X(\ell) = \left\{ j : \mu_j(\ell_X^{-1}(j)) = 1 \right\}.$$

The measures $\{\mu_j, j \in A_X(\ell)\}$ being concentrated on disjoint sets, they are by definition mutually singular, so that $A_X(\ell) \in \mathcal{T}$. Therefore

$$\begin{aligned}\sup_{\ell} \log \left[\mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_j(\ell_X^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right] &= \sup_{\ell} \log \left[\mathbb{P}_X \left(\sum_{j \in A_X(\ell)} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right] \\ &\leq \log \left[\mathbb{P}_X \left(\sup_{A \in \mathcal{T}} \left\{ \sum_{j \in A} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right\} \right) \right].\end{aligned}$$

On the other hand, when the measures $\{\mu_j, j \in A\}$ are mutually singular, that is when $A \in \mathcal{T}$, we can find disjoint measurable sets $\{B_j(A) \subset \mathcal{Y}, j \in A\}$, such that $\mu_j(B_j(A)) =$

$1, j \in A$, and therefore we can find a measurable function $\ell_A : \mathbb{Y} \rightarrow \llbracket 1, k \rrbracket$ such that $\mu_j(\ell_A^{-1}(j)) = 1$, for any $j \in A$. Let

$$A_x = \arg \max_{A \in \mathcal{T}} \sum_{j \in A} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X=x}) \right], \quad x \in \mathfrak{X}.$$

Let us define ℓ^* by the formula $\ell_x^* = \ell_{A_x}$, $x \in \mathfrak{X}$. Since

$$x \mapsto \sum_{j \in A} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X=x}) \right]$$

is measurable, $x \mapsto A_x$ is measurable and therefore $x \mapsto \ell_{A_x}$ is measurable, taking its values in a finite set of measurable functions, implying that ℓ^* is measurable. Remark that

$$\begin{aligned} \log \left[\mathbb{P}_X \left(\sup_{A \in \mathcal{T}} \left\{ \sum_{j \in A} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right\} \right) \right] &= \log \left[\mathbb{P}_X \left(\sum_{j \in A_X} \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right] \\ &\leq \log \left[\mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_j(\ell_{A_X}^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right] \\ &= \log \left[\mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_j(\ell_X^*{}^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right] \\ &\leq \sup_{\ell} \log \left[\mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_j(\ell_X^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right] \right) \right]. \end{aligned}$$

This proves that

$$\begin{aligned} \inf \left\{ \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) : Q_{X,Y} \text{ satisfies (3.7) for some } \ell \right\} \\ = -\log \left[\mathbb{P}_X \left(\sup_{A \in \mathcal{T}} \left\{ \sum_{j \in A} \mathcal{K}(\mu_j, \mathbb{P}_{Y|X}) \right\} \right) \right]. \end{aligned}$$

To prove identity (3.9), it is enough to remark that

$$\begin{aligned} \mathcal{K}(\rho_W, \mathbb{P}_{X|Y,W}) &= \rho_W \left[\log \left(\frac{d\rho_W}{d\mathbb{P}_{X|Y,W}} \right) \right] \\ &= \rho_W \left[\log \left(\frac{d\rho_W}{d\mathbb{P}_{X|W}} \right) \right] - \rho_W \left[\log \left(\frac{d\mathbb{P}_{X|Y,W}}{d\mathbb{P}_{X|W}} \right) \right] \\ &= \rho_W \left[\log \left(\frac{d\rho_W}{d\mathbb{P}_{X|W}} \right) \right] - \rho_W \left[\log \left(\frac{d\mathbb{P}_{Y|X,W}}{d\mathbb{P}_{Y|W}} \right) \right]. \end{aligned}$$

□

3.3. SIGNAL FRAGMENTATION

Consider a random signal $X : \Omega \rightarrow \mathbb{R}^d$. Let us assume that $S : \Omega \rightarrow \llbracket 1, d \rrbracket$ and $V : \Omega \rightarrow \mathbb{R}$ are such that

$$\mathbb{P}_{S,V|X} = \mathbb{P}_{S|X} \mathcal{N}(X_S, \sigma^2),$$

where $\text{supp}(\mathbb{P}_{S|X}) = [\![1, d]\!]$ and where σ is a positive real parameter. Note that X is a function of $\mathbb{P}_{S,V|X}$, given by the identity

$$X_s = \mathbb{P}_{V|X, S=s}(V), \quad 1 \leq s \leq d.$$

In other words, $\mathbb{P}_{S,V|X}$ is a lossless representation of the signal X . This representation is not unique, since we can choose the smoothing parameter σ and the site distribution $\mathbb{P}_{S|X}$ as we please to represent the same X . Usually we will take $\mathbb{P}_{S|X}$ to be the uniform probability measure on $[\![1, d]\!]$, this choice being independent from X .

We are looking for labels $W = \ell(X, S) \in [\![1, k]\!]$ and for a probability measure $Q \in \mathcal{M}_+^1(\Omega)$ such that $Q_{X, S, V|W} = Q_{X|W} Q_{S, V|W}$, Q_W almost surely, minimizing

$$\mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V}).$$

The process $Q_{X, S, V}$ can be understood as a patch process approximating $\mathbb{P}_{X, S, V}$. Indeed,

$$Q_{S, V|X} = Q_{W|X}(Q_{S, V|W}),$$

meaning that the signal $Q_{S, V|X}$ is a mixture of the k patches $Q_{S, V|W=j}$, $1 \leq j \leq k$.

PROPOSITION 16 (GENERALIZED k -MEANS ALGORITHM) *Consider k centers $\rho_j \in \mathcal{M}_+^1(\mathbb{R}^d)$, $1 \leq j \leq k$. Define*

$$\mathfrak{T}_2 = \left\{ B \subset [\![1, k]\!] : \rho_i \perp \rho_j, i \neq j \in B \right\},$$

the set of tilings by mutually singular probability measures ρ_j . The partial minimum

$$\inf_Q \mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V})$$

taken on all probability measures Q , such that for some measurable function $\ell : \mathbb{R}^d \times [\![1, d]\!] \rightarrow [\![1, k]\!]$

$$Q \left[Q_{X|S, V, \ell(X, S)} = \rho_{\ell(X, S)} \right] = 1. \quad (3.10)$$

is equal to

$$\begin{aligned} - \sup_{\ell} \log \mathbb{P}_S \left(\sum_{j=1}^k \mathbb{1} \left[\rho_j(\ell_S^{-1}(j)) = 1 \right] \exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \text{Var}_{\rho_j}(X_S)/(2\sigma^2) \right] \right) \\ = - \log \mathbb{P}_S \left(\sup_{B \in \mathfrak{T}_2} \sum_{j \in B} \exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \text{Var}_{\rho_j}(X_S)/(2\sigma^2) \right] \right), \end{aligned}$$

where

$$\begin{aligned} \ell_s : \mathbb{R}^d &\rightarrow [\![1, k]\!] \\ x &\mapsto \ell(x, s). \end{aligned}$$

For any choice of ℓ , and in particular for the optimal one, putting $W = \ell(X, S)$, the optimum in $Q_{W, S, V}$ is reached when

$$\frac{dQ_{W, S, V}}{d\mathbb{P}_{W, S} \otimes \lambda_V} = Z^{-1} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W, S}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2) \right] g_{\sigma, \rho_W(X_S)}(V), \quad (3.11)$$

where λ_V is the Lebesgue measure on \mathbb{R} and

$$g_{\sigma, m}(v) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(v-m)^2}{2\sigma^2}\right).$$

In particular, for the optimal choice of $Q_{W, S, V}$, $Q_{V|S, W} = \mathcal{N}(\rho_W(X_S), \sigma^2)$ is a Gaussian probability measure and

$$\frac{dQ_{S|W}}{d\mathbb{P}_{S|W}} = Z_W^{-1} \exp\left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W, S}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2)\right].$$

On the other hand, consider k centers $\mu_{S, V}^{(j)} \in \mathcal{M}_+^1([\![1, d]\!] \times \mathbb{R})$, $1 \leq j \leq k$ such that

$$\mu_{V|S}^{(j)} = \mathcal{N}(\mu_{V|S}^{(j)}(V), \sigma^2), \quad 1 \leq j \leq k.$$

Define

$$\mathcal{T}_1 = \left\{ A \subset [\![1, k]\!]: \mu_S^{(i)} \perp \mu_S^{(j)}, i \neq j \in A \right\},$$

the set of tilings by mutually singular probability measures $\mu_S^{(j)}$ (or equivalently by mutually singular probability measures $\mu_{S, V}^{(j)}$).

The partial minimum

$$\inf_Q \mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V})$$

taken on all probability measures $Q \in \mathcal{M}_+^1(\Omega)$ such that, for some measurable function $\ell: \mathbb{R}^d \times [\![1, d]\!] \rightarrow [\![1, k]\!]$

$$Q\left[Q_{S, V|X, \ell(X, S)} = \mu_{S, V}^{(\ell(X, S))}\right] = 1, \quad (3.12)$$

is equal to

$$\begin{aligned} & - \sup_{\ell} \log \mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_S^{(j)} \left(\ell_X^{-1}(j) \right) = 1 \right] \right. \\ & \quad \times \exp \left\{ -\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) - \mu_S^{(j)} \left[\left(\mu_{V|S}^{(j)}(V) - X_S \right)^2 / (2\sigma^2) \right] \right\} \left. \right) \\ & = - \log \mathbb{P}_X \left(\sup_{A \in \mathcal{T}_1} \sum_{j \in A} \exp \left\{ -\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) - \mu_S^{(j)} \left[\left(\mu_{V|S}^{(j)}(V) - X_S \right)^2 / (2\sigma^2) \right] \right\} \right), \end{aligned}$$

where

$$\begin{aligned} \ell_x: [\![1, d]\!] & \rightarrow [\![1, k]\!] \\ s & \mapsto \ell(x, s). \end{aligned}$$

For any value of ℓ , and in particular for the optimal one, considering $W = \ell(X, S)$, the minimum in $Q_{X, W}$ is reached when

$$\frac{dQ_{X, W}}{d\mathbb{P}_{X, W}} = Z^{-1} \exp \left\{ -\mathcal{K}(\mu_S^{(W)}, \mathbb{P}_{S|X, W}) - \mu_S^{(W)} \left[\left(\mu_{V|S}^{(W)}(V) - X_S \right)^2 / (2\sigma^2) \right] \right\}. \quad (3.13)$$

Alternating these two partial optimization steps, we can converge to a local minimum for the optimization problem

$$\inf_Q \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V}),$$

where the infimum is taken over probability measures $Q \in \mathcal{M}_+^1(\Omega)$ satisfying, for some measurable classification function $\ell : \mathbb{R}^d \times \llbracket 1, d \rrbracket \rightarrow \llbracket 1, k \rrbracket$,

$$Q \left[Q_{X,S,V|\ell(X,S)} = Q_{X|\ell(X,S)} \otimes Q_{S,V|\ell(X,S)} \right] = 1. \quad (3.14)$$

PROOF. Assume for the time being that the classification function ℓ is fixed and note that

$$Q_{X|S,V,\ell(X,S)=j} = Q_{X|S,V,X \in \ell_S^{-1}(j)}.$$

Let us set

$$Q_{X|S,V,\ell(X,S)=j} = \rho_j$$

whenever $\rho_j(\ell_S^{-1}(j)) = 1$ and give it any arbitrary value otherwise. Set $W = \ell(X, S)$ and define $Q_{S,V,W}$ by equation (3.11) on page 52. This defines $Q_{X,S,V}$. Condition (3.10) on page 52 is satisfied. Indeed,

$$\frac{dQ_{W|S}}{d\mathbb{P}_{W|S}} = Z_S^{-1} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2) \right].$$

Remark that

$$\mathbb{P}_{X|S,W=j} = \mathbb{P}_{X|S,X \in \ell_S^{-1}(j)},$$

so that $\mathbb{P}_{X|S,W=j}(\ell_S^{-1}(j)) = 1$. Therefore, when $\rho_j(\ell_S^{-1}(j)) < 1$, ρ_j is not absolutely continuous with respect to $\mathbb{P}_{X|S,W=j}$, $\mathcal{K}(\rho_j, \mathbb{P}_{X|S,W=j}) = +\infty$ and $Q_{W|S}(j) = 0 = Q(\ell(X, S) = j | S)$. Thus $\rho_W(\ell_S^{-1}(W)) = 1$, Q almost surely, and therefore $Q_{X|S,V,\ell(X,S)} = \rho_{\ell(X,S)}$, Q almost surely, as required by condition (3.10) on page 52.

Let us prove that

$$\frac{dQ_{S,V,W}}{d\mathbb{P}_{S,V,W}} = Z^{-1} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,V,W}) \right]. \quad (3.15)$$

Remark that

$$\begin{aligned} \mathcal{K}(\rho_W, \mathbb{P}_{X|W,S,V}) &= \mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \rho_W \left[\log \left(\frac{d\mathbb{P}_{X|W,S,V}}{d\mathbb{P}_{X|W,S}} \right) \right] \\ &= \mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \rho_W \left[\log \left(\frac{d\mathbb{P}_{V|X,W,S}}{d\mathbb{P}_{V|W,S}} \right) \right] \\ &= \mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \rho_W \left[\log \left(\frac{d\mathbb{P}_{V|X,S}}{d\mathbb{P}_{V|W,S}} \right) \right], \quad (\text{since } W = \ell(X, S)). \end{aligned}$$

Hence

$$\exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,V,W}) \right]$$

$$\begin{aligned}
&= \exp \left\{ -\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) + \rho_W \left[\log \left(\frac{d\mathbb{P}_{V|X,S}}{d\lambda_V} \right) \right] \right\} \frac{d\lambda_V}{d\mathbb{P}_{V|W,S}} \\
&= \frac{d\lambda_V}{d\mathbb{P}_{V|W,S}} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \rho_W [(V - X_S)^2]/(2\sigma^2) - \log(\sigma\sqrt{2\pi}) \right] \\
&= \frac{d\lambda_V}{d\mathbb{P}_{V|W,S}} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2) \right] \times \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{(V - \rho_W(X_S))^2}{2\sigma^2} \right] \\
&= \frac{d\lambda_V}{d\mathbb{P}_{V|W,S}} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|W,S}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2) \right] g_{\sigma, \rho_W(X_S)}(V). \quad (3.16)
\end{aligned}$$

Thus, from the definition of $Q_{S,V,W}$,

$$\frac{dQ_{W,S,V}}{d\mathbb{P}_{W,S} \otimes \lambda_V} = Z^{-1} \frac{d\mathbb{P}_{V|W,S}}{d\lambda_V} \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,V,W}) \right],$$

proving (3.15).

According to the decomposition formula for the divergence (Lemma 1 on page 5), for any probability measure $Q'_{X,S,V}$ satisfying condition (3.10) on page 52, for the same classification function ℓ as Q ,

$$\mathcal{K}(Q'_{S,V,X}, \mathbb{P}_{S,V,X}) = \mathcal{K}(Q'_{S,V,W}, \mathbb{P}_{S,V,W}) + Q'_{S,V,W} \left[\mathcal{K}(\rho_W, \mathbb{P}_{X|S,V,W}) \right],$$

so that from Lemma 6 on page 37 and from (3.15),

$$\inf_{Q'_{S,V,W}} \mathcal{K}(Q'_{S,V,X}, \mathbb{P}_{S,V,X}) = -\log \left\{ \mathbb{P}_{S,V,W} \left[\exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,V,W}) \right] \right] \right\}$$

is reached when $Q'_{X,S,V} = Q_{X,S,V}$. Moreover, from (3.16),

$$\mathbb{P}_{V|S,W} \left[\exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,V,W}) \right] \right] = \exp \left[-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,W}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2) \right].$$

Thus

$$\begin{aligned}
\inf_{Q'_{S,V,X}} \mathcal{K}(Q'_{S,V,X}, \mathbb{P}_{S,V,X}) &= -\log \mathbb{P}_{S,W} \left[\exp \left(-\mathcal{K}(\rho_W, \mathbb{P}_{X|S,W}) - \text{Var}_{\rho_W}(X_S)/(2\sigma^2) \right) \right] \\
&= -\log \mathbb{P}_S \left(\sum_{j=1}^k \mathbb{P}_{W|S}(j) \exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S,W=j}) - \text{Var}_{\rho_j}(X_S)/(2\sigma^2) \right] \right).
\end{aligned}$$

Remark now that whenever $\mathbb{P}_{W|S}(j) > 0$,

$$\begin{aligned}
\mathcal{K}(\rho_j, \mathbb{P}_{X|S,W=j}) &= \mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \rho_j \left[\log \left(\frac{d\mathbb{P}_{X|S,W=j}}{d\mathbb{P}_{X|S}} \right) \right] \\
&= \mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \rho_j \left[\log \left(\frac{\mathbb{P}_{W|X,S}(j)}{\mathbb{P}_{W|S}(j)} \right) \right],
\end{aligned}$$

so that

$$\exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S,W=j}) \right] = \frac{\mathbb{1}[\rho_j(\ell_S^{-1}(j)) = 1]}{\mathbb{P}_{W|S}(j)} \exp \left[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) \right]$$

Therefore,

$$\begin{aligned}
& \inf_{Q'_{S,V,X}} \mathcal{K}(Q'_{S,V,X}, \mathbb{P}_{S,V,X}) \\
&= -\log \mathbb{P}_S \left(\sum_{j=1}^k \mathbb{1}[\mathbb{P}_{W|S}(j) > 0] \mathbb{1}[\rho_j(\ell_S^{-1}(j)) = 1] \right. \\
&\quad \left. \times \exp[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \text{Var}_{\rho_j}(X_S)/(2\sigma^2)] \right) \\
&= -\log \mathbb{P}_S \left(\sum_{j=1}^k \mathbb{1}[\rho_j(\ell_S^{-1}(j)) = 1] \exp[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \text{Var}_{\rho_j}(X_S)/(2\sigma^2)] \right),
\end{aligned}$$

because $\mathbb{P}_{W|S}(j) = \mathbb{P}_{X|S}(\ell_S^{-1}(j)) > 0$ when $\rho_j(\ell_S^{-1}(j)) = 1$ and $\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) < \infty$ and consequently $\rho_j \ll \mathbb{P}_{X|S}$.

Reasoning as in the proof of Proposition 15 on page 48, we conclude that

$$\begin{aligned}
& \inf_{\ell} \inf_{Q'_{S,V,X}} \mathcal{K}(Q'_{S,V,X}, \mathbb{P}_{S,V,X}) \\
&= -\sup_{\ell} \log \mathbb{P}_S \left(\sum_{j=1}^k \mathbb{1}[\rho_j(\ell_S^{-1}(j)) = 1] \exp[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \text{Var}_{\rho_j}(X_S)/(2\sigma^2)] \right) \\
&= -\log \mathbb{P}_S \left(\sup_{B \in \mathcal{I}_2} \sum_{j \in B} \exp[-\mathcal{K}(\rho_j, \mathbb{P}_{X|S}) - \text{Var}_{\rho_j}(X_S)] \right).
\end{aligned}$$

The proof of the second half of the proposition is in the same spirit. We construct an optimal solution for a given classification function ℓ setting

$$Q_{S,V|X, \ell(X,S)=j} = \mu_{S,V}^{(j)}$$

when $\mu_S^{(j)}(\ell_X^{-1}(j)) = 1$, and any value otherwise. We complete the definition of $Q_{X,S,V}$ defining $Q_{X,W}$ as in equation (3.13) on page 53. We then remark that $\mathbb{P}_{S|X,W=j} = \mathbb{P}_{S|X, S \in \ell_X^{-1}(j)}$, so that when $\mu_S^{(j)}(\ell_X^{-1}(j)) < 1$, $\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X,W=j}) = +\infty$, implying that

$$Q_{X,S} \left[\mu_S^{(\ell(X,S))}(\ell_X^{-1}(\ell(X,S))) = 1 \right] = 1,$$

and therefore condition (3.12) on page 53 is satisfied. Let us prove now that

$$\frac{dQ_{X,W}}{d\mathbb{P}_{X,W}} = Z^{-1} \exp \left[-\mathcal{K}(\mu_{S,V}^{(W)}, \mathbb{P}_{S,V|X,W}) \right]. \quad (3.17)$$

Indeed,

$$\begin{aligned}
\mathcal{K}(\mu_{S,V}^{(W)}, \mathbb{P}_{S,V|X,W}) &= \mathcal{K}(\mu_S^{(W)}, \mathbb{P}_{S|X,W}) + \mu_S^{(W)} \left[\mathcal{K}(\mu_{V|S}^{(W)}, \mathbb{P}_{V|S,X,W}) \right] \\
&= \mathcal{K}(\mu_S^{(W)}, \mathbb{P}_{S|X,W}) + \mu_S^{(W)} \left[\mathcal{K}(\mu_{V|S}^{(W)}, \mathbb{P}_{V|S,X}) \right], \quad (\text{since } W = \ell(X,S)) \\
&= \mathcal{K}(\mu_S^{(W)}, \mathbb{P}_{S|X,W}) + \mu_S^{(W)} \left[(\mu_{V|S}^{(W)}(V) - X_S)^2 / (2\sigma^2) \right].
\end{aligned}$$

From the decomposition property of the divergence,

$$\begin{aligned}
\mathcal{K}(Q'_{X,S,V}, \mathbb{P}_{X,S,V}) &= \mathcal{K}(Q'_{X,S,V,W}, \mathbb{P}_{X,S,V,W}) \\
&= \mathcal{K}(Q'_{X,W}, \mathbb{P}_{X,W}) + Q'_{X,W} \left[\mathcal{K}(Q'_{S,V|X,W}, \mathbb{P}_{S,V|X,W}) \right] \\
&= \mathcal{K}(Q'_{X,W}, \mathbb{P}_{X,W}) + Q'_{X,W} \left[\mathcal{K}(\mu_{S,V}^{(W)}, \mathbb{P}_{S,V|X,W}) \right],
\end{aligned}$$

so that according to equation (3.17) and Lemma 6 on page 37, Q is optimal for ℓ fixed, and

$$\begin{aligned}
\inf_{Q'} \mathcal{K}(Q'_{X,S,V}, \mathbb{P}_{X,S,V}) &= -\log \mathbb{P}_{X,W} \left[\exp \left[-\mathcal{K}(\mu_{S,V}^{(W)}, \mathbb{P}_{S,V|X,W}) \right] \right] \\
&= -\log \mathbb{P}_{X,W} \left[\exp \left\{ -\mathcal{K}(\mu_S^{(W)}, \mathbb{P}_{S|X,W}) - \mu_S^{(W)} \left[(\mu_{V|S}^{(W)}(V) - X_S)^2 / (2\sigma^2) \right] \right\} \right] \\
&= -\log \mathbb{P}_X \left(\sum_{j=1}^k \mathbb{1} \left[\mu_S^{(j)}(\ell_X^{-1}(j)) = 1 \right] \exp \left\{ -\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) - \mu_S^{(j)} \left[(\mu_{V|S}^{(j)}(V) - X_S)^2 / (2\sigma^2) \right] \right\} \right),
\end{aligned}$$

where the last identity is proved as in the case of the first half of the proposition. As in the proof of Proposition 15 on page 48, we conclude that

$$\begin{aligned}
\inf_{\ell} \inf_{Q'} \mathcal{K}(Q'_{X,S,V}, \mathbb{P}_{X,S,V}) \\
&= -\log \mathbb{P}_X \left(\sup_{A \in \mathcal{T}_1} \sum_{j \in A} \exp \left\{ -\mathcal{K}(\mu_S^{(j)}, \mathbb{P}_{S|X}) - \mu_S^{(j)} \left[(\mu_{V|S}^{(j)}(V) - X_S)^2 / (2\sigma^2) \right] \right\} \right).
\end{aligned}$$

□

CHAPTER 4

PAC-Bayesian bounds for information k -means and information fragmentation

4.1. A PAC-BAYESIAN BOUND FOR INFORMATION k -MEANS

In this section, we consider the setting described in Section 3.1 on page 35, and more specifically in Proposition 13 on page 44.

We first derive a non-asymptotic dimension free bound for the representation of the problem in the separable Hilbert space H . Then, we translate this result to the case of the original information k -means risk.

Our proofs are based on the following PAC-Bayesian lemma.

LEMMA 17 *Consider two measurable spaces \mathcal{T} and \mathcal{W} , a prior probability measure $\pi \in \mathcal{M}_+^1(\mathcal{T})$ defined on \mathcal{T} , and a measurable function $h : \mathcal{T} \times \mathcal{W} \rightarrow \mathbb{R}$. Let $W \in \mathcal{W}$ be a random variable and let (W_1, \dots, W_n) be a sample made of n independent copies of W . Let λ be a positive real parameter.*

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \exp \left[\sup_{\rho \in \mathcal{M}_+^1(\mathcal{T})} \sup_{\eta \in \mathbb{N}} \left\{ \int \min \left\{ \eta, -\lambda \sum_{i=1}^n h(\theta', W_i) \right. \right. \right. \right. \right. \\ \left. \left. \left. \left. \left. \left. - n \log \left[\mathbb{P}_W \exp[-\lambda h(\theta', W)] \right] \right\} d\rho(\theta') - \mathcal{K}(\rho, \pi) \right\} \right] \right\} \leq 1. \quad (4.1)$$

Consequently, for any $\delta \in]0, 1[$, with probability at least $1 - \delta$,

$$\sup_{\rho \in \mathcal{M}_+^1(\mathcal{T})} \sup_{\eta \in \mathbb{N}} \left\{ \int \min \left\{ \eta, -\lambda \sum_{i=1}^n h(\theta', W_i) \right. \right. \\ \left. \left. - n \log \left[\mathbb{P}_W \exp[-\lambda h(\theta', W)] \right] \right\} d\rho(\theta') - \mathcal{K}(\rho, \pi) \right\} \leq \log(\delta^{-1}). \quad (4.2)$$

Note that the role of η in this formula is to give a meaning to the integration with respect to ρ in all circumstances.

PROOF. We follow here the same arguments as in the proof of Proposition 1.7 in [Giu15]. Remark that the supremum in ρ can be restricted to the case when $\mathcal{K}(\rho, \pi) < \infty$, and recall that in this case $\rho \ll \pi$ and $\mathcal{K}(\rho, \pi) = \int \log\left(\frac{d\rho}{d\pi}(\theta')\right) d\rho(\theta')$. Note also that

$$\int \mathbb{1}\left(\frac{d\rho}{d\pi}(\theta') > 0\right) d\rho(\theta') = \int \mathbb{1}\left(\frac{d\rho}{d\pi}(\theta') > 0\right) \frac{d\rho}{d\pi}(\theta') d\pi(\theta') = \int \frac{d\rho}{d\pi}(\theta') d\pi(\theta') = 1.$$

Applying Jensen's inequality, we get

$$\begin{aligned} & \exp \left\{ \sup_{\rho \in \mathcal{M}_+^1(\mathcal{T})} \sup_{\eta \in \mathbb{N}} \int \min \left\{ \eta, -\lambda \sum_{i=1}^n h(\theta', W_i) \right. \right. \\ & \quad \left. \left. - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} d\rho(\theta') - \mathcal{K}(\rho, \pi) \right\} \\ & \leq \sup_{\eta \in \mathbb{N}} \sup_{\substack{\rho \in \mathcal{M}_+^1(\mathcal{T}) \\ \mathcal{K}(\rho, \pi) < \infty}} \int \exp \left\{ \min \left\{ \eta, -\lambda \sum_{i=1}^n h(\theta', W_i) \right. \right. \\ & \quad \left. \left. - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} \right\} \frac{d\rho}{d\pi}(\theta')^{-1} d\rho(\theta') \\ & = \sup_{\eta \in \mathbb{N}} \sup_{\substack{\rho \in \mathcal{M}_+^1(\mathcal{T}) \\ \mathcal{K}(\rho, \pi) < \infty}} \int \exp \left\{ \min \left\{ \eta, \right. \right. \\ & \quad \left. \left. -\lambda \sum_{i=1}^n h(\theta', W_i) - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} \right\} \mathbb{1}\left(\frac{d\rho}{d\pi}(\theta') > 0\right) d\pi(\theta') \\ & \leq \sup_{\eta \in \mathbb{N}} \int \exp \left\{ \min \left\{ \eta, -\lambda \sum_{i=1}^n h(\theta', W_i) - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} \right\} d\pi(\theta') \\ & \stackrel{\text{monotone convergence}}{=} \int \exp \left\{ -\lambda \sum_{i=1}^n h(\theta', W_i) - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} d\pi(\theta'). \end{aligned}$$

Let us put

$$\begin{aligned} Y' &= \sup_{\rho \in \mathcal{M}_+^1(\mathcal{T})} \sup_{\eta \in \mathbb{N}} \left\{ \int \min \left\{ \eta, -\lambda \sum_{i=1}^n h(\theta', W_i) \right. \right. \\ & \quad \left. \left. - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} d\rho(\theta') - \mathcal{K}(\rho, \pi) \right\} \text{ and} \\ Y &= \log \int \exp \left\{ -\lambda \sum_{i=1}^n h(\theta', W_i) - n \log [\mathbb{P}_W \exp[-\lambda h(\theta', W)]] \right\} d\pi(\theta'). \end{aligned}$$

We just proved that $Y' \leq Y$. Moreover, Y is measurable, according to Fubini's theorem for non-negative functions. Therefore Y is a random variable. Note that we did not prove that Y' itself is measurable. Remark now that

$$\mathbb{P}_{W_1, \dots, W_n} [\exp(Y)]$$

$$\begin{aligned}
&= \mathbb{P}_{W_1, \dots, W_n} \int \exp \left\{ -\lambda \sum_{i=1}^n h(\theta', W_i) - n \log \left[\mathbb{P}_W \exp[-\lambda h(\theta', W)] \right] \right\} d\pi(\theta'), \\
&\stackrel{\text{Fubini}}{=} \int \mathbb{P}_{W_1, \dots, W_n} \exp \left\{ -\lambda \sum_{i=1}^n h(\theta', W_i) - n \log \left[\mathbb{P}_W \exp[-\lambda h(\theta', W)] \right] \right\} d\pi(\theta') \\
&= \int \left(\mathbb{1} \left(\mathbb{P}_W \left[\exp(-\lambda h(\theta', W)) \right] < +\infty \right) \prod_{i=1}^n \frac{\mathbb{P}_{W_i} \left[\exp(-\lambda h(\theta', W_i)) \right]}{\mathbb{P}_W \left[\exp(-\lambda h(\theta', W)) \right]} \right) d\pi(\theta') \leq 1,
\end{aligned}$$

proving the first part of the lemma. From Markov's inequality,

$$\mathbb{P}(Y \geq \log(\delta^{-1})) \leq \delta \mathbb{P}_{W_1, \dots, W_n} [\exp(Y)] \leq \delta.$$

Consequently $\mathbb{P}(Y \leq \log(\delta^{-1})) \geq 1 - \delta$. We have proved that the non necessarily measurable event $Y' \leq \log(\delta^{-1})$ contains the measurable event $Y \leq \log(\delta^{-1})$ whose probability is at least $1 - \delta$. \square

LEMMA 18 *Let W be a bounded random vector in a separable Hilbert space H . Let $\Theta \subset H$ be a closed and bounded set of parameters. Define $\|\Theta\| = \sup\{\|\theta\| : \theta \in \Theta\}$ and $\|W\|_\infty = \text{ess sup}\|W\|$.*

Assume that

$$\mathbb{P} \left(\inf_{\theta \in \Theta} \langle \theta, W \rangle \geq 0 \right) = 1. \quad (4.3)$$

Let W_1, \dots, W_n be a statistical sample made of n independent copies of W . Consider any number of centers $k \geq 2$, any sample size $n \geq 8k/\log(k)$ and any probability level $\delta \geq \exp(-n \log(k))$. With probability at least $1 - \delta$, for any $\theta \in \Theta^k$,

$$\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \leq \overline{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \left(\sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n} \right)^{1/4} \right) \|\Theta\| \|W\|_\infty,$$

where $\overline{\mathbb{P}}_W = \frac{1}{n} \sum_{i=1}^n \delta_{W_i}$ is the empirical measure of the sample. In expectation

$$\mathbb{P}_{W_1, \dots, W_n} \left[\sup_{\theta \in \Theta^k} \left(\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) - \overline{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \right) \right] \leq \left(\frac{8k \log(k)}{n} \right)^{1/4} \|\Theta\| \|W\|_\infty,$$

Note that condition (4.3) is not essential. Its role is only to provide slightly better constants. PROOF. Let $\Phi : H \rightarrow \ell_2 \subset \mathbb{R}^{\mathbb{N}}$ be an isometry (obtained by considering an orthonormal basis of H). Let us consider an infinite sequence of normal random variables $\varepsilon \sim \mathcal{N}(0, 1)^{\otimes \mathbb{N}} \in \mathcal{M}_+^1(\mathbb{R}^{\mathbb{N}})$, independent from the random variable X , and therefore from W . For any two random variables $U, V : \Omega \rightarrow \mathbb{R}^{\mathbb{N}}$, let us define the following extension of the inner product

$$\langle U, V \rangle = \begin{cases} \lim_{s \rightarrow \infty} \sum_{t=0}^s U_t V_t, & \text{when } \limsup_{s \rightarrow \infty} \sum_{t=0}^s U_t V_t = \liminf_{s \rightarrow \infty} \sum_{t=0}^s U_t V_t \in \mathbb{R}, \\ 0, & \text{otherwise.} \end{cases} \quad (4.4)$$

Note that the fact that U and V are measurable implies that $\langle U, V \rangle : \Omega \rightarrow \mathbb{R}$ is measurable. Remark nonetheless that this extension of the inner product is not bilinear, due to the introduction of a condition depending on the existence of the limit. Note that our construction is related to the Gaussian process

$$G(f, g) = \langle \Phi(f) + \beta^{-1/2} \varepsilon, \Phi(g) \rangle, \quad (f, g) \in H^2. \quad (4.5)$$

For any $\theta \in H$, introduce $\rho_\theta = \mathbb{P}_{\Phi(\theta) + \beta^{-1/2} \varepsilon} \in \mathcal{M}_+^1(\mathbb{R}^{\mathbb{N}})$, and put, for any $\theta \in H^k$, $\rho_\theta = \bigotimes_{j=1}^k \rho_{\theta_j} \in \mathcal{M}_+^1\left(\left(\mathbb{R}^{\mathbb{N}}\right)^k\right)$. Note that for any $\theta, \tilde{\theta} \in H$,

$$\mathcal{K}(\rho_\theta, \rho_{\tilde{\theta}}) = \sum_{i \in \mathbb{N}} \mathcal{K}(\mathbb{P}_{\Phi(\theta)_i + \beta^{-1/2} \varepsilon_i}, \mathbb{P}_{\Phi(\tilde{\theta})_i + \beta^{-1/2} \varepsilon_i}) = \sum_{i \in \mathbb{N}} \frac{\beta}{2} [\Phi(\theta)_i - \Phi(\tilde{\theta})_i]^2 = \frac{\beta \|\theta - \tilde{\theta}\|^2}{2}.$$

In the same way, for any $\theta, \tilde{\theta} \in H^k$,

$$\mathcal{K}(\rho_\theta, \rho_{\tilde{\theta}}) = \frac{\beta}{2} \sum_{j=1}^k \|\theta_j - \tilde{\theta}_j\|^2 = \frac{\beta}{2} \|\theta - \tilde{\theta}\|^2.$$

We will use a PAC-Bayesian inequality deduced from Lemma 17 on page 59 to be stated later. It requires to upper bound the following expressions, that can be seen as perturbations of exponential moments :

$$M(\theta, \lambda) = \int \log \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] d\rho_\theta(\theta'), \quad \theta \in \Theta^k, \quad \lambda > 0.$$

Note that $M(\theta, \lambda)$ being the expectation with respect to ρ_θ of a non positive random variable is always well defined in $\mathbb{R}_- \cup \{-\infty\}$. We can use Jensen's inequality to move the perturbation inside the logarithm, and Fubini's theorem for non-negative functions, to move it inside the expectation, obtaining

$$M(\theta, \lambda) \leq \log \mathbb{P}_W \left[\int \exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) d\rho_\theta(\theta') \right].$$

We can then linearize the minimum in j , remarking that

$$\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \leq \inf_{\alpha \geq 1} \left(\sum_{j=1}^k \exp \left(-\alpha \lambda \langle \theta'_j, \Phi(W) \rangle \right) \right)^{1/\alpha}.$$

Note that

$$\rho_\theta \circ (\theta'_j \mapsto \langle \theta'_j, \Phi(W) \rangle)^{-1} = \mathcal{N} \left(\langle \theta_j, W \rangle, \|W\|^2 / \beta \right),$$

meaning that under the probability measure $d\rho_\theta(\theta')$, the random variable $\langle \theta'_j, \Phi(W) \rangle$ (where we use the extension of the scalar product) is a scalar Gaussian random variable with mean $\langle \theta_j, W \rangle$ (where we use the scalar product in H) and variance $\|W\|^2 / \beta$. Using Jensen's inequality again and the formula giving the Laplace transform of a Gaussian measure, we deduce that

$$\begin{aligned}
M(\theta, \lambda) &\leq \log \mathbb{P}_W \left[\inf_{\alpha \geq 1} \left(\sum_{j=1}^k \int \exp(-\alpha \lambda \langle \theta'_j, \Phi(W) \rangle) d\rho_\theta(\theta') \right)^{1/\alpha} \right] \\
&= \log \mathbb{P}_W \left[\inf_{\alpha \geq 1} \left(\sum_{j=1}^k \exp \left(-\alpha \lambda \langle \theta_j, W \rangle + \frac{\alpha^2 \lambda^2}{2\beta} \|W\|^2 \right) \right)^{1/\alpha} \right] \\
&\leq \log \mathbb{P}_W \exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle + \inf_{\alpha \geq 1} \frac{\alpha \lambda^2}{2\beta} \|W\|^2 + \frac{\log(k)}{\alpha} \right) \\
&\leq \log \mathbb{P}_W \exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle + \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\| \right) \\
&\leq \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \log \mathbb{P}_W \exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right),
\end{aligned}$$

under the condition that $\alpha^2 = \frac{2\beta \log(k)}{\lambda^2 \|W\|^2} \geq 1$ almost surely, that is

$$\lambda^2 \|W\|_\infty^2 \leq 2\beta \log(k). \quad (4.6)$$

Notice that

$$\mathbb{P}_W \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \in [-\lambda \|\Theta\| \|W\|_\infty, 0] \right) = 1.$$

Therefore, according to Hoeffding's lemma,

$$\log \mathbb{P}_W \exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \leq -\lambda \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \frac{\lambda^2}{8} \|\Theta\|^2 \|W\|_\infty^2.$$

Indeed, for any measurable function f such that $\mathbb{P}(a \leq f(W) \leq b) = 1$, a Taylor expansion with integral remainder of $u \mapsto \log \mathbb{P}_W \left(\exp(uf(W)) \right)$ between 0 and 1 gives

$$\log \mathbb{P}_W \left(\exp(f(W)) \right) = \mathbb{P}_W(f(W)) + \int_0^1 (1-u) \mathbf{Var}_u(f(W)) du,$$

where, putting

$$\mathbb{P}_{W|u}(h(W)) = \frac{\mathbb{P}_W \left(\exp[uf(W)] h(W) \right)}{\mathbb{P}_W \left(\exp[uf(W)] \right)},$$

$$\mathbf{Var}_u(f(W)) = \mathbb{P}_{W|u} \left[\left(f(W) - \mathbb{P}_{W|u} f(W) \right)^2 \right] \leq \mathbb{P}_{W|u} \left[\left(f(W) - (a+b)/2 \right)^2 \right] \leq \frac{(b-a)^2}{4}.$$

We refer to [Cat14] for more details concerning the proof of Hoeffding's lemma and related bounds. We get

$$M(\theta, \lambda) \leq -\lambda \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \frac{\lambda^2}{8} \|\Theta\|^2 \|W\|_\infty^2 + \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty. \quad (4.7)$$

Let us apply the PAC-Bayesian inequality of Lemma 17 on page 59 to our problem. Choose $\pi = \rho_{\tilde{\theta}}$ for some deterministic value $\tilde{\theta}$ of the parameter (that we will set equal to zero later). We obtain that with probability at least $1 - \delta$,

$$\sup_{\theta \in \Theta_B^k} \sup_{\eta \in \mathbb{N}} \int \min \left\{ \eta, -\lambda \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W_i) \rangle - n \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\} \right\} d\rho_\theta(\theta') - \frac{\beta}{2} \|\theta - \tilde{\theta}\|^2 \leq \log(\delta^{-1}).$$

When θ' is distributed according to ρ_θ , $\left(\langle \theta'_j, \Phi(W_i) \rangle, j \in \llbracket 1, k \rrbracket \right) \sim \bigotimes_{j=1}^k \mathcal{N}(\langle \theta_j, W_i \rangle, \beta^{-1} \|W_i\|^2)$

is a Gaussian vector in \mathbb{R}^k with independent components and it is elementary from this remark to check that $\min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W_i) \rangle$ is integrable with respect to ρ_θ , for any (fixed) value of $W_i \in H$. Therefore, from what we proved already the negative part of

$$-\lambda \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W_i) \rangle - n \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\}$$

is integrable with respect to ρ_θ , so that the integral of this expression with respect to ρ_θ is well defined in $\mathbb{R} \cup \{+\infty\}$ and the monotone convergence theorem applied to its positive part ensures that

$$\begin{aligned} & \sup_{\eta \in \mathbb{N}} \int \min \left\{ \eta, -\lambda \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W_i) \rangle - n \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\} \right\} d\rho_\theta(\theta') \\ &= \int \left\{ -\lambda \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W_i) \rangle - n \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\} \right\} d\rho_\theta(\theta') \\ &= -\lambda \sum_{i=1}^n \int \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W_i) \rangle d\rho_\theta(\theta') - n \int \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\} d\rho_\theta(\theta') \\ &\geq -\lambda \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \int \langle \theta'_j, \Phi(W_i) \rangle d\rho_\theta(\theta') - n \int \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\} d\rho_\theta(\theta') \\ &= -\lambda \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W_i \rangle - \underbrace{n \int \log \left\{ \mathbb{P}_W \left[\exp \left(-\lambda \min_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) \right] \right\} d\rho_\theta(\theta')}_{= M(\theta, \lambda)}. \end{aligned}$$

Thus, combining the upper bound (4.7) on page 63 with Lemma 17 on page 59, we get with probability at least $1 - \delta$, for any $\theta \in \Theta^k$,

$$\begin{aligned} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) &\leq \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W_i \rangle \\ &\quad + \frac{\lambda}{8} \|\Theta\|^2 \|W\|_\infty^2 + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \frac{\beta \|\theta - \tilde{\theta}\|^2 + 2 \log(\delta^{-1})}{2n\lambda}. \end{aligned}$$

Choose $\tilde{\theta} = 0$. We obtain, with probability at least $1 - \delta$, for any $\theta \in \Theta^k$,

$$\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) - \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W_i \rangle$$

$$\begin{aligned}
&\leq \frac{\lambda}{8} \|\Theta\|^2 \|W\|_\infty^2 + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \frac{k\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{2n\lambda} \\
&\leq \sqrt{\frac{1}{4n} \left(k\beta \|\Theta\|^2 + 2 \log(\delta^{-1}) \right)} \|\Theta\| \|W\|_\infty + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty \\
&\leq \left(\sqrt{\frac{\log(\delta^{-1})}{2n}} + \sqrt{\frac{k\beta \|\Theta\|^2}{4n}} \right) \|\Theta\| \|W\|_\infty + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty \\
&\leq \left(\sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n} \right)^{1/4} \right) \|\Theta\| \|W\|_\infty,
\end{aligned}$$

where we have taken

$$\begin{aligned}
\beta &= \sqrt{\frac{8n \log(k)}{k}} \|\Theta\|^{-2}, \\
\text{and } \lambda &= 2 \sqrt{\frac{k\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{n}} \|\Theta\|^{-1} \|W\|_\infty^{-1} \\
&= 2 \sqrt{\sqrt{\frac{8k \log(k)}{n}} + \frac{2 \log(\delta^{-1})}{n}} \|\Theta\|^{-1} \|W\|_\infty^{-1}.
\end{aligned}$$

Our computations require condition (4.6) on page 63 that translates as

$$\begin{aligned}
\sqrt{\frac{2k \log(k)}{n}} + \frac{\log(\delta^{-1})}{n} &\leq \log(k) \sqrt{\frac{n \log(k)}{2k}} \\
\iff \delta &\geq \exp \left[-n \log(k) \sqrt{\frac{n \log(k)}{2k}} \left(1 - \frac{2k}{n \log(k)} \right) \right].
\end{aligned}$$

The condition is for example satisfied when $k \geq 2$, $n \geq 8k/\log(k)$ and $\delta \geq \exp(-n \log(k))$.

□

LEMMA 19 *Let W be a bounded random vector in a separable Hilbert space H . Let $\Theta \subset H$ be a closed and bounded set of parameters. Define $\|\Theta\| = \sup\{\|\theta\| : \theta \in \Theta\}$ and $\|W\|_\infty = \text{ess sup}\|W\|$. Let W_1, \dots, W_n be a statistical sample made of n independent copies of W . Consider any number $k \geq 2$ of centers, any sample size $n \geq 2k/\log(k)$ and any probability level $\delta \geq \exp(-n \log(k))$.*

Let $\bar{\mathbb{P}}_W$ be the expectation with respect to the empirical measure $\frac{1}{n} \sum_{i=1}^n \delta_{W_i}$.

With probability at least $1 - \delta$, for any $\theta \in \Theta^k$,

$$\begin{aligned}
&\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) - \inf_{\theta \in \Theta^k} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \\
&\leq \bar{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) - \inf_{\theta \in \Theta^k} \bar{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \\
&\quad + \left(\sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4} \right) \|\Theta\| \|W\|_\infty.
\end{aligned}$$

Consequently, if

$$\widehat{\theta}(W_1, \dots, W_n) \in \arg \min_{\theta \in \Theta^k} \overline{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right)$$

is an empirical minimizer, with probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \widehat{\theta}_j, W \rangle \mid W_1, \dots, W_n \right) &\leq \inf_{\theta \in \Theta^k} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \\ &\quad + \left(\sqrt{\frac{2 \log(\delta^{-1})}{n}} + 2 \left(\frac{2k \log(k)}{n} \right)^{1/4} \right) \|\Theta\| \|W\|_\infty. \end{aligned}$$

PROOF. In this proof, we will bound

$$(\mathbb{P}_W - \overline{\mathbb{P}}_W) \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right),$$

where $\overline{\mathbb{P}}_W$ is the expectation with respect to the empirical measure $\frac{1}{n} \sum_{i=1}^n \delta_{W_i}$, and where

$$\theta_j^* \in \arg \min_{\theta \in \Theta^k} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right).$$

We will use the same line of proof as in the case of the previous proposition. With the same notation, we need to bound the quantities

$$M(\theta, \lambda) = \int \log \mathbb{P}_W \left[\exp \left(\lambda \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j', \Phi(W) \rangle \right) \right) \right] d\rho_\theta(\theta'), \quad \theta \in \Theta^k, \quad \lambda > 0.$$

Acting exactly as in the previous proof, we get that

$$\begin{aligned} M(\theta, \lambda) &\leq \log \mathbb{P}_W \exp \left(\lambda \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\| \right) \\ &\leq \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \log \mathbb{P}_W \exp \left(\lambda \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \right). \end{aligned}$$

under condition (4.6) on page 63. Remark that \mathbb{P}_W almost surely

$$-\|\Theta\| \|W\|_\infty \leq \lambda \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \leq \|\Theta\| \|W\|_\infty.$$

Apply Hoeffding's inequality to deduce that

$$M(\theta, \lambda) \leq \lambda \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \frac{\lambda^2}{2} \|\Theta\|^2 \|W\|_\infty^2.$$

Operating in the same way as in the previous proof, we obtain that with probability at least $1 - \delta$ for any $\theta \in \Theta^k$,

$$\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) - \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right) - \overline{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \overline{\mathbb{P}}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right)$$

$$\begin{aligned}
&\leq \frac{\lambda}{2} \|\Theta\|^2 \|W\|_\infty^2 + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \frac{\beta \|\theta\|^2 + 2 \log(\delta^{-1})}{2n\lambda} \\
&\leq \sqrt{\frac{k\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{n}} \|\Theta\| \|W\|_\infty + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty \\
&\leq \sqrt{\frac{2 \log(\delta^{-1})}{n}} \|\Theta\| \|W\|_\infty + \sqrt{\frac{k\beta}{n}} \|\Theta\|^2 \|W\|_\infty + \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty \\
&\leq \left(\sqrt{\frac{2 \log(\delta^{-1})}{n}} + 2 \left(\frac{2k \log(k)}{n} \right)^{1/4} \right) \|\Theta\| \|W\|_\infty.
\end{aligned}$$

In these inequalities, we have chosen

$$\begin{aligned}
\beta &= \sqrt{\frac{2n \log(k)}{k}} \|\Theta\|^{-2} \\
\text{and } \lambda &= \sqrt{\frac{k\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{n}} \|\Theta\|^{-1} \|W\|_\infty^{-1} \\
&= \sqrt{\sqrt{\frac{2k \log(k)}{n}} + \frac{2 \log(\delta^{-1})}{n}} \|\Theta\|^{-1} \|W\|_\infty^{-1}.
\end{aligned}$$

Condition (4.6) on page 63 reads

$$\begin{aligned}
\sqrt{\frac{2k \log(k)}{n}} + \frac{2 \log(\delta^{-1})}{n} &\leq 2 \log(k) \sqrt{\frac{2n \log(k)}{k}} \\
\iff \delta &\geq \exp \left(-n \log(k) \left(\sqrt{\frac{2n \log(k)}{k}} - \sqrt{\frac{k}{2n \log(k)}} \right) \right).
\end{aligned}$$

It is satisfied when $k \geq 2$, $n \geq 2k/\log(k)$ and $\delta \geq \exp(-n \log(k))$. \square

4.1.1. EXPLICIT BOUND IN THE INFORMATION k -MEANS SETTING. As proved in Lemma 12 on page 43, we can use the previous bounds in the kernel space H to analyze the original information k -means minimizer.

PROPOSITION 20 *Assume that*

$$\text{ess sup}_X \left(\int p_X^2 d\nu \right) < +\infty \quad \text{and} \quad \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right) < +\infty.$$

Consider the information radius

$$R = \inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X)$$

and the bounds

$$B = \text{ess sup}_X \left(\int p_X^2 d\nu \right)^{1/2} \exp(R)$$

and $C = \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right)^{1/2}$.

Note that $R \leq \text{ess sup}_X \mathcal{K}(1, p_X) \leq C$.

Introduce the parameter space

$$\mathbf{Q}_B = \left\{ q \in \mathbb{L}_{+,1}^1(\nu) \cap \mathbb{L}^2(\nu) : \int q^2 d\nu \leq B^2 \right\}.$$

Given (X_1, \dots, X_n) , a sample made of n independent copies of X , with probability at least $1 - \delta$, for any $q \in \mathbf{Q}_B^k$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) &\leq \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_{X_i}) \\ &\quad + \left(\sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n} \right)^{1/4} \right) (BC + 2 \log(B)). \end{aligned}$$

Consider an empirical risk minimizer $\hat{q}(X_1, \dots, X_n) \in \mathbf{Q}_B^k$ satisfying

$$\hat{q} \in \arg \min_{q \in \mathbf{Q}_B^k} \overline{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right).$$

With probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(\hat{q}, p_X) \right) &\leq \inf_{q \in \left(\mathbb{L}_{+,1}^1(\nu) \right)^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) \\ &\quad + \left(\sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4} \right) (BC + 2 \log(B)). \end{aligned}$$

PROOF. The proof consists in applying Lemmas 12 on page 43, 18 on page 61 and 19 on page 65. Introduce $\theta = (q, \mathcal{K}(q, 1)) \in H$ and $W = (-\log(p_X), \mu^{-1}) \in H$. Let

$$\Theta = \{ \theta \in H : q \in \mathbf{Q}_B \}.$$

Remark that

$$\|W\|_\infty^2 = \text{ess sup}_X \|\log(p_X)\|_\nu^2 + \mu^{-1} \text{ and that } \|\theta\|^2 = \|q\|_\nu^2 + \mu \mathcal{K}(q, 1)^2.$$

According to Jensen's inequality,

$$\mathcal{K}(q, 1) = \int q \log(q) d\nu \leq \log \left(\int q^2 d\nu \right) = \log(\|q\|_\nu^2).$$

Thus $\|\Theta\|_H^2 \leq B^2 + \mu \log(B^2)^2$. Accordingly

$$\begin{aligned} \|\Theta\|^2 \|W\|_\infty^2 &\leq (C^2 + \mu^{-1})(B^2 + \mu \log(B^2)^2) \\ &= B^2 C^2 + \log(B^2)^2 + \mu^{-1} B^2 + \mu C^2 \log(B^2)^2 = (BC + 2 \log(B))^2, \end{aligned}$$

taking $\mu = \frac{B}{C \log(B^2)}$. The proposition then follows from Lemmas 18 on page 61 and 19 on page 65, in view of Lemma 12 on page 43. \square

4.1.2. CLASSICAL k -MEANS QUANTIZATION IN A SEPARABLE HILBERT SPACE. In this section, we investigate how to adapt the previous proofs in the case of the classical k -means risk in order to obtain non-asymptotic dimension-free bounds. Let us consider a separable Hilbert space $(\mathfrak{X}, \|\cdot\|)$. Let (X_1, \dots, X_n) be n independent copies of a random vector $X \sim \mathbb{P}_X$ such that $\|X\| \leq B$, almost surely with $B > 0$ and introduce the codebook (or k centers) $\mu = (\mu_1, \dots, \mu_k) \in \mathfrak{X}^k$. In this framework, the k -means risk is given by

$$\mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right),$$

and its empirical counterpart is

$$\frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \|X_i - \mu_j\|^2.$$

Let us consider the closed ball

$$\mathcal{B} = \{\mu \in \mathfrak{X} : \|\mu\| \leq B\},$$

then we have the following lemma

LEMMA 21

$$\inf_{\mu_1, \dots, \mu_k \in \mathfrak{X}^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) = \inf_{\mu_1, \dots, \mu_k \in \mathcal{B}^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right).$$

PROOF. Recall that, for a fixed $\ell : \mathfrak{X} \rightarrow \llbracket 1, k \rrbracket$, we have

$$\mathbb{P}_X \left(\|X - \mu_{\ell(X)}\|^2 \right) = \mathbb{P}_X \left(\|X - \mathbb{P}_{X|\ell(X)}(X)\|^2 \right) + \mathbb{P}_X \left(\|\mathbb{P}_{X|\ell(X)}(X) - \mu_{\ell(X)}\|^2 \right).$$

Thus optimal centers μ_j^* are given by conditional means $\mathbb{P}_{X|\ell(X)=j}(X)$. Notice that, by Jensen's inequality, $\|\mathbb{P}_{X|\ell(X)=j}(X)\| \leq \mathbb{P}_{X|\ell(X)=j}(\|X\|) \leq B$, since $\mathbb{P}_X(\|X\| \leq B) = 1$. Therefore we can restrict the optimization to the case when $\mu_j \in \mathcal{B}$. \square

Remark also that

$$\mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) = \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|\mu_j\|^2 - 2\langle \mu_j, X \rangle + \|X\|^2 \right).$$

In view of this, we can introduce the new separable Hilbert space $H = \mathfrak{X} \times \mathbb{R}^2$ endowed with the inner product

$$\langle h, h' \rangle_H = 2\langle h_1, h'_1 \rangle_{\mathfrak{X}} + B^{-2}h_2 h'_2 + B^2h_3 h'_3.$$

Define

$$\begin{aligned} \theta_j &= (\mu_j, \|\mu_j\|^2, 1) \quad \in \Theta \subset H \\ \text{where } \Theta &= \{(\mu, \|\mu\|^2, 1) : \mu \in \mathcal{B}\} \\ \text{and } W &= (-X, B^2, B^{-2}\|X\|^2) \quad \in H. \end{aligned}$$

We obtain that $\langle \theta_j, W \rangle = \|X - \mu_j\|^2$, so that

$$\inf_{\mu \in \mathcal{X}^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) = \inf_{\theta \in \Theta^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right),$$

with $\langle \theta_j, W \rangle \geq 0$. Therefore, we are in a situation where we can apply Proposition 18 on page 61 and Proposition 19 on page 65.

Remark that

$$\begin{aligned} \|W\|^2 &= 2\|X\|^2 + B^2 + B^{-2}\|X\|^4, \\ \|\theta\|^2 &= 2\|\mu\|^2 + B^{-2}\|\mu\|^4 + B^2, \end{aligned}$$

so that $\|\Theta\| \|W\|_\infty \leq 4B^2$.

PROPOSITION 22 *Let $X_i, 1 \leq i \leq n$ be a sample made of n independent copies of X . Assume that $k \geq 2$, $n \geq 2k \log(k)$ and $\delta \geq \exp(-n/4)$. With probability at least $1 - \delta$, for any $\mu = (\mu_1, \dots, \mu_k) \in \mathcal{B}^k$,*

$$\mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) \leq \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \|X_i - \mu_j\|^2 + 4B^2 \left(\sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n} \right)^{1/4} \right).$$

With probability at least $1 - \delta$, for any $\mu \in \mathcal{B}^k$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) - \inf_{\mu \in \mathcal{B}^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) \\ \leq \bar{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) - \inf_{\mu \in \mathcal{B}^k} \bar{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) \\ + 4B^2 \left(\sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4} \right), \end{aligned}$$

where $\bar{\mathbb{P}}_X = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$. In particular, if

$$\hat{\mu}(X_1, \dots, X_n) \in \arg \min_{\mu \in \mathcal{B}^k} \bar{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right),$$

with probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \hat{\mu}_j\|^2 \right) &\leq \inf_{\mu \in \mathcal{X}^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) \\ &\quad + 4B^2 \left(\sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4} \right). \end{aligned}$$

4.1.3. DISCUSSION ABOUT THE BOUNDS. We get a bound for the excess risk in the classical k -means setting, and more generally in the information k -means setting, of order

$$\mathbf{O} \left(\frac{\log(k)k}{n} \right)^{1/4}. \tag{4.8}$$

Remark that the information k -means setting is a new extension of the classical k -means setting, where we minimize the Kullback divergence with respect to its first argument. Minimization with respect to the second argument is done instead in [BDG04], within the general framework of Bregman divergence. Besides, [Fis10] develops the approach of [BDG04] using Bregman divergence and provides a non asymptotic bound for the excess risk of the same order as in [BDL08], that is

$$\mathcal{O}\left(\frac{k}{\sqrt{n}}\right). \quad (4.9)$$

Moreover, [BFL20] proves also a deviation bound of the same order as in [BDL08] in the case of robust k -means with Bregman divergence. However, the dependence in k is not explicit and depends on a constant based on the characteristics of the underlying data distribution. We will prove later on faster bounds of order

$$\mathcal{O}\left(\log\left(\frac{n}{k}\right)\sqrt{\frac{k\log(k)}{n}}\right). \quad (4.10)$$

Note that (4.9) goes to zero whenever k^2/n goes to zero, whereas (4.10) goes to zero whenever $k\log(k)\log(\log(k))/n$ goes to zero and that (4.8) goes to zero whenever $k\log(k)/n$ goes to zero, so that, although (4.8) is slower than (4.10), it provides a slightly better consistency condition.

In addition, [Lev13] derives a bound of order

$$\mathcal{O}\left(\frac{k^3}{n}\right),$$

with a better dependence in n and a worse dependence in k , under margin conditions. Margin conditions are beyond the scope of the present study, where we focus on the use of the k -means criterion as a vector quantization tool, a situation where margin conditions are unlikely to hold true. In this perspective, equation (4.8) is the most significant bound, although it is the slowest one with respect to n alone, since it provides the best sufficient condition for consistency (namely that $k\log(k)/n$ tends to zero).

4.2. A BOUNDED CRITERION FOR INFORMATION k -MEANS

In this section, we still consider a couple of random variables $(X, Y) : \Omega \rightarrow \mathcal{X} \times \mathcal{Y}$, where \mathcal{X} and \mathcal{Y} are two Polish spaces. We assume that for some reference probability measure ν , $\mathbb{P}(\mathbb{P}_{Y|X} \ll \nu) = 1$. Remark that the sigma algebra associated with \mathcal{Y} is countably generated, so that $\mathbb{L}^2(\nu)$ is separable (see prop 3.4.5 in [Coh]). We let $p_X = \frac{d\mathbb{P}_{Y|X}}{d\nu}$. For any family of probability densities $q_j \in \mathbb{L}_{+,1}^1$, $1 \leq j \leq k$, we consider a classification function

$$\ell(X) \in \arg \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X)$$

and the model

$$\mathfrak{Q}_q = \left\{ Q \in \mathcal{M}_+^1(\Omega) : Q_{Y|X} \ll \nu \text{ and } \frac{dQ_{Y|X}}{d\nu} = q_{\ell(X)} \right\}.$$

LEMMA 23 For any family $q = (q_1, \dots, q_k) \in (\mathbb{L}_{+,1}^1(\nu))^k$ of k centers, define the bounded criterion

$$\mathcal{C}(q) = 1 - \exp \left[- \inf_{Q \in \Omega_q} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) \right].$$

It can be expressed as

$$\mathcal{C}(q) = 1 - \mathbb{P}_X \left(\exp \left[- \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right] \right).$$

PROOF. Combining Lemma 1 on page 5 and Lemma 6 on page 37, we get

$$\begin{aligned} \inf_{Q \in \Omega_q} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) &= \inf_{Q \in \Omega_q} \left\{ \mathcal{K}(Q_X, P_X) + Q_X [\mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X})] \right\} \\ &= \min_{q_1, \dots, q_k} \left\{ - \log \mathbb{P}_X \left(\exp \left[- \mathcal{K}(q_{\ell(X)}, p_X) \right] \right) \right\} = - \log \mathbb{P}_X \left(\exp \left[- \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right] \right). \end{aligned}$$

This completes the proof. \square

From the Aronszajn theorem ([Aro50]), there is a mapping $\Psi : \mathbb{L}^2(\nu) \rightarrow H$ to a separable Hilbert space H such that

$$\langle \Psi(f), \Psi(g) \rangle_H = \exp \left(\int f g \, d\nu \right), \quad f, g \in \mathbb{L}^2(\nu).$$

Notice that H is separable since $K(f, g) = \exp \left(\int f g \, d\nu \right)$ is a continuous kernel (see lemma 4.33 in [CS08]). Introduce a positive real parameter μ to be chosen afterwards and remark that

$$\exp \left(- \mathcal{K}(q_j, p_X) \right) = \left\langle \exp(-\mathcal{K}(q_j, 1)) \Psi(\mu q_j), \Psi(\mu^{-1} \log(p_X)) \right\rangle_H$$

Accordingly, if we let $\theta_j = \exp(-\mathcal{K}(q_j, 1)) \Psi(\mu q_j)$ and $W = \Psi(\mu^{-1} \log(p_X))$, we obtain that

$$\mathcal{C}(q) = 1 - \mathbb{P}_X \left(\max_{j \in \llbracket 1, k \rrbracket} \underbrace{\langle \theta_j, W \rangle_H}_{\in [0,1]} \right).$$

To proceed, we need the following lemma in kernel space H .

LEMMA 24 Consider a separable Hilbert space H and a random vector $W \in H$. Let W_1, \dots, W_n be a sample made of n independent copies of W and let $\Theta \subset H$ be a bounded set of parameters.

Assume that $\mathbb{P}_W \left(\langle \theta, W \rangle \in [0, 1], \theta \in \Theta \right) = 1$.

For any number of centers $k \geq 2$, any sample size $n \geq 4k\|\Theta\|^2\|W\|_\infty^2/\log(k)$ and any probability level $\delta \geq \exp(-\log(k)n/\sqrt{2})$, with probability at least $1 - \delta$, for any $\theta \in \Theta^k$,

$$\overline{\mathbb{P}}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \leq \mathbb{P}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2}.$$

For any number of centers $k \geq 2$, any sample size $n \geq k\|\Theta\|^2\|W\|_\infty^2/\log(k)$, any probability level $\delta \geq \exp(-n\log(k)/\sqrt{2})$, any non random family of centers $\theta^* \in \Theta^k$, with probability at least $1 - \delta$, for any $\theta \in \Theta^k$

$$(\bar{\mathbb{P}}_W - \mathbb{P}_W) \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle - \max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right) \leq \sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2}.$$

Consequently if

$$\theta^* \in \arg \max_{\theta \in \Theta^k} \mathbb{P}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right)$$

and $\hat{\theta}(W_1, \dots, W_n) \in \arg \max_{\theta \in \Theta^k} \bar{\mathbb{P}}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right)$

with probability at least $1 - \delta$,

$$\mathbb{P}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right) \leq \mathbb{P}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \hat{\theta}_j, W \rangle \right) + \sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2}.$$

PROOF. The proof is almost the same as the proofs of Propositions 18 on page 61 and 19 on page 65. Reasoning in the same way, we see that

$$\begin{aligned} \int \log \mathbb{P}_W \exp \left(\lambda \max_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, W \rangle \right) d\rho_\theta(\theta') &\leq \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \log \mathbb{P}_W \exp \left(\lambda \max_{j \in \llbracket 1, k \rrbracket} \underbrace{\langle \theta_j, W \rangle}_{\in [0, 1]} \right) \\ &\leq \lambda \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \lambda \mathbb{P}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) + \frac{\lambda^2}{8}, \end{aligned}$$

under condition (4.6) on page 63. Remarking that moreover

$$\int \bar{\mathbb{P}}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta'_j, \Phi(W) \rangle \right) d\rho_\theta(\theta') \geq \bar{\mathbb{P}}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right),$$

we deduce that

$$\begin{aligned} (\bar{\mathbb{P}}_W - \mathbb{P}_W) \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) &\leq \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \frac{\lambda}{8} + \frac{k\beta\|\Theta\|^2 + 2\log(\delta^{-1})}{2n\lambda} \\ &\leq \sqrt{\frac{2 \log(k)}{\beta}} \|W\|_\infty + \sqrt{\frac{k\beta\|\Theta\|^2 + 2\log(\delta^{-1})}{4n}} \\ &\leq \sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k\log(k)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2}, \end{aligned}$$

where we have chosen

$$\beta = \sqrt{\frac{8n\log(k)}{k}} \|\Theta\|^{-1} \|W\|_\infty,$$

$$\begin{aligned}\lambda &= 2\sqrt{\frac{k\beta\|\Theta\|^2 + 2\log(\delta^{-1})}{n}} \\ &= 2\sqrt{\sqrt{\frac{8k\log(k)}{n}}\|\Theta\|\|W\|_\infty + \frac{2\log(\delta^{-1})}{n}}.\end{aligned}$$

Accordingly condition (4.6) on page 63 reads

$$\begin{aligned}\sqrt{\frac{8k\log(k)}{n}}\|\Theta\|\|W\|_\infty + \frac{2\log(\delta^{-1})}{n} &\leq \frac{1}{2}\log(k)\sqrt{\frac{8n\log(k)}{k}}\|\Theta\|^{-1}\|W\|_\infty^{-1} \\ \iff \delta &\geq \exp\left(-n\log(k)\sqrt{\frac{n\log(k)}{2k}}\|\Theta\|^{-1}\|W\|_\infty^{-1}\left(1 - \frac{2k}{n\log(k)}\|\Theta\|^2\|W\|_\infty^2\right)\right).\end{aligned}$$

Therefore the condition is satisfied when $k \geq 2$, $n \geq 4k\|\Theta\|^2\|W\|_\infty^2/\log(k)$ and $\delta \geq \exp(-n\log(k)/\sqrt{2})$. This completes the proof of the first part of the lemma. To prove the second part, remark in the same way that

$$\begin{aligned}(\bar{\mathbb{P}}_W - \mathbb{P}_W)\underbrace{\left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle - \max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle\right)}_{\in [-1, 1]} &\leq \sqrt{\frac{2\log(k)}{\beta}}\|W\|_\infty + \frac{\lambda}{2} + \frac{k\beta\|\Theta\|^2 + 2\log(\delta^{-1})}{2n\lambda} \\ &\leq \sqrt{\frac{2\log(k)}{\beta}}\|W\|_\infty + \sqrt{\frac{k\beta\|\Theta\|^2 + 2\log(\delta^{-1})}{n}} \\ &\leq \sqrt{\frac{2\log(\delta^{-1})}{n}} + \left(\frac{32k\log(k)}{n}\right)^{1/4}\|\Theta\|^{1/2}\|W\|_\infty^{1/2},\end{aligned}$$

where we have set

$$\begin{aligned}\beta &= \sqrt{\frac{2\log(k)n}{k}}\|W\|_\infty\|\Theta\|^{-1} \\ \text{and } \lambda &= \sqrt{\frac{k\beta\|\Theta\|^2 + 2\log(\delta^{-1})}{n}} \\ &= \sqrt{\sqrt{\frac{2k\log(k)}{n}}\|\Theta\|\|W\|_\infty + \frac{2\log(\delta^{-1})}{n}}.\end{aligned}$$

We have also to satisfy condition (4.6) on page 63 that reads

$$\begin{aligned}\sqrt{\frac{2k\log(k)}{n}}\|\Theta\|\|W\|_\infty + \frac{2\log(\delta^{-1})}{n} &\leq 2\log(k)\sqrt{\frac{2n\log(k)}{k}}\|\Theta\|^{-1}\|W\|_\infty^{-1} \\ \iff \delta &\geq \exp\left(-n\log(k)\sqrt{\frac{2n\log(k)}{k}}\|\Theta\|^{-1}\|W\|_\infty^{-1}\left(1 - \frac{k}{2n\log(k)}\|\Theta\|^2\|W\|_\infty^2\right)\right).\end{aligned}$$

It is satisfied when $k \geq 2$, $n \geq k\|\Theta\|^2\|W\|_\infty^2/\log(k)$ and $\delta \geq \exp(-n\log(k)/\sqrt{2})$. \square

We need now an equivalent of Lemma 12 on page 43 for the bounded criterion defined in Lemma 23 on page 72.

LEMMA 25 Assume that $\text{ess sup}_X \int p_X^2 d\nu < +\infty$ and $\text{ess sup}_X \int \log(p_X)^2 d\nu < +\infty$. Consider the information radius

$$R = \inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X)$$

and the bounds

$$B = \text{ess sup}_X \left(\int p_X^2 d\nu \right)^{1/2} \exp(R),$$

$$C = \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right)^{1/2}$$

Note that $R \leq \text{ess sup}_X \mathcal{K}(1, p_X) \leq C < +\infty$, so that $B < +\infty$. Consider the set

$$\mathcal{B} = \{q \in \mathbb{L}_{+,1}^1(\nu) : \int q^2 d\nu \leq B^2\}$$

The minimization of the bounded criterion $\mathcal{C}(q)$ defined in Lemma 23 on page 72 can be restricted to \mathcal{B} , in the sense that

$$\inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathcal{C}(q) = \inf_{q \in \mathcal{B}^k} \mathcal{C}(q).$$

PROOF. Remark that

$$\begin{aligned} \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathcal{C}(q) &= \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \left\{ 1 - \exp \left[- \inf_{Q_X} \left(\mathcal{K}(Q_X, \mathbb{P}_X) + Q_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) \right) \right] \right\} \\ &= 1 - \exp \left\{ - \inf_{Q_X} \left[\mathcal{K}(Q_X, \mathbb{P}_X) + \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} Q_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) \right] \right\} \end{aligned}$$

and apply Lemma 12 on page 43 to restrict the minimization to $q \in \mathcal{B}^k$. \square

We have seen that

$$\mathcal{C}(q) = 1 - \mathbb{P}_X \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle_H \right),$$

where $\theta_j = \exp(-\mathcal{K}(q_j, 1)) \Psi(\mu q_j)$ and $W = \Psi(\mu^{-1} \log(p_X))$. Note that for any $q \in \mathcal{B}^k$,

$$\begin{aligned} \|\theta_j\|^2 &\leq \|\Psi(\mu q_j)\|^2 = \exp(\mu^2 \|q_j\|_\nu^2) \leq \exp(\mu^2 B^2) \\ \text{and } \|W\|^2 &= \exp(\mu^{-2} \|\log(p_X)\|_\nu^2) \leq \exp(\mu^{-2} C^2), \\ \text{so that } \|\theta_j\|^2 \|W\|^2 &\leq \exp(\mu^2 B^2 + \mu^{-2} C^2) = \exp(2BC), \end{aligned}$$

if we choose $\mu = \sqrt{C/B}$. In view of Lemma 24 on page 72, this leads to

PROPOSITION 26 In the situation described at the beginning of this section, consider a sample X_1, \dots, X_n made of n independent copies of X . For any family of k probability densities $q \in (\mathbb{L}_{+,1}^1(\nu))^k$, consider the bounded criterion $\mathcal{C}(q)$ defined in Lemma 23 on page 72 and its empirical counterpart

$$\bar{\mathcal{C}}(q) = 1 - \bar{\mathbb{P}}_X \left(\exp \left[- \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right] \right).$$

Assume that $k \geq 2$, $n \geq 4k \exp(2BC)/\log(k)$ and $\delta \geq \exp(-\log(k)n/\sqrt{2})$. With probability at least $1 - \delta$, for any $q \in \mathcal{B}^k$,

$$\mathcal{C}(q) \leq \bar{\mathcal{C}}(q) + \sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n}\right)^{1/4} \exp(BC/2).$$

Let $q^* \in \mathcal{B}^k$ be a non random family of centers, and assume that $k \geq 2$, $n \geq k \exp(2BC)/\log(k)$, and $\delta \geq \exp(-n \log(k)/\sqrt{2})$. With probability at least $1 - \delta$, for any $q \in \mathcal{B}^k$,

$$\mathcal{C}(q) - \mathcal{C}(q^*) \leq \bar{\mathcal{C}}(q) - \bar{\mathcal{C}}(q^*) + \sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n}\right)^{1/4} \exp(BC/2).$$

Consequently, if

$$\hat{q}(X_1, \dots, X_n) \in \arg \min_{q \in \mathcal{B}^k} \bar{\mathcal{C}}(q),$$

with probability at least $1 - \delta$,

$$\mathcal{C}(\hat{q}) \leq \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathcal{C}(q) + \sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n}\right)^{1/4} \exp(BC/2).$$

4.3. A BOUNDED CRITERION FOR THE EUCLIDEAN k -MEANS

In this section we are given a random vector $X \in \mathbb{R}^d$ and a sample X_1, \dots, X_n made of n independent copies of X . Introduce the random variable Y whose joint distribution with X is given by $\mathbb{P}_{Y|X} = \mathcal{N}(X, \sigma^2 I_d)$, and consider a family of k centers μ_1, \dots, μ_k . Consider a nearest neighbour classification function $\ell : \mathbb{R}^d \rightarrow \llbracket 1, k \rrbracket$ that satisfies therefore $\|x - \mu_{\ell(x)}\| = \min_{j \in \llbracket 1, k \rrbracket} \|x - \mu_j\|$. Consider the conditional probability measure $Q_{Y|X}$ defined as $Q_{Y|X} = \mathcal{N}(\mu_{\ell(X)}, \sigma^2 I_d)$. Remark that the k -means Euclidean criterion can be written as

$$\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 = 2\sigma^2 \mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X}).$$

Inspired by this identity, we can introduce another loss function with the help of the model

$$\mathcal{Q}_\mu = \left\{ Q_{X,Y} : Q_{Y|X} = \mathcal{N}(\mu_{\ell(X)}, \sigma^2 I_d) \right\}.$$

It is based on the identity

$$\begin{aligned} \inf_{Q \in \mathcal{Q}_\mu} \mathcal{K}(Q_{X,Y}, \mathbb{P}_{X,Y}) &= -\log \mathbb{P}_X \left(-\mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X}) \right) \\ &= -\log \mathbb{P}_X \exp \left(-\frac{1}{2\sigma^2} \min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right). \end{aligned}$$

In view of this, one may be incited to introduce the loss function

$$\mathcal{C}(\mu) = 1 - \exp \left(-\inf_{Q \in \mathcal{Q}_\mu} \mathcal{K}(Q_{Y|X}, \mathbb{P}_{Y|X}) \right) = \mathbb{P}_X \left[\underbrace{1 - \exp \left(-\frac{1}{2\sigma^2} \min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right)}_{\in [0,1]} \right]. \quad (4.11)$$

It is the expectation of a loss function ranging in the unit interval and the corresponding empirical risk function is

$$\bar{\mathcal{C}}(\mu) = \bar{\mathbb{P}}_X \left[1 - \exp \left(-\frac{1}{2\sigma^2} \min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right) \right].$$

Observing also that

$$\mathcal{C}(\mu) = 1 - \max_{j \in \llbracket 1, k \rrbracket} \mathbb{P}_X \left(\exp \left(-\frac{1}{2\sigma^2} \|X - \mu_j\|^2 \right) \right),$$

from the Aronszajn theorem, there is a mapping $\Psi : \mathbb{R}^d \rightarrow H$ to a separable Hilbert space H such that

$$\begin{aligned} \mathcal{C}(\mu) &= 1 - \mathbb{P}_X \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \Psi(X), \Psi(\mu_j) \rangle \right) \\ &= 1 - \mathbb{P}_X \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right), \end{aligned}$$

where $\theta_j = \Psi(\mu_j)$ and $W = \Psi(X)$. Due to the Gaussian kernel, it is interesting to remark that Ψ maps the whole Euclidean space \mathbb{R}^d to the unit sphere of H . Therefore, we will not need any boundedness of even integrability assumption on X to be in a situation to apply Lemma 24 on page 72.

PROPOSITION 27 *Consider any $k \geq 2$, any $n \geq 4k/\log(k)$ and any $\delta \geq \exp(-n\log(k)/\sqrt{2})$. With probability at least $1 - \delta$, for any $\mu \in \mathbb{R}^{d \times k}$,*

$$\mathcal{C}(\mu) \leq \bar{\mathcal{C}}(\mu) + \sqrt{\frac{\log(\delta^{-1})}{2n}} + \left(\frac{8k \log(k)}{n} \right)^{1/4}.$$

Assume now that $k \geq 2$, $n \geq k/\log(k)$ and $\delta \geq \exp(-n\log(k)/\sqrt{2})$. For any non random family of centers $\mu^ \in \mathbb{R}^{d \times k}$, with probability at least $1 - \delta$, for any $\mu \in \mathbb{R}^{d \times k}$,*

$$\mathcal{C}(\mu) - \mathcal{C}(\mu^*) \leq \bar{\mathcal{C}}(\mu) - \bar{\mathcal{C}}(\mu^*) + \sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4}.$$

Consequently, if

$$\hat{\mu}(X_1, \dots, X_n) \in \arg \min_{\mu \in \mathbb{R}^{d \times k}} \bar{\mathcal{C}}(\mu),$$

with probability at least $1 - \delta$,

$$\mathcal{C}(\hat{\mu}) \leq \inf_{\mu \in \mathbb{R}^{d \times k}} \mathcal{C}(\mu) + \sqrt{\frac{2 \log(\delta^{-1})}{n}} + \left(\frac{32 k \log(k)}{n} \right)^{1/4}.$$

Note that the risk $\mathcal{C}(\mu)$ can be defined by the right-hand side of equation (4.11) in the case when X belongs to a separable Hilbert space and that in this case also Lemma 24 applies and Proposition 27 holds true, although in this case the definition of $\mathbb{P}_{Y|X}$ requires the construction of a Gaussian process similar to (4.5) on page 62.

4.4. PAC-BAYESIAN BOUNDS FOR INFORMATION FRAGMENTATION

In this section, we consider a random signal $X \in \mathbb{R}^d$, a statistical sample X_1, \dots, X_n made of n independent copies of X and a set of k fragments $\mu_j \in \mathbb{R}^d$, $1 \leq j \leq k$ that we would like to optimize to get the best possible approximation of X in terms of quantization.

More precisely, letting B_j be the support of μ_j , defined as

$$B_j = \{s \in \llbracket 1, d \rrbracket : \mu_{j,s} \neq 0\}, \quad 1 \leq j \leq k,$$

introduce for each $K \leq k$ the family of subsets

$$\mathcal{T}_{\mu,K} = \left\{ A \subset \llbracket 1, k \rrbracket : |A| \leq K \text{ and } B_i \cap B_j = \emptyset, i \neq j \in A \right\}.$$

We are willing to approximate X by $\sum_{j \in A} \mu_j$ for the best possible choice of $A \in \mathcal{T}_{\mu,K}$ depending on X . This can be seen as a structured instance of k -means, where the set of centers is

$$\sum_{j \in A} \mu_j, \quad A \in \mathcal{T}_{\mu,K}.$$

Some care should be taken though, due to the fact that $\mathcal{T}_{\mu,K}$ depends on μ and is therefore not a constant size index set. Besides, we will take advantage of the special structure of this fragment k -means problem to derive specific generalization bounds.

Let us first address the fact that $\mathcal{T}_{\mu,K}$ depends on μ . To circumvent this, we will describe the possible values of $\mathcal{T}_{\mu,K}$ first, and then we will describe the range of values of μ that corresponds to each value of $\mathcal{T}_{\mu,K}$. In other words we will condition our description of the parameter space on the value of $\mathcal{T}_{\mu,K}$.

Remark that the value of $\mathcal{T}_{\mu,K}$ depends only on the symmetric graph of the intersections of the supports

$$\mathbf{J}_\mu = \left\{ (i, j) \in \llbracket 1, k \rrbracket^2 : B_i \cap B_j \neq \emptyset \right\}.$$

Therefore, we can describe the problem in the following way. Consider the set \mathbf{G} of symmetric and reflexive graphs on the set of vertices $\llbracket 1, k \rrbracket$ (we mean by reflexive that the graph contains the diagonal of $\llbracket 1, k \rrbracket^2$). There are $2^{k(k-1)/2}$ such graphs (there are $\binom{k}{2}$ distinct pairs of vertices, so we can make $2^{\binom{k}{2}}$ possible connections), so that

$$\log(|\mathbf{G}|) = \frac{k(k-1)}{2} \log(2).$$

For each $g \in \mathbf{G}$, and each maximum number of components $K \leq k$, introduce the set of subsets

$$\mathcal{T}_{g,K} = \left\{ A \subset \llbracket 1, k \rrbracket : |A| \leq K \text{ and } A^2 \cap g = \mathbf{diag}(A^2) \right\}.$$

(In other words the set of disconnected subsets of at most K vertices.) Consider the fragment model

$$\mathcal{M}_g = \left\{ \mu \in \mathbb{R}^{d \times k} : \mathbf{J}_\mu \subset g \right\}.$$

We see that $\mathcal{T}_{g,K} \subset \mathcal{T}_{\mu,K}$, that $\mathcal{T}_{\mathcal{G}_\mu,K} = \mathcal{T}_{\mu,K}$ and that $\mu \in \mathcal{M}_{\mathcal{G}_\mu}$. Therefore

$$\left\{ \left(\sum_{j \in A} \mu_j, A \in \mathcal{T}_{\mu,K} \right) : \mu \in \mathbb{R}^{d \times k} \right\} \subset \left\{ \left(\sum_{j \in A} \mu_j, A \in \mathcal{T}_{g,K} \right) : g \in \mathcal{G}, \mu \in \mathcal{M}_g \right\}.$$

Let us now describe possible risk functions for our problem. We assume in the following discussion that the parameters k and K are fixed. The most obvious risk is the Euclidean criterion

$$\mathcal{C}_1(g, \mu) = \frac{1}{d} \mathbb{P}_X \left(\min_{A \in \mathcal{T}_{g,K}} \|X - \sum_{j \in A} \mu_j\|^2 \right), \quad g \in \mathcal{G}, \mu \in \mathcal{M}_g.$$

We normalize by $1/d$ to scale nicely with the dimension of the signal. Introducing the representation of the problem by the triplet of random variables (X, S, V) , where $\mathbb{P}_{S|X}(s) = 1/d$ and $\mathbb{P}_{V|S,X} = \mathcal{N}(X_S, \sigma^2)$, we can rewrite \mathcal{C}_1 as

$$\begin{aligned} \mathcal{C}_1(g, \mu) &= \mathbb{P}_X \left[\min_{A \in \mathcal{T}_{g,K}} \mathbb{P}_S \left(\left(X_S - \sum_{j \in A} \mu_{j,S} \right)^2 \right) \right] \\ &= 2\sigma^2 \mathbb{P}_X \left[\min_{A \in \mathcal{T}_{g,K}} \mathbb{P}_S \left(\mathcal{K}(Q_{V|S}^{(\mu, A)}, \mathbb{P}_{V|X,S}) \right) \right] \\ &= 2\sigma^2 \inf_{Q \in \mathbf{Q}_1(g, \mu)} \mathcal{K}(Q_{X,S,V}, \mathbb{P}_{X,S,V}), \end{aligned}$$

$$\text{where } Q_{V|S}^{(\mu, A)} = \mathcal{N} \left(\sum_{j \in A} \mu_{j,S}, \sigma^2 \right),$$

$$\text{and where } \mathbf{Q}_1(g, \mu) = \left\{ Q_{X,S,V} : Q_{X,S} = \mathbb{P}_{X,S}, Q_{V|X,S} = Q_{V|S}^{(\mu, A(X))}, A(X) \in \mathcal{T}_{g,K} \right\}.$$

Based on this interpretation, we can define smaller risk functions by relaxing the constraint that $Q_{X,S} = \mathbb{P}_{X,S}$. For each $\mu \in \mathcal{M}_g$ and each $A \in \mathcal{T}_{g,K}$, choose a classification function

$$\ell_{\mu, A} : \llbracket 1, d \rrbracket \longrightarrow A$$

such that

$$\text{supp}(\mu_j) \subset \ell_{\mu, A}^{-1}(j), \quad j \in A.$$

Note that

$$\sum_{j \in A} \mu_j, s = \mu \ell_{\mu, A}(s), s, \quad s \in \llbracket 1, d \rrbracket.$$

Define

$$B_j = \bigcap_{\substack{A \in \mathcal{T}_{g,K} \\ : j \in A}} \ell_{\mu, A}^{-1}(j), \quad j \in \llbracket 1, k \rrbracket.$$

and remark that

$$\text{supp}(\mu_j) \subset B_j, \quad j \in \llbracket 1, k \rrbracket.$$

Let us introduce the models

$$\mathbf{Q}_2(g, \mu) = \left\{ Q_{X,S,V} : Q_{S|X} = \mathbb{P}_S, Q_{V|X,S} = Q_{V|S}^{(A(X), \mu)}, A(X) \in \mathcal{T}_{g,K} \right\}, \quad (Q_X \text{ is free})$$

$$\mathbf{Q}_3(g, \mu) = \left\{ Q_{X, S, V} : Q_{S|X, \ell(\mu, A(X), S)=j} = \mathbb{P}_{S|S \in B_j}, Q_{V|X, S} = Q_{V|S}^{(A(X), \mu)}, A(X) \in \mathbf{T}_{g, K} \right\},$$

$(Q_{X, \ell(\mu, A(X), S)} \text{ is free})$

$$\mathbf{Q}_4(g, \mu) = \left\{ Q_{X, S, V} : Q_{S|X} \left(\bigcup_{j \in A(X)} B_j \right) = 1, Q_{V|X, S} = Q_{V|S}^{(A(X), \mu)}, A(X) \in \mathbf{T}_{g, K} \right\},$$

$(Q_{X, S|S \in \bigcup_{j \in A(X)} B_j} \text{ is free})$ and the risk functions

$$\mathbf{C}_t(g, \mu) = 2\sigma^2 \left[1 - \exp \left(- \inf_{Q \in \mathbf{Q}_t(g, \mu)} \mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V}) \right) \right], \quad t \in \llbracket 2, 4 \rrbracket.$$

LEMMA 28 For any $g \in \mathbf{G}$ and any $\mu \in \mathbf{M}_g$,

$$\mathbf{C}_2(g, \mu) \leq \mathbf{C}_1(g, \mu)$$

and

$$\mathbf{C}_4(g, \mu) \leq \mathbf{C}_3(g, \mu).$$

When the fragmentation is complete in the sense that

$$\bigcup_{j \in A} B_j = \llbracket 1, d \rrbracket, \quad A \in \overline{\mathbf{T}}_{g, K},$$

where

$$\overline{\mathbf{T}}_{g, K} = \left\{ A \in \mathbf{T}_{g, K} : A \subset A' \in \mathbf{T}_{g, K} \Rightarrow A' = A \right\}$$

are the maximal sets of $\mathbf{T}_{g, K}$, then

$$\mathbf{C}_4(g, \mu) \leq \mathbf{C}_3(g, \mu) \leq \mathbf{C}_2(g, \mu) \leq \mathbf{C}_1(g, \mu).$$

PROOF. When the fragmentation is complete

$$\mathbf{Q}_1(g, \mu) \subset \mathbf{Q}_2(g, \mu) \subset \mathbf{Q}_3(g, \mu) \subset \mathbf{Q}_4(g, \mu),$$

implying that

$$\begin{aligned} \mathbf{C}_4(g, \mu) &\leq \mathbf{C}_3(g, \mu) \leq \mathbf{C}_2(g, \mu) \leq 2\sigma^2 \left[1 - \exp \left(- \inf_{Q \in \mathbf{Q}_1(g, \mu)} \mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V}) \right) \right] \\ &\leq 2\sigma^2 \inf_{Q \in \mathbf{Q}_1(g, \mu)} \mathcal{K}(Q_{X, S, V}, \mathbb{P}_{X, S, V}) = \mathbf{C}_1(g, \mu). \end{aligned}$$

In the general case, the central inclusion $\mathbf{Q}_2(g, \mu) \subset \mathbf{Q}_3(g, \mu)$ does not hold and we cannot compare \mathbf{C}_1 or \mathbf{C}_2 with \mathbf{C}_3 or \mathbf{C}_4 . \square

LEMMA 29 For any $g \in \mathbf{G}$ and any $\mu \in \mathbf{M}_g$,

$$\mathbf{C}_2(g, \mu) = 2\sigma^2 \mathbb{P}_X \left[1 - \max_{A \in \mathbf{T}_{g, K}} \exp \left(- \frac{1}{2\sigma^2} \mathbb{P}_S \left[\left(X_S - \sum_{j \in A} \mu_{j, S} \right)^2 \right] \right) \right],$$

$$\mathbf{C}_3(g, \mu) = 2\sigma^2 \mathbb{P}_X \left[1 - \max_{A \in \mathbf{T}_{g, K}} \sum_{j \in A} \mathbb{P}_S(B_j) \exp \left(- \frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} \left[\left(X_S - \mu_{j, S} \right)^2 \right] \right) \right],$$

$$\mathbf{C}_4(g, \mu) = 2\sigma^2 \mathbb{P}_X \left[1 - \max_{A \in \mathbf{T}_{g, K}} \mathbb{P}_S \left(\sum_{j \in A} \mathbf{1}(S \in B_j) \exp \left[- \frac{1}{2\sigma^2} \left(X_S - \mu_{j, S} \right)^2 \right] \right) \right].$$

We define the empirical counterparts of our three risk functions as

$$\bar{\mathcal{C}}_1(g, \mu) = \bar{\mathbb{P}}_X \left[\min_{A \in \mathcal{T}_{g,K}} \mathbb{P}_S \left(\left(X_S - \sum_{j \in A} \mu_{j,S} \right)^2 \right) \right], \quad (4.12)$$

$$\bar{\mathcal{C}}_2(g, \mu) = 2\sigma^2 \bar{\mathbb{P}}_X \left[1 - \max_{A \in \mathcal{T}_{g,K}} \exp \left(-\frac{1}{2\sigma^2} \mathbb{P}_S \left[\left(X_S - \sum_{j \in A} \mu_{j,S} \right)^2 \right] \right) \right], \quad (4.13)$$

$$\bar{\mathcal{C}}_3(g, \mu) = 2\sigma^2 \bar{\mathbb{P}}_X \left[1 - \max_{A \in \mathcal{T}_{g,K}} \sum_{j \in A} \mathbb{P}_S(B_j) \exp \left(-\frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} \left[\left(X_S - \mu_{j,S} \right)^2 \right] \right) \right], \quad (4.14)$$

$$\bar{\mathcal{C}}_4(g, \mu) = 2\sigma^2 \bar{\mathbb{P}}_X \left[1 - \max_{A \in \mathcal{T}_{g,K}} \mathbb{P}_S \left(\sum_{j \in A} \mathbb{1}(S \in B_j) \exp \left[-\frac{1}{2\sigma^2} \left(X_S - \mu_{j,S} \right)^2 \right] \right) \right]. \quad (4.15)$$

We put $\mathcal{C}_t(\mu) = \mathcal{C}_t(\mathcal{I}_\mu, \mu)$ and consider the four optimization problems

$$\inf \left\{ \mathcal{C}_t(g, \mu) : g \in \mathcal{G}, \mu \in \mathcal{M}_g \right\} = \inf \left\{ \mathcal{C}_t(\mu) : \mu \in \mathbb{R}^{d \times k} \right\}, \quad 1 \leq t \leq 4,$$

based on the observation of the sample (X_1, \dots, X_n) , assuming that the data distribution \mathbb{P}_X is unknown to the statistician.

We can readily state generalization bounds for the first two criteria, \mathcal{C}_1 and \mathcal{C}_2 , using results for the k -means algorithm for each value of g and taking a union bound. By union bound, we mean that if $\mathfrak{B}(g, \delta)$ denotes an excess risk bound (whether for \mathcal{C}_1 or \mathcal{C}_2), for a fixed $g \in \mathcal{G}$, such that $\mathbb{P}(\mathfrak{B}(g, \delta)) \geq 1 - \delta$. Then, by the union bound we get that

$$\mathbb{P} \left(\bigcap_{g \in \mathcal{G}} \mathfrak{B}(g, \delta/|\mathcal{G}|) \right) \geq 1 - \sum_{g \in \mathcal{G}} \mathbb{P} \left(\Omega \setminus \mathfrak{B}(g, \delta/|\mathcal{G}|) \right) \geq 1 - \delta,$$

where in our case $|\mathcal{G}| = 2^{k(k-1)/2}$.

Proceeding in this way and using Propositions 22 on page 70 and 27 on page 77 leads to the following proposition.

PROPOSITION 30 Define $\bar{\mathcal{C}}_j(\mu) = \bar{\mathcal{C}}_j(\mathcal{I}_\mu, \mu)$. Assume that

$$\mathbb{P}_X \left(\sup_{s \in \llbracket 1, d \rrbracket} |X_s| \leq B \right) = 1.$$

Consider any $k \geq 2$ and any $\delta \geq \exp(-n/4)$. With probability at least $1 - \delta$, for any $\mu \in [-B, B]^{d \times k}$,

$$\mathcal{C}_1(\mu) \leq \bar{\mathcal{C}}_1(\mu) + 4B^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{4n}} + \left(\frac{8|\mathcal{T}_{\mu,K}| \log(|\mathcal{T}_{\mu,K}|)}{n} \right)^{1/4} \right).$$

For any non random set of k fragments $\mu^* \in [-B, B]^{d \times k}$, with probability at least $1 - \delta$, for any $\mu \in [-B, B]^{d \times k}$,

$$\begin{aligned} \mathcal{C}_1(\mu) - \mathcal{C}_1(\mu^*) - \bar{\mathcal{C}}_1(\mu) + \bar{\mathcal{C}}_1(\mu^*) \\ \leq 4B^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32|\mathcal{T}_{\mu,K}| \log(|\mathcal{T}_{\mu,K}|)}{n} \right)^{1/4} \right). \end{aligned}$$

Consequently, if

$$\hat{\mu} \in \arg \min_{\mu \in [-B, B]^{d \times k}} \bar{\mathcal{C}}_1(\mu) + 4B^2 \left(\frac{32 |\mathcal{T}_{\mu, K}| \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4},$$

with probability at least $1 - \delta$,

$$\mathcal{C}_1(\hat{\mu}) \leq \inf_{\mu \in \mathbb{R}^{d \times k}} \mathcal{C}_1(\mu) + 4B^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32 |\mathcal{T}_{\mu, K}| \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right).$$

PROPOSITION 31 Consider any $k \geq 2$ and any $\delta \geq \exp(-n/4)$. With probability at least $1 - \delta$, for any $\mu \in \mathbb{R}^{d \times k}$,

$$\mathcal{C}_2(\mu) \leq \bar{\mathcal{C}}_2(\mu) + 2\sigma^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{4n}} + \left(\frac{8 |\mathcal{T}_{\mu, K}| \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right).$$

For any non random set of k fragments $\mu^* \in \mathbb{R}^{d \times k}$, with probability at least $1 - \delta$, for any $\mu \in \mathbb{R}^{d \times k}$,

$$\begin{aligned} \mathcal{C}_2(\mu) - \mathcal{C}_2(\mu^*) - \bar{\mathcal{C}}_2(\mu) + \bar{\mathcal{C}}_2(\mu^*) \\ \leq 2\sigma^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32 |\mathcal{T}_{\mu, K}| \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right). \end{aligned}$$

Consequently, if

$$\hat{\mu} \in \arg \min_{\mu \in \mathbb{R}^{d \times k}} \bar{\mathcal{C}}_2(\mu) + 2\sigma^2 \left(\frac{32 |\mathcal{T}_{\mu, K}| \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4},$$

with probability at least $1 - \delta$,

$$\mathcal{C}_2(\hat{\mu}) \leq \inf_{\mu \in \mathbb{R}^{d \times k}} \mathcal{C}_2(\mu) + 2\sigma^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32 |\mathcal{T}_{\mu, K}| \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right).$$

The previous bounds depend on the factor $|\mathcal{T}_{\mu, K}|$, that is of order 2^k , and therefore too large in many situations. To get rid of this term, we will use the following lemma, that is specific to the fragmentation setting.

LEMMA 32 Let $W = (W_j, 1 \leq j \leq k)$ be a random vector in the product H^k , where H is a separable Hilbert space (that we can take as being ℓ_2 if we want). Consider a sample $(W^{(1)}, \dots, W^{(n)})$ made of n independent copies of W . Consider a bounded parameter set $\Theta \subset H^k$ and a set \mathcal{T} of subsets of $\llbracket 1, k \rrbracket$. Assume that

$$\mathbb{P}_W \left(\sum_{j \in A} \langle \theta_j, W_j \rangle \in [a, b], A \in \mathcal{T}, \theta \in \Theta \right) = 1.$$

Consider the risk

$$\mathcal{C}(\theta) = \mathbb{P}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right), \quad \theta \in \Theta,$$

and its empirical counterpart

$$\bar{\mathcal{C}}(\theta) = \bar{\mathbb{P}}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right), \quad \theta \in \Theta$$

Put

$$K(\mathcal{T}) = \max_{A \in \mathcal{T}} |A|, \quad \|\Theta\| = \sup_{\theta \in \Theta} \left(\sum_{j=1}^k \|\theta_j\|^2 \right)^{1/2} \quad \text{and} \quad \|W\|_\infty = \max_{j \in \llbracket 1, k \rrbracket} \text{ess sup}_{\mathbb{P}_W} \|W_j\|.$$

Assume that $|\mathcal{T}| \geq 3$ and that $\delta \geq \exp[-n \log(|\mathcal{T}|)/\sqrt{2}]$. With probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\mathcal{C}(\theta) \leq \bar{\mathcal{C}}(\theta) + \sqrt{\frac{\log(\delta^{-1})}{2n}}(b-a) + \left(\frac{8K(\mathcal{T}) \log(|\mathcal{T}|)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2} (b-a)^{1/2}.$$

Consider any non random value of the parameter $\theta^* \in \Theta$. With probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\begin{aligned} \mathcal{C}(\theta) - \mathcal{C}(\theta^*) &\leq \bar{\mathcal{C}}(\theta) - \bar{\mathcal{C}}(\theta^*) \\ &\quad + \sqrt{\frac{2 \log(\delta^{-1})}{n}}(b-a) + \left(\frac{32 K(\mathcal{T}) \log(|\mathcal{T}|)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2} (b-a)^{1/2}. \end{aligned}$$

Consequently, if

$$\hat{\theta} \in \arg \min_{\theta \in \Theta} \bar{\mathcal{C}}(\theta),$$

with probability at least $1 - \delta$,

$$\mathcal{C}(\hat{\theta}) \leq \inf_{\theta \in \Theta} \mathcal{C}(\theta) + \sqrt{\frac{2 \log(\delta^{-1})}{n}}(b-a) + \left(\frac{32 K(\mathcal{T}) \log(|\mathcal{T}|)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2} (b-a)^{1/2}.$$

In expectation

$$\mathbb{P}_{W^{(1)}, \dots, W^{(n)}} \left(\mathcal{C}(\hat{\theta}) \right) \leq \inf_{\theta \in \Theta} \mathcal{C}(\theta) + \left(\frac{32 K(\mathcal{T}) \log(|\mathcal{T}|)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2} (b-a)^{1/2}.$$

PROOF. The proof follows the same line as Lemma 18 on page 61 and 19 on page 65. To prove the first part of the lemma, we need to bound

$$M(\theta, \lambda) = \int \log \left[\mathbb{P}_W \left(\exp \left(-\lambda \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta'_j, \Phi(W_j) \rangle \right) \right) \right] d\rho_\theta(\theta'), \quad \theta \in \Theta, \lambda > 0.$$

and to apply Lemma 17 on page 59. First remark that

$$M(\theta, \lambda) \leq \int \log \mathbb{P}_W \left[\inf_{\alpha \geq 1} \left(\sum_{A \in \mathcal{T}} \exp \left(-\alpha \lambda \sum_{j \in A} \langle \theta'_j, \Phi(W_j) \rangle \right) \right)^{1/\alpha} \right] d\rho_\theta(\theta').$$

Then use Jensen's inequality and Fubini's theorem to move the integration with respect to ρ_θ inside to get

$$M(\theta, \lambda) \leq \log \mathbb{P}_W \left[\inf_{\alpha \geq 1} \left(\sum_{A \in \mathcal{T}} \int \exp \left(-\alpha \lambda \sum_{j \in A} \langle \theta'_j, \Phi(W_j) \rangle \right) d\rho_\theta(\theta') \right)^{1/\alpha} \right]$$

$$\begin{aligned}
&= \log \mathbb{P}_W \left[\inf_{\alpha \geq 1} \left(\sum_{A \in \mathcal{T}} \exp \left(-\alpha \lambda \sum_{j \in A} \langle \theta_j, W_j \rangle + \frac{\alpha^2 \lambda^2}{2\beta} \sum_{j \in A} \|W_j\|^2 \right) \right)^{1/\alpha} \right] \\
&\leq \log \mathbb{P}_W \left[\inf_{\alpha \geq 1} \exp \left(\log(|\mathcal{T}|)/\alpha - \lambda \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle + \frac{\alpha \lambda^2}{2\beta} K(\mathcal{T}) \|W\|_\infty^2 \right) \right] \\
&= \lambda \|W\|_\infty \sqrt{\frac{2K(\mathcal{T}) \log(|\mathcal{T}|)}{\beta}} + \log \mathbb{P}_W \left[\exp \left(-\lambda \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right) \right].
\end{aligned}$$

The above inequalities require that $\alpha \geq 1$, that is

$$\lambda^2 K(\mathcal{T}) \|W\|_\infty^2 \leq 2\beta \log(|\mathcal{T}|). \quad (4.16)$$

Remembering Hoeffding's lemma (used in the course of the proof of Lemma 18 on page 61), we get

$$M(\theta, \lambda) \leq \lambda \|W\|_\infty \sqrt{\frac{2K(\mathcal{T}) \log(|\mathcal{T}|)}{\beta}} - \lambda \mathbb{P}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right) + \frac{\lambda^2}{8} (b-a)^2.$$

Thus according to Lemma 17 on page 59, with probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\begin{aligned}
\mathbb{P}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right) &\leq \int \bar{\mathbb{P}}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta'_j, \Phi(W_j) \rangle \right) d\rho_\theta(\theta') \\
&\quad + \|W\|_\infty \sqrt{\frac{2K(\mathcal{T}) \log(|\mathcal{T}|)}{\beta}} + \frac{\lambda}{8} (b-a)^2 + \frac{\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{2n\lambda}.
\end{aligned}$$

Note that

$$\begin{aligned}
\int \bar{\mathbb{P}}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta'_j, \Phi(W_j) \rangle \right) d\rho_\theta(\theta') \\
&\leq \bar{\mathbb{P}}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \int \langle \theta'_j, \Phi(W_j) \rangle d\rho_\theta(\theta') \right) = \bar{\mathbb{P}}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right).
\end{aligned}$$

Take

$$\lambda = 2 \sqrt{\frac{\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{n(b-a)^2}}$$

to obtain

$$\begin{aligned}
&(\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right) \\
&\leq \|W\|_\infty \sqrt{\frac{2K(\mathcal{T}) \log(|\mathcal{T}|)}{\beta}} + (b-a) \sqrt{\frac{\beta \|\Theta\|^2 + 2 \log(\delta^{-1})}{4n}} \\
&\leq \|W\|_\infty \sqrt{\frac{2K(\mathcal{T}) \log(|\mathcal{T}|)}{\beta}} + (b-a) \sqrt{\frac{\beta \|\Theta\|^2}{4n}} + (b-a) \sqrt{\frac{\log(\delta^{-1})}{2n}}.
\end{aligned}$$

Then choose

$$\beta = \|W\|_\infty \|\Theta\|^{-1} (b-a)^{-1} \sqrt{8nK(\mathcal{T}) \log(|\mathcal{T}|)}$$

to get

$$\begin{aligned} (\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right) \\ \leq \sqrt{\frac{\log(\delta^{-1})}{2n}} (b-a) + \left(\frac{8K(\mathcal{T}) \log(|\mathcal{T}|)}{n} \right)^{1/4} \|\Theta\|^{1/2} \|W\|_\infty^{1/2} (b-a)^{1/2}. \end{aligned}$$

and

$$\lambda = \frac{2}{(b-a)} \sqrt{\frac{\|W\|_\infty \|\Theta\|}{(b-a)} \sqrt{\frac{8K(\mathcal{T}) \log(|\mathcal{T}|)}{n}} + \frac{2 \log(\delta^{-1})}{n}}.$$

We have to satisfy condition (4.16) on page 84 that reads

$$\begin{aligned} \frac{\|W\|_\infty \|\Theta\|}{(b-a)} \sqrt{\frac{8K(\mathcal{T}) \log(|\mathcal{T}|)}{n}} + \frac{2 \log(\delta^{-1})}{n} \leq \frac{\log(|\mathcal{T}|)}{2} \sqrt{\frac{8n \log(|\mathcal{T}|)}{K(\mathcal{T})}} \frac{(b-a)}{\|W\|_\infty \|\Theta\|} \\ \iff \delta \geq \exp \left(-n \log(|\mathcal{T}|) \sqrt{\frac{n \log(|\mathcal{T}|)}{2K(\mathcal{T})}} \frac{(b-a)}{\|W\|_\infty \|\Theta\|} \left(1 - \frac{2K(\mathcal{T}) \|W\|_\infty^2 \|\Theta\|^2}{n \log(|\mathcal{T}|) (b-a)^2} \right) \right). \end{aligned}$$

The condition is satisfied when $|\mathcal{T}| \geq 2$,

$$n \geq \frac{4K(\mathcal{T}) \|W\|_\infty^2 \|\Theta\|^2}{\log(|\mathcal{T}|) (b-a)^2} \quad (4.17)$$

and $\delta \geq \exp[-n \log(|\mathcal{T}|)/\sqrt{2}]$. When $|\mathcal{T}| \geq 3$ and condition (4.17) is not satisfied, the generalization bound is larger than $b-a$ and therefore trivially true. Thus we can drop condition (4.17) if we assume that $|\mathcal{T}| \geq 3$.

To prove the second part of the lemma, we have got to study

$$M(\theta, \lambda) = \int \log \left[\mathbb{P}_W \left(\exp \left(\lambda \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j^*, W_j \rangle - \lambda \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j', W_j \rangle \right) \right) \right] d\rho_\theta(\theta'), \quad \theta \in \Theta, \lambda > 0.$$

We can remark that \mathbb{P}_W almost surely

$$\left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j^*, W_j \rangle - \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j', W_j \rangle \right) \in [a-b, b-a]$$

and proceed as in the first part of the proof to establish that

$$M(\theta, \lambda) \leq \lambda \|W\|_\infty \sqrt{\frac{2K(\mathcal{T}) \log(|\mathcal{T}|)}{\beta}} + \lambda \mathbb{P}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j^*, W_j \rangle - \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j', W_j \rangle \right) + \frac{\lambda^2}{2} (b-a)^2.$$

So the situation is the same as in the first part with a variance term increased by a factor four, and except for this small difference the computations are the same. \square

To apply Lemma 32 on page 82, we need to put the risk functions \mathbf{C}_t , $t \in \{1, 3, 4\}$ in a suitable form, that is to express the different risks in terms of a sum of inner products of parameters θ_j and random vectors W_j belonging to some separable Hilbert space. Let us notice that we will not be able to express criterion \mathbf{C}_2 in this form, since the summation over $j \in A$ is inside the exponential.

Let us start with \mathbf{C}_1 , assuming that $X \in [-B, B]^d$, so that we can also assume that $\mu \in [-B, B]^{d \times k}$, since shrinking μ to that range decreases the risk. Remark accordingly that

$$\mathbf{C}_1(g, \mu) = \mathbb{P}_X \left\{ \min_{A \in \mathcal{T}_{g, K}} \left[\mathbb{P}_S \left(\left(\sum_{j \in A} \mu_{j,S} \right)^2 \right) - 2 \sum_{j \in A} \mathbb{P}_S(X_S \mu_{j,S}) \right] \right\} + \mathbb{P}_X \left(\mathbb{P}_S(X_S^2) \right)$$

Moreover, since the supports are disjoint,

$$\left(\sum_{j \in A} \mu_{j,S} \right)^2 = \sum_{j \in A} \mu_{j,S}^2, \quad A \in \mathcal{T}_{g, K}.$$

Thus

$$\mathbf{C}_1(g, \mu) = \mathbb{P}_X \left(\min_{A \in \mathcal{T}_{g, K}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right) + \mathbb{P}_X \left(\mathbb{P}_S(X_S^2) \right),$$

where $\theta_j = (\mu_j, B^{-1} \mathbb{P}_S(\mu_{j,S}^2)) \in \mathbb{L}^2(\mathbb{P}_S) \times \mathbb{R}$ and $W_j = (-2X, B)$, in the same space. Note that

$$\begin{aligned} \|\theta_j\|^2 &= \mathbb{P}_S(\mu_{j,S}^2) + B^{-2} \mathbb{P}_S(\mu_{j,S}^2)^2 \leq 2B^2 \mathbb{P}_S(\text{supp}(\mu_j)), \\ \text{and that } \|W_j\|^2 &= 4\mathbb{P}_S(X_S^2) + B^2 \leq 5B^2. \end{aligned}$$

Moreover

$$\begin{aligned} -B^2 \leq -\mathbb{P}_S(X_S^2) &\leq \sum_{j \in A} \langle \theta_j, W_j \rangle \\ &\leq \mathbb{P}_S \left[\left(X_S - \sum_{j \in A} \mu_{j,S} \right)^2 \right] - \mathbb{P}_S(X_S^2) \leq \mathbb{P}_S(X_S^2) + 2\mathbb{P}_S \left[\left(\sum_{j \in A} \mu_{j,S} \right)^2 \right] \leq 3B^2. \end{aligned}$$

Consider the model

$$\mathcal{M}(\mathbf{S}) = \left\{ \mu \in [-B, B]^{d \times k} : \sum_{j=1}^k \mathbb{P}_S(\text{supp}(\mu_j)) \leq \mathbf{S} \right\}$$

and apply Lemma 32 on page 82 to the corresponding parameter set Θ . Taking into account the fact that

$$\|\Theta\|^2 \leq 2B^2 \mathbf{S},$$

we obtain

PROPOSITION 33 *Consider any $k \geq 3$, any $K \geq 1$, any $\delta \geq \exp(-n \log(k)/\sqrt{2})$ and any $\mathbf{S} \in [1, k]$. With probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathbf{S})$,*

$$\begin{aligned} \mathcal{C}_1(\mu) - \mathbb{P}_X[\mathbb{P}_S(X_S^2)] &\leq \overline{\mathcal{C}}_1(\mu) - \overline{\mathbb{P}}_X[\mathbb{P}_S(X_S^2)] \\ &\quad + 4B^2 \left(\sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{4n}} + \left(\frac{5K\mathbf{S}\log|\mathcal{T}_{\mu,K}|}{n} \right)^{1/4} \right). \end{aligned}$$

For any non random set of fragments $\mu^* \in \mathbf{M}(\mathcal{S})$, with probability at least $1 - \delta$, for any $\mu \in \mathbf{M}(\mathcal{S})$,

$$\begin{aligned} \mathcal{C}_1(\mu) - \mathcal{C}_1(\mu^*) - \overline{\mathcal{C}}_1(\mu) + \overline{\mathcal{C}}_1(\mu^*) \\ \leq 4B^2 \left(\sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{n}} + \left(\frac{20K\mathbf{S}\log|\mathcal{T}_{\mu,K}|}{n} \right)^{1/4} \right). \end{aligned}$$

Consequently, if

$$\widehat{\mu} \in \arg \min_{\mu \in \mathbf{M}(\mathcal{S})} \overline{\mathcal{C}}_1(\mu) + 4B^2 \left(\frac{20K\mathbf{S}\log|\mathcal{T}_{\mu,K}|}{n} \right)^{1/4},$$

with probability at least $1 - \delta$,

$$\mathcal{C}_1(\widehat{\mu}) \leq \inf_{\mu \in \mathbf{M}(\mathcal{S})} \mathcal{C}_1(\mu) + 4B^2 \left(\sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{n}} + \left(\frac{20K\mathbf{S}\log|\mathcal{T}_{\mu,K}|}{n} \right)^{1/4} \right).$$

Let us now turn our attention to the risk function \mathcal{C}_3 . Consider a fix set of subsets $\{B_j \subset \llbracket 1, d \rrbracket : 1 \leq j \leq k\}$. Define accordingly

$$\mathcal{T}_{B,K} = \left\{ A \subset \llbracket 1, k \rrbracket : |A| \leq K, B_i \cap B_j = \emptyset, (i, j) \in A^2, i \neq j \right\}.$$

Consider some separable Hilbert space H (for instance ℓ_2 , the space of square integrable real valued sequences). For each $j \in \llbracket 1, k \rrbracket$, there is a mapping

$$\Psi_j : \mathbb{R}^d \longrightarrow H$$

such that

$$\langle \Psi_j(x), \Psi_j(y) \rangle = \exp \left(-\frac{1}{2\sigma^2} \mathbb{P}_{S|S \in B_j} [(x_S - y_S)^2] \right).$$

Introduce the model

$$\mathbf{M}_B = \left\{ \mu \in \mathbb{R}^{d \times k} : \text{supp}(\mu_j) \subset B_j \right\}.$$

For any (B, μ) , the risk $\mathcal{C}_3(B, \mu)$ can be written as

$$\mathcal{C}_3(B, \mu) = 1 + \mathbb{P}_X \left(\min_{A \in \mathcal{T}_{B,K}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right),$$

where $\theta_j = \mathbb{P}_S(B_j)\Psi_j(\mu_j)$ and $W_j = -\Psi_j(X)$. Note that

$$\sum_{j=1}^k \|\theta_j\|^2 = \sum_{j=1}^k \mathbb{P}_S(B_j)$$

and that

$$\|W_j\| = 1,$$

while almost surely

$$\sum_{j \in A} \langle \theta_j, W_j \rangle \in [-1, 0].$$

Remark also that $\mathbf{C}_3(\mu) = \mathbf{C}_3(\mathbf{J}_\mu, \mu) = \mathbf{C}_3(\text{supp}(\mu), \mu)$. Applying Lemma 32 on page 82 gives

PROPOSITION 34 *Consider the model*

$$\mathbf{M}(\mathbf{S}) = \{\mu \in \mathbb{R}^{d \times k} : \sum_{j=1}^k \mathbb{P}_S(\text{supp}(\mu_j)) \leq \mathbf{S}, |\mathcal{T}_{\mu, K}| \geq 3\}.$$

Assume that $\delta \geq \exp(-n \log(3)/\sqrt{2})$ and that $\mathbf{S} \in [1, k]$. With probability at least $1 - \delta$, for any $\mu \in \mathbf{M}(\mathbf{S})$,

$$\mathbf{C}_3(\mu) \leq \bar{\mathbf{C}}_3(\mu) + 2\sigma^2 \left(\sqrt{\frac{kd \log(2) + \log(\delta^{-1})}{2n}} + \left(\frac{8K\mathbf{S} \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right)$$

Consider a non random set of fragments $\mu^* \in \mathbf{M}(\mathbf{S})$. With probability at least $1 - \delta$, for any $\mu \in \mathbf{M}(\mathbf{S})$,

$$\begin{aligned} \mathbf{C}_3(\mu) - \mathbf{C}_3(\mu^*) &\leq \bar{\mathbf{C}}_3(\mu) - \bar{\mathbf{C}}_3(\mu^*) \\ &\leq 2\sigma^2 \left(\sqrt{\frac{2kd \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32K\mathbf{S} \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right). \end{aligned}$$

Consequently, if

$$\hat{\mu} \in \arg \min_{\mu \in \mathbf{M}(\mathbf{S})} \bar{\mathbf{C}}_3(\mu) + 2\sigma^2 \left(\frac{32K\mathbf{S} \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4},$$

with probability at least $1 - \delta$,

$$\mathbf{C}_3(\hat{\mu}) \leq \inf_{\mu \in \mathbf{M}(\mathbf{S})} \mathbf{C}_3(\mu) + 2\sigma^2 \left(\sqrt{\frac{2kd \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32K\mathbf{S} \log(|\mathcal{T}_{\mu, K}|)}{n} \right)^{1/4} \right).$$

The strong point of this proposition is that we do not have to assume that the signal X or the fragments μ_j are bounded. The weak point is that the union bound on B_j (that is essentially on the choice of $\text{supp}(\mu_j)$) introduce a dependence of the bound on the dimension d of the signal. Accordingly, when X is a digital image of dimension $d = 10^6$ or so, the proposition is not very meaningful, although it is still interesting within a submodel $\mathbf{M}(B, \mathbf{S})$ where B is fixed and accordingly the term $kd \log(2)$ is not present.

To get a generalization bound that is independent of the dimension d and does not require boundedness or integrability assumptions on the signal, we can turn to the smaller and therefore less demanding risk function \mathbf{C}_4 .

There is a mapping $\Psi : \mathbb{R} \rightarrow H$ of the real line into a reproducing kernel separable Hilbert space H such that

$$\langle \Psi(x), \Psi(y) \rangle_H = \exp\left(-\frac{1}{2\sigma^2}(y - x)^2\right), \quad x, y \in \mathbb{R}.$$

We can then consider $\mathbb{L}^2(H^{[1,d]}, \mathbb{P}_S)$, another separable Hilbert space whose scalar product is defined as

$$\langle f, g \rangle = \mathbb{P}_S \left(\langle f(S), g(S) \rangle_H \right), \quad f, g : \llbracket 1, d \rrbracket \rightarrow H.$$

Define for any (g, μ) such that $g \in \mathfrak{G}$ and $\mu \in \mathcal{M}_g$

$$\begin{aligned} \theta_j(s) &= \mathbb{1}(s \in B_j) \Psi(\mu_{j,s}), \\ X_j(s) &= -\Psi(X_s), \quad 1 \leq s \leq d, \quad 1 \leq j \leq k. \end{aligned}$$

We see that

$$\mathfrak{C}_4(g, \mu) = 2\sigma^2 \left[1 + \mathbb{P}_X \left(\min_{A \in \mathcal{T}_{g,K}} \underbrace{\sum_{j \in A} \langle \theta_j, W_j \rangle}_{\in [-1,0]} \right) \right].$$

Moreover $\|\theta_j\|^2 = \mathbb{P}_S(B_j)$ and $\|W_j\| = 1$. We obtain in view of Lemma 32 on page 82

PROPOSITION 35 *Define the model*

$$\mathcal{M}(\mathfrak{S}) = \left\{ \mu \in \mathbb{R}^{d \times k} : \sum_{j=1}^k \mathbb{P}_S(\text{supp}(\mu_j)) \leq \mathfrak{S}, |\mathcal{T}_{\mu,K}| \geq 3 \right\},$$

with $\mathfrak{S} \in [1, k]$ and assume that $\delta \geq \exp(-n \log(3)/\sqrt{2})$. With probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathfrak{S})$,

$$\mathfrak{C}_4(\mu) \leq \bar{\mathfrak{C}}_4(\mu) + 2\sigma^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{4n}} + \left(\frac{8K\mathfrak{S} \log(|\mathcal{T}_{\mu,K}|)}{n} \right)^{1/4} \right).$$

Moreover, if $\mu^* \in \mathcal{M}(\mathfrak{S})$ is a non random set of fragments, with probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathfrak{S})$,

$$\begin{aligned} \mathfrak{C}_4(\mu) - \mathfrak{C}_4(\mu^*) - \bar{\mathfrak{C}}_4(\mu) + \bar{\mathfrak{C}}_4(\mu^*) \\ \leq 2\sigma^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32K\mathfrak{S} \log(|\mathcal{T}_{\mu,K}|)}{n} \right)^{1/4} \right). \end{aligned}$$

Consequently, if

$$\hat{\mu} \in \arg \min_{\mu \in \mathcal{M}(\mathfrak{S})} \bar{\mathfrak{C}}_4(\mu) + 2\sigma^2 \left(\frac{32K\mathfrak{S} \log(|\mathcal{T}_{\mu,K}|)}{n} \right)^{1/4},$$

with probability at least $1 - \delta$

$$\mathfrak{C}_4(\hat{\mu}) \leq \inf_{\mu \in \mathcal{M}(\mathfrak{S})} \mathfrak{C}_4(\mu) + 2\sigma^2 \left(\sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} + \left(\frac{32K\mathfrak{S} \log(|\mathcal{T}_{\mu,K}|)}{n} \right)^{1/4} \right).$$

We see from this proposition that when we work with the robust risk \mathfrak{C}_4 , we can obtain a dimension free generalization bound that does not require any boundedness or integrability condition on the signal X , while \mathfrak{C}_4 still carries a meaningful notion of quantization. Another useful remark to understand the bound is that

$$\begin{aligned}
\log(|\mathcal{T}_{\mu,K}|) &\leq \log \left[\sum_{m=0}^K \binom{k}{m} \right] \leq \log \left[\sum_{m=0}^K \frac{k^m}{m!} \right] \\
&\leq \log \left[\left(\frac{k}{K} \right)^K \sum_{m=0}^K \frac{K^m}{m!} \right] \leq K \log \left(\frac{ek}{K} \right).
\end{aligned}$$

Thus the complexity term is bounded by

$$\left(\frac{32 K^2 \mathbf{S} \log(ek/K)}{n} \right)^{1/4}.$$

A second important remark we can make is that the upper bounds we obtained are increasing with respect to the term \mathbf{S} , which represents the maximum area covered by the fragments. In that way, if we diminish \mathbf{S} , we obtained a tighter bound. Accordingly, the bounds advocate to some extent the use of the fragmentation algorithm described in section 2.3 on page 25, since the fragmentation algorithm decreases \mathbf{S} at each step.

Besides, one can derive generalization bounds that are uniform with respect to \mathbf{S} , using a union bound over an appropriate grid containing a geometric progression of \mathbf{S} values. This requires to pay a small cost in the bound.

4.5. FASTER BOUNDS

In this section, we derive faster bounds for the different quantization criteria we developed so far. To this aim, we will borrow ideas from the classical chaining method used to upper bound the expected supremum of Gaussian processes (see [BLM13]). However, we will choose a PAC-Bayesian approach to this chaining technique in the sense that we will use a sequence of perturbations of the parameter parametrized by a variance ranging in a geometric grid, which will play the role of the δ -covering sets in the classical chaining argument.

4.5.1. FASTER BOUNDS FOR INFORMATION k -MEANS AND CLASSICAL k -MEANS. We can reach faster speeds than in Lemma 18 on page 61 and the likes through some kind of PAC-Bayesian chaining.

LEMMA 36 *Let W be a random vector in a separable Hilbert space H . Let (W_1, \dots, W_n) be a sample made of n independent copies of W . Let $\Theta \subset H^k$ be a closed and bounded set of parameters. Define*

$$\|\Theta\| = \sup \left\{ \left(\sum_{j=1}^k \|\theta_j\|^2 \right)^{1/2} : \theta \in \Theta \right\}$$

and assume that

$$\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \in [a, b] \text{ for all } \theta \in \Theta \right) = 1.$$

For any $k \geq 2$, any $n \geq 2k$ and any $\delta \in]0, 1[$, with probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\begin{aligned}
\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) &\leq \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W_i \rangle \\
&\quad + \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8 \log(k)}{n}} + 2 \sqrt{\frac{\log(k)}{n}} \right) \|\Theta\| \|W\|_\infty \\
&\quad + \sqrt{\frac{(\sqrt{2}+1) \left(k(b-a)^2 + 2 \log(ek) \|W\|_\infty^2 \|\Theta\|^2 \right)}{n}} + \sqrt{\frac{\log(\delta^{-1})}{2n}} (b-a).
\end{aligned}$$

If $\theta^* \in \Theta$ is a non random value of the parameter, with probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\begin{aligned}
&\left(\mathbb{P}_W - \bar{\mathbb{P}}_W \right) \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle - \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j^*, W \rangle \right) \\
&\leq \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8 \log(k)}{n}} + 2 \sqrt{\frac{\log(k)}{n}} \right) \|\Theta\| \|W\|_\infty \\
&\quad + \sqrt{\frac{(\sqrt{2}+1) \left(k(b-a)^2 + 2 \log(ek) \|W\|_\infty^2 \|\Theta\|^2 \right)}{n}} + \sqrt{\frac{2 \log(\delta^{-1})}{n}} (b-a).
\end{aligned}$$

Therefore in the case when

$$\hat{\theta} \in \arg \min_{\theta \in \Theta} \bar{\mathbb{P}} \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right),$$

$\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \hat{\theta}_j, W \rangle \right) - \inf_{\theta \in \Theta} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right)$ satisfies the same bound with the same probability.

Moreover, the expected excess risk satisfies

$$\begin{aligned}
&\mathbb{P}_{W_1, \dots, W_n} \left[\mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \hat{\theta}_j, W \rangle \right) - \inf_{\theta \in \Theta} \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) \right] \\
&\leq \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8 \log(k)}{n}} + 2 \sqrt{\frac{\log(k)}{n}} \right) \|\Theta\| \|W\|_\infty \\
&\quad + \sqrt{\frac{(\sqrt{2}+1) \left(k(b-a)^2 + 2 \log(ek) \|W\|_\infty^2 \|\Theta\|^2 \right)}{n}}.
\end{aligned}$$

PROOF. Let

$$\rho_{\theta' \mid \theta} = \mathbb{P}_{\theta_i + \beta^{-1/2} \varepsilon_i, i \in \mathbb{N}}$$

be a Gaussian conditional probability distribution with values in $\mathcal{M}_+^1(\mathbb{R}^{\mathbb{N}})$, where ε_i , $i \in \mathbb{N}$ is an infinite sequence of independent standard normal random variables. When θ and $\theta' \in \mathbb{R}^{\mathbb{N} \times k}$ are made of k infinite sequences of real numbers, let

$$\rho_{\theta' \mid \theta} = \bigotimes_{j=1}^k \rho_{\theta'_j \mid \theta_j}$$

be the tensor product of the previously defined conditional probability distributions. Let W be a random vector in the separable Hilbert space $\ell_2 \subset \mathbb{R}^{\mathbb{N}}$. Consider the functions

$$f(\theta, w) = \min_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, w \rangle, \quad \theta \in \mathbb{R}^{\mathbb{N} \times k}, w \in \mathbb{R}^{\mathbb{N}},$$

where the scalar product is extended beyond ℓ_2 as already explained in a measurable but not bilinear way (see equation (4.4) on page 61). Let

$$\bar{f}(\theta, w) = f(\theta, w) - \mathbb{P}_W(f(\theta, W)), \quad \theta \in \mathbb{R}^{\mathbb{N} \times k}, w \in \mathbb{R}^{\mathbb{N}},$$

be the centered loss function. Previously introduced PAC-Bayesian inequalities (see Lemma 17 on page 59) show that

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \exp \sup_{\theta \in \ell_2^k} \left[n\lambda (\mathbb{P}_W - \bar{\mathbb{P}}_W) (\rho_{\theta'|\theta} - \rho_{\theta'|\theta}^2) f(\theta', W) \right. \right. \\ \left. \left. - n\rho_{\theta'|\theta} \left[\log \left(\mathbb{P}_W \left[\exp \left(-\lambda (\delta_{\theta''|\theta'} - \rho_{\theta''|\theta'}) \bar{f}(\theta'', W) \right) \right] \right) \right] - \frac{\beta \|\theta\|^2}{2} \right] \right\} \leq 1.$$

Apply Jensen's inequality and devide by $n\lambda$ to get

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \ell_2^k} \left[(\mathbb{P}_W - \bar{\mathbb{P}}_W) (\rho_{\theta'|\theta} - \rho_{\theta'|\theta}^2) f(\theta', W) \right. \right. \\ \left. \left. - \lambda^{-1} \rho_{\theta'|\theta} \left[\log \left(\mathbb{P}_W \left[\exp \left(-\lambda (\delta_{\theta''|\theta'} - \rho_{\theta''|\theta'}) \bar{f}(\theta'', W) \right) \right] \right) \right] - \frac{\beta \|\theta\|^2}{2n\lambda} \right] \right\} \leq 0.$$

Remark that

$$(\delta_{\theta'|\theta} - \rho_{\theta'|\theta}) f(\theta', W) = \rho_{\theta'|\theta} \left(\min_j \langle \theta_j, W \rangle - \min_j \langle \theta'_j, W \rangle \right) \\ \leq \rho_{\theta'|\theta} \left(\max_j \langle \theta_j - \theta'_j, W \rangle \right) \leq \sqrt{2 \log(k)/\beta} \|W\|_{\infty},$$

where the last inequality follows from the classical maximal inequality of the expectation of the maximum of i.i.d Gaussian random variables (see section 2.5 in [BLM13]).

Writing a symmetric inequality for the opposite, we deduce that

$$\left| (\delta_{\theta'|\theta} - \rho_{\theta'|\theta}) \bar{f}(\theta', W) \right| \leq 2\sqrt{2 \log(k)/\beta} \|W\|_{\infty}. \quad (4.18)$$

Using Hoeffding's inequality, and considering a closed bounded subset $\Theta \subset \ell_2^k$, we deduce that

$$\mathbb{P}_{W_1, \dots, W_n} \left[\sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) (\rho_{\theta'|\theta} - \rho_{\theta'|\theta}^2) f(\theta', W) \right] \leq \frac{4\lambda}{\beta} \log(k) \|W\|_{\infty}^2 + \frac{\beta \|\Theta\|^2}{2n\lambda}.$$

In view of this, choose

$$\lambda = \frac{\beta \|\Theta\|}{\sqrt{8n \log(k)} \|W\|_{\infty}}$$

and define

$$F = \|W\|_\infty \|\Theta\| \sqrt{\frac{8 \log(k)}{n}}. \quad (4.19)$$

For any integer h ,

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left[(\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\rho_{\theta'|\theta}^{2^h} - \rho_{\theta'|\theta}^{2^{h+1}} \right) f(\theta', W) \right] \right\} \leq F.$$

Summing up for $h = 0$ to $H - 1$, where H is to be chosen later, and exchanging \sum_h and \sup_θ , we deduce that

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left[(\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\rho_{\theta'|\theta} - \rho_{\theta'|\theta}^{2^H} \right) f(\theta', W) \right] \right\} \leq HF.$$

As we are interested in bounding $(\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W)$, there remains to upper bound

$$(\mathbb{P}_W - \bar{\mathbb{P}}_W) (\delta_{\theta'|\theta} - \rho_{\theta'|\theta}) f(\theta', W) \quad (4.20)$$

$$\text{and } (\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta}^{2^H} f(\theta', W), \quad (4.21)$$

or with a change of notation

$$(\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta} f(\theta', W). \quad (4.22)$$

An almost sure bound for (4.20) is provided by equation (4.18). To bound (4.22), introduce the influence function

$$\psi(x) = \begin{cases} \log(1 + x + x^2/2), & x \geq 0, \\ -\log(1 - x + x^2/2), & x \leq 0 \end{cases} \quad (4.23)$$

and put

$$\tilde{f}(\theta, W) = f(\theta, W) - \frac{a+b}{2}.$$

Decompose (4.22) into

$$(\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta} f(\theta', W) = \rho_{\theta'|\theta} \left[\mathbb{P}_W \tilde{f}(\theta', W) - \bar{\mathbb{P}}_W \left(\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] \right) \right] \quad (4.24)$$

$$+ \rho_{\theta'|\theta} \bar{\mathbb{P}}_W \left[\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] - \tilde{f}(\theta', W) \right]. \quad (4.25)$$

In order to bound (4.25), note that from lemma 7.2 in [Cat12]

$$|x - \psi(x)| \leq \frac{x^2}{4(1 + \sqrt{2})}, \quad x \in \mathbb{R}. \quad (4.26)$$

Therefore, from the inequality $(a + b)^2 \leq 2a^2 + 2b^2$ and the properties of the variance, \mathbb{P}_W almost surely,

$$\rho_{\theta'|\theta} \left[\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] - \tilde{f}(\theta', W) \right] \leq \frac{\lambda}{4(1 + \sqrt{2})} \rho_{\theta'|\theta} [\tilde{f}(\theta', W)^2]$$

$$\leq \frac{\lambda}{2(1+\sqrt{2})} \left[\left(\min_j \langle \theta_j, W \rangle - (a+b)/2 \right)^2 + \rho_{\theta' \mid 0} \left(\max_j \langle \theta'_j, W \rangle^2 \right) \right].$$

At this point, it remains to bound the variance term $\rho_{\theta' \mid 0} \left(\max_j \langle \theta'_j, W \rangle^2 \right)$. Let us remark that

$$\rho_{\theta' \mid 0} \circ (\theta'_j \mapsto \langle \theta'_j, W \rangle)^{-1} = \mathbf{N}(0, \|W\|^2/\beta).$$

Next, we need the following maximal inequality.

LEMMA 37 *Let $(\varepsilon_1, \dots, \varepsilon_k)$ be a sequence of Gaussian random variables such that $\varepsilon_j \sim \mathbf{N}(0, \sigma^2)$. We have*

$$\mathbb{E} \left(\max_{1 \leq j \leq k} \varepsilon_j^2 \right) \leq 2\sigma^2 \log(ke).$$

PROOF.

$$\begin{aligned} \mathbb{E} \left(\max_{1 \leq j \leq k} \varepsilon_j^2 \right) &= \int_{\mathbb{R}_+} \mathbb{P} \left(\max_{1 \leq j \leq k} \varepsilon_j^2 > t \right) dt \\ &\leq \int_{\mathbb{R}_+} \min \left\{ \sum_{j=1}^k \mathbb{P}(\varepsilon_j^2 > t), 1 \right\} dt \leq \int_{\mathbb{R}_+} \min \left\{ 2k \mathbb{P}(\varepsilon_1 > \sqrt{t}), 1 \right\} dt \\ &\leq \int_{\mathbb{R}_+} \min \left\{ k \exp \left(-\frac{t}{2\sigma^2} \right), 1 \right\} dt \leq 2\sigma^2 \log(k) + \int_{2\sigma^2 \log(k)}^{+\infty} k \exp \left(-\frac{t}{2\sigma^2} \right) dt \\ &\leq 2\sigma^2 \log(k) + 2\sigma^2 = 2\sigma^2 \log(ke). \end{aligned}$$

□

Accordingly, we obtain \mathbb{P}_W almost surely,

$$\begin{aligned} \rho_{\theta' \mid \theta} \left[\lambda^{-1} \psi \left[\lambda \tilde{f}(\theta', W) \right] - \tilde{f}(\theta', W) \right] \\ \leq \frac{\lambda}{2(1+\sqrt{2})} \left[\left(\min_j \langle \theta_j, W \rangle - (a+b)/2 \right)^2 + \rho_{\theta' \mid 0} \left(\max_j \langle \theta'_j, W \rangle^2 \right) \right] \\ \leq \frac{\lambda}{2(1+\sqrt{2})} \left[(b-a)^2/4 + 2 \log(ek) \|W\|_\infty^2 / \beta \right]. \quad (4.27) \end{aligned}$$

The right-hand side of this inequality provides an almost sure upper bound for (4.25). To bound (4.24), or rather the expectation of an exponential moment of (4.24), we can write a PAC-Bayesian bound using the influence function ψ .

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \exp \left[-n \lambda \rho_{\theta' \mid \theta} \bar{\mathbb{P}}_W \left(\lambda^{-1} \psi \left[\lambda \tilde{f}(\theta', W) \right] \right) \right. \right. \\ \left. \left. - n \rho_{\theta' \mid \theta} \left[\log \left(\mathbb{P}_W \left[\exp \left(\psi \left[-\lambda \tilde{f}(\theta', W) \right] \right) \right] \right) \right] - \frac{\beta \|\theta\|^2}{2} \right] \right\} \leq 1. \end{aligned}$$

Remarking that

$$\psi(x) \leq \log(1 + x + x^2/2), \quad x \in \mathbb{R},$$

and removing the exponential according to Jensen's inequality, we obtain

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left[\rho_{\theta' \mid \theta} \left[\mathbb{P}_W \left(\tilde{f}(\theta', W) \right) - \bar{\mathbb{P}}_W \left(\lambda^{-1} \psi [\lambda \tilde{f}(\theta', W)] \right) \right] - \frac{\lambda}{2} \rho_{\theta' \mid \theta} \left[\mathbb{P}_W \left(\tilde{f}(\theta', W)^2 \right) \right] \right] \right\} \leq \frac{\beta \|\Theta\|^2}{2}.$$

Using the same maximal inequality in relation to the maximum of squared Gaussian random variables, used in the course of equation (4.27) to bound the variance term, we get

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \rho_{\theta' \mid \theta} \left[\mathbb{P}_W \left(\tilde{f}(\theta', W) \right) - \bar{\mathbb{P}}_W \left(\lambda^{-1} \psi [\lambda \tilde{f}(\theta', W)] \right) \right] \right\} \\ \leq \lambda \left[(b-a)^2/4 + 2 \log(ek) \|W\|_\infty^2 / \beta \right] + \frac{\beta \|\Theta\|^2}{2n\lambda}. \end{aligned}$$

This provides an upper bound for (4.24). Combining it with the upper bound for (4.25) gives an upper bound for (4.22) that reads

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left(\mathbb{P}_W - \bar{\mathbb{P}}_W \right) \rho_{\theta' \mid \theta} f(\theta', W) \right\} \\ \leq \frac{(\sqrt{2}+1)\lambda}{2} \left[(b-a)^2/4 + 2 \log(ek) \|W\|_\infty^2 / \beta \right] + \frac{\beta \|\Theta\|^2}{2n\lambda}. \end{aligned}$$

Choosing

$$\lambda = \sqrt{\frac{4\beta \|\Theta\|^2}{(\sqrt{2}+1) \left[(b-a)^2 + 8 \log(ek) \|W\|_\infty^2 / \beta \right] n}}$$

gives

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left(\mathbb{P}_W - \bar{\mathbb{P}}_W \right) \rho_{\theta' \mid \theta} f(\theta', W) \right\} \\ \leq \tilde{F}(\beta) \stackrel{\text{def}}{=} \sqrt{\frac{(\sqrt{2}+1) \left(\beta(b-a)^2 + 8 \log(ek) \|W\|_\infty^2 \right) \|\Theta\|^2}{4n}}. \end{aligned}$$

Putting everything together,

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left(\mathbb{P}_W - \bar{\mathbb{P}}_W \right) f(\theta, W) \right\} \leq 2\sqrt{2 \log(k) / \beta} \|W\|_\infty + \tilde{F}(2^{-H}\beta) + HF,$$

where F is defined by equation (4.19) on page 93.

Let us choose $\beta = 2n\|\Theta\|^{-2}$ and $H = \lfloor \log(n/k) / \log(2) \rfloor$, so that

$$2^{-H}\beta \leq 4k\|\Theta\|^{-2}.$$

We get

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left(\mathbb{P}_W - \bar{\mathbb{P}}_W \right) f(\theta, W) \right\} \\ \leq \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8 \log(k)}{n}} + 2\sqrt{\frac{\log(k)}{n}} \right) \|\Theta\| \|W\|_\infty \end{aligned}$$

$$+ \sqrt{\frac{(\sqrt{2}+1)\left(k(b-a)^2 + 2\log(ek)\|W\|_\infty^2\|\Theta\|^2\right)}{n}}.$$

The upper deviations from this mean are controled by the extension of Hoeffding's bound called the bounded difference inequality (see section 6.1 and theorem 6.2 in [BLM13]). It gives with probability at least $1 - \delta$

$$\sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W) \leq \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W) \right\} + \sqrt{\frac{2\log(\delta^{-1})}{n}}(b-a).$$

This proves the first statement of the lemma. To get the second one, add to the previous inequality

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ (\bar{\mathbb{P}}_W - \mathbb{P}_W) f(\theta^*, W) \right\} = 0$$

to get

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) (f(\theta, W) - f(\theta^*, W)) \right\} \\ \leq \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8\log(k)}{n}} + 2\sqrt{\frac{\log(k)}{n}} \right) \|\Theta\| \|W\|_\infty \\ + \sqrt{\frac{(\sqrt{2}+1)\left(k(b-a)^2 + 2\log(ek)\|W\|_\infty^2\|\Theta\|^2\right)}{n}}. \end{aligned}$$

and apply the bounded difference inequality to get the deviations. \square

Let us first apply this lemma to the information k -means problem.

PROPOSITION 38 *Assume that*

$$\text{ess sup}_X \left(\int p_X^2 d\nu \right) < +\infty \quad \text{and} \quad \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right) < +\infty.$$

Consider the information radius

$$R = \inf_{q \in \mathbb{L}_{+,1}^1(\nu)} \text{ess sup}_X \mathcal{K}(q, p_X)$$

and the bounds

$$\begin{aligned} B &= \text{ess sup}_X \left(\int p_X^2 d\nu \right)^{1/2} \exp(R) \\ \text{and } C &= \text{ess sup}_X \left(\int \log(p_X)^2 d\nu \right)^{1/2}. \end{aligned}$$

Introduce the parameter space

$$\Omega_B = \left\{ q \in \mathbb{L}_{+,1}^1(\nu) \cap \mathbb{L}^2(\nu) : \int q^2 d\nu \leq B^2 \right\}.$$

Given (X_1, \dots, X_n) , a sample made of n independent copies of X , with probability at least $1 - \delta$, for any $q \in \Omega_B^k$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) &\leq \frac{1}{n} \sum_{i=1}^n \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_{X_i}) \\ &+ \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2 \sqrt{\frac{k \log(k)}{n}} \right. \\ &\quad \left. + \sqrt{\frac{(\sqrt{2}+1)k(3+2\log(k))}{n}} + \sqrt{\frac{\log(\delta^{-1})}{2n}} \right) (BC + 2\log(B)). \end{aligned}$$

Consider an empirical risk minimizer $\hat{q}(X_1, \dots, X_n) \in \mathbf{Q}_B^k$ satisfying

$$\hat{q} \in \arg \min_{q \in \mathbf{Q}_B^k} \overline{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right).$$

With probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(\hat{q}, p_X) \mid X_1, \dots, X_n \right) &\leq \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right) \\ &+ \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2 \sqrt{\frac{k \log(k)}{n}} \right. \\ &\quad \left. + \sqrt{\frac{(\sqrt{2}+1)k(3+2\log(k))}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}} \right) (BC + 2\log(B)). \end{aligned}$$

PROOF. Consider the parametrization used in the proof of Proposition 20 on page 67, that is $\theta = (q, \mathcal{K}(q, 1)) \in H$ and $W = (-\log(p_X), \mu^{-1}) \in H$. This puts the information k -means risk in a suitable form to apply Lemma 36 on page 90. Taking into account that $\|\Theta\| \|W\|_\infty \leq \sqrt{k}(BC + 2\log(B))$ for a suitable choice of μ ends the proof. \square

We can also apply the previous lemma to the Euclidean k -means algorithm.

PROPOSITION 39 Consider a random vector X in a separable Hilbert space H . Let (X_1, \dots, X_n) be a sample made of n independent copies of X . Consider the ball of radius B

$$\mathcal{B} = \{x \in H : \|x\| \leq B\}.$$

Assume that $\mathbb{P}(X \in \mathcal{B}) = 1$ and consider an estimator

$$\hat{\mu} \in \arg \min_{\mu \in H^k} \overline{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \mu_j\|^2 \right).$$

Assume that $n \geq 2k$ and $k \geq 2$. With probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \hat{\mu}_j\|^2 \right) &\leq \inf_{\mu \in H^k} \mathbb{P}_X \left(\|X - \mu_j\|^2 \right) \\ &+ B^2 \log \left(\frac{n}{k} \right) \sqrt{\frac{k \log(k)}{n}} \left(\underbrace{\frac{6\sqrt{2}}{\log(2)} + \frac{6}{\log(n/k)} + \frac{1}{\log(n/k)} \sqrt{\frac{2(\sqrt{2}+1)(17+9\log(k))}{\log(k)}}}_{\leq 12.3} \right) \end{aligned}$$

$$+ 4B^2 \sqrt{\frac{2 \log(\delta^{-1})}{n}}.$$

Consequently, with probability at least $1 - \delta$,

$$\begin{aligned} \mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \hat{\mu}_j\|^2 \right) &\leq \inf_{\mu \in H^k} \mathbb{P}_X \left(\|X - \mu_j\|^2 \right) \\ &\quad + 16 B^2 \log \left(\frac{n}{k} \right) \sqrt{\frac{k \log(k)}{n}} + 4 B^2 \sqrt{\frac{2 \log(\delta^{-1})}{n}}. \end{aligned}$$

In expectation

$$\mathbb{P}_{X_1, \dots, X_n} \left[\mathbb{P}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|X - \hat{\mu}_j\|^2 \right) \right] \leq \inf_{\mu \in H^k} \mathbb{P}_X \left(\|X - \mu_j\|^2 \right) + 16 B^2 \log \left(\frac{n}{k} \right) \sqrt{\frac{k \log(k)}{n}}.$$

In conclusion a chaining argument yields a dimension free non asymptotic generalization bound that decreases in expectation as $\sqrt{k/n}$ up to logarithmic factors.

PROOF. Remark that

$$\hat{\mu} \in \arg \min_{\mu \in \mathcal{B}^k} \overline{\mathbb{P}}_X \left(\min_{j \in \llbracket 1, k \rrbracket} \|\mu_j\|^2 - 2 \langle X, \mu_j \rangle \right)$$

and apply the lemma to $W = (-2X, \gamma B) \in H \times \mathbb{R}$ and $\theta_j = (\mu_j, \gamma^{-1} \|\mu_j\|^2 B^{-1})$. Note that

$$\|W\|^2 \|\theta_j\|^2 \leq B^4 (4 + \gamma^2) (1 + \gamma^{-2}) = B^4 (5 + \gamma^2 + 4\gamma^{-2}).$$

Choose the optimal value $\gamma^2 = 2$ to get

$$\|W\|^2 \|\Theta\|^2 \leq 9kB^4$$

Remark also that

$$\langle \theta_j, W \rangle \in [-B^2, +3B^2].$$

This gives the first statement of the proposition, according to Lemma 36 on page 90. Since $4B^2$ is a trivial bound, we have now to prove that for any $k \geq 2$ and any $n \geq 2k$,

$$\begin{aligned} \min \left\{ 4, \log \left(\frac{n}{k} \right) \sqrt{\frac{k \log(k)}{n}} \left(\frac{6\sqrt{2}}{\log(2)} + \frac{6}{\log(n/k)} + \frac{1}{\log(n/k)} \sqrt{\frac{2(\sqrt{2}+1)(17+9\log(k))}{\log(k)}} \right) \right\} \\ \leq 16 \log \left(\frac{n}{k} \right) \sqrt{\frac{k \log(k)}{n}}. \end{aligned}$$

Putting $a = \frac{6\sqrt{2}}{\log(2)}$, $b = 16$, $\rho = n/k$,

$$\begin{aligned} \eta &= 6 + \sqrt{\frac{2(\sqrt{2}+1)(17+9\log(k))}{\log(k)}}, \\ f(\rho, k) &= \sqrt{\log(k)/\rho} (a \log(\rho) + \eta(k)) \\ \text{and } g(\rho, k) &= b \sqrt{\log(k)/\rho} \log(\rho), \end{aligned}$$

we have to prove that

$$\min\{4, f(\rho, k)\} \leq g(\rho, k), \quad \rho \geq 2, k \geq 2.$$

In other words, we have to prove that, when $g(\rho, k) < f(\rho, k)$, then $g(\rho, k) \geq 4$. This can also be written as

$$g(\rho, k) \geq 4, \quad \min\{\rho, k\} \geq 2, \quad g(\rho, k) < f(\rho, k).$$

According to the definitions, this is also equivalent to

$$\log(\rho) - 2 \log(\log(\rho)) \leq 2 \log(b/4) + \log(\log(k)), \quad \min\{\rho, k\} \geq 2, \quad (b-a) \log(\rho) \leq \eta(k).$$

Since η is decreasing and since $k \mapsto \log(\log(k))$ is increasing, if the statement is true for $k = 2$, it is true for any $k \geq 2$. Thus we have to prove that

$$\log(\rho) - 2 \log(\log(\rho)) \leq 2 \log(b/4) + \log(\log(2)), \quad \log(2) \leq \log(\rho) \leq \eta(2)/(b-a).$$

Putting $\xi = \log(\rho)$, we have to prove that

$$\xi - 2 \log(\xi) \leq 2 \log(b/4) + \log(\log(2)), \quad \log(2) \leq \xi \leq \eta(2)/(b-a).$$

Since $\xi \mapsto \xi - 2 \log(\xi)$ is convex, it is enough to check the inequality at the two ends of the interval, that is when $\xi \in \{\log(2), \eta(2)/(b-a)\}$, which can be done numerically. More precisely, we have to check that

$$\begin{aligned} & 2 \log(b/4) + \log(\log(2)) \\ & - \max\left\{\log(2) - 2 \log(\log(2)), \eta(2)/(b-a) - 2 \log[\eta(2)/(b-a)]\right\} \geq 0, \end{aligned}$$

and we get numerically that the left-hand side is larger than the minimum of 0.9 and 0.6. \square

4.5.2. FASTER BOUNDS FOR BOTH THE BOUNDED INFORMATION k -MEANS AND THE BOUNDED k -MEANS CRITERION. Let us derive faster generalization bounds concerning the bounded information k -means and the bounded k -means criterion, that are defined respectively in section 4.2 on page 71 and 4.3 on page 76.

PROPOSITION 40 *Let X_1, \dots, X_n be made of n independent copies of X . For any family of k probability densities $q \in (\mathbb{L}_{+,1}^1(\nu))^k$, consider the bounded criterion $\mathbf{C}(q)$ defined in Lemma 23 on page 72 and its empirical counterpart*

$$\bar{\mathbf{C}}(q) = 1 - \bar{\mathbb{P}}_X \left(\exp \left[- \min_{j \in \llbracket 1, k \rrbracket} \mathcal{K}(q_j, p_X) \right] \right).$$

Define the bounds B and C and the set \mathcal{B} as in Lemma 25 on page 75. Assume that $k \geq 2$ and $n \geq 2k$. With probability at least $1 - \delta$, for any $q \in \mathcal{B}^k$,

$$\begin{aligned}\mathcal{C}(q) \leq \bar{\mathcal{C}}(q) + \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2\sqrt{\frac{k \log(k)}{n}} \right) \exp(BC) \\ + \sqrt{\frac{(\sqrt{2}+1)k [1 + 2\log(ek) \exp(2BC)]}{n}} + \sqrt{\frac{\log(\delta^{-1})}{2n}}.\end{aligned}$$

Let $q^* \in \mathcal{B}^k$ be a non random family of centers, and assume that $k \geq 2$ and $n \geq 2k$. With probability at least $1 - \delta$, for any $q \in \mathcal{B}^k$,

$$\begin{aligned}\mathcal{C}(q) - \mathcal{C}(q^*) \leq \bar{\mathcal{C}}(q) - \bar{\mathcal{C}}(q^*) + \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2\sqrt{\frac{k \log(k)}{n}} \right) \exp(BC) \\ + \sqrt{\frac{(\sqrt{2}+1)k [1 + 2\log(ek) \exp(2BC)]}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}}.\end{aligned}$$

Consequently, if

$$\hat{q}(X_1, \dots, X_n) \in \arg \min_{q \in \mathcal{B}^k} \bar{\mathcal{C}}(q),$$

with probability at least $1 - \delta$,

$$\begin{aligned}\mathcal{C}(\hat{q}) \leq \inf_{q \in (\mathbb{L}_{+,1}^1(\nu))^k} \mathcal{C}(q) + \left(\frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2\sqrt{\frac{k \log(k)}{n}} \right) \exp(BC) \\ + \sqrt{\frac{(\sqrt{2}+1)k [1 + 2\log(ek) \exp(2BC)]}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}}.\end{aligned}$$

One can get also a similar bound for the expected excess risk.

PROOF. Remember from section 4.2 on page 71 that the risk $\mathcal{C}(q)$ can be expressed as

$$\mathcal{C}(q) = 1 - \mathbb{P}_W \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) = 1 + \mathbb{P}_W \left(\min_{j \in \llbracket 1, k \rrbracket} \langle -\theta_j, W \rangle \right),$$

where $\theta_j = \exp(-\mathcal{K}(q_j, 1)) \Psi(\mu q_j)$ and $W = \Psi(\mu^{-1} \log(p_X))$. Therefore we can apply Lemma 36 on page 90, using the inequality $\|\theta_j\|^2 \|W\|^2 \leq \exp(2BC)$. \square

Now, let us look at the bounded k -means criterion.

PROPOSITION 41 Consider the same situation as in Proposition 27 on page 77. Consider any $k \geq 2$ and any $n \geq 2k$. With probability at least $1 - \delta$, for any $\mu \in \mathbb{R}^{d \times k}$,

$$\begin{aligned}\mathcal{C}(\mu) \leq \bar{\mathcal{C}}(\mu) + \frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2\sqrt{\frac{k \log(k)}{n}} \\ + \sqrt{\frac{(\sqrt{2}+1)k (3 + 2\log(k))}{n}} + \sqrt{\frac{\log(\delta^{-1})}{2n}}.\end{aligned}$$

For any non random family of centers $\mu^* \in \mathbb{R}^{d \times k}$, with probability at least $1 - \delta$, for any $\mu \in \mathbb{R}^{d \times k}$,

$$\begin{aligned} \mathcal{C}(\mu) - \mathcal{C}(\mu^*) &\leq \bar{\mathcal{C}}(\mu) - \bar{\mathcal{C}}(\mu^*) + \frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2\sqrt{\frac{k \log(k)}{n}} \\ &\quad + \sqrt{\frac{(\sqrt{2}+1)k(3+2\log(k))}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}}. \end{aligned}$$

Consequently, if

$$\hat{\mu}(X_1, \dots, X_n) \in \arg \min_{\mu \in \mathbb{R}^{d \times k}} \bar{\mathcal{C}}(\mu),$$

with probability at least $1 - \delta$,

$$\begin{aligned} \mathcal{C}(\hat{\mu}) &\leq \inf_{\mu \in \mathbb{R}^{d \times k}} \mathcal{C}(\mu) + \frac{\log(n/k)}{\log(2)} \sqrt{\frac{8k \log(k)}{n}} + 2\sqrt{\frac{k \log(k)}{n}} \\ &\quad + \sqrt{\frac{(\sqrt{2}+1)k(3+2\log(k))}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}}. \end{aligned}$$

PROOF. From section 4.3 on page 76, we can express the risk as

$$\mathcal{C}(\mu) = 1 - \mathbb{P}_X \left(\max_{j \in \llbracket 1, k \rrbracket} \langle \theta_j, W \rangle \right) = 1 + \mathbb{P} \left(\min_{j \in \llbracket 1, k \rrbracket} \langle -\theta_j, W \rangle \right),$$

where $\theta_j = \Psi(\mu_j)$ and $W = \Psi(X)$, and where Ψ is the feature map associated to the Gaussian kernel. Accordingly, θ_j and W belongs to the unit ball in the corresponding reproducing kernel Hilbert space H , so that $\|W\|_\infty \leq 1$ and $\|\Theta\| \leq \sqrt{k}$. \square

4.5.3. FASTER BOUNDS FOR INFORMATION FRAGMENTATION. Similarly to what we have done previously, we can also obtain faster bounds for information fragmentation. However, we need first to establish the equivalent of Lemma 36 on page 90 for the information fragmentation risk.

LEMMA 42 *Let $W = (W_j, 1 \leq j \leq k)$ be a random vector in the product H^k , where H is a separable Hilbert space (that we can take as being ℓ_2 if we want). Consider a sample $(W^{(1)}, \dots, W^{(n)})$ made of n independent copies of W . Consider a bounded parameter set $\Theta \subset H^k$ and a set \mathcal{T} of subsets of $\llbracket 1, k \rrbracket$. Assume that*

$$\mathbb{P}_W \left(\sum_{j \in A} \langle \theta_j, W_j \rangle \in [a, b], A \in \mathcal{T}, \theta \in \Theta \right) = 1.$$

Consider the risk

$$\mathcal{C}(\theta) = \mathbb{P}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right), \quad \theta \in \Theta,$$

and its empirical counterpart

$$\bar{\mathcal{C}}(\theta) = \bar{\mathbb{P}}_W \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle \right), \quad \theta \in \Theta.$$

Put

$$K(\mathcal{T}) = \max_{A \in \mathcal{T}} |A|, \quad \|\Theta\| = \sup_{\theta \in \Theta} \left(\sum_{j=1}^k \|\theta_j\|^2 \right)^{1/2} \quad \text{and} \quad \|W\|_\infty = \max_{j \in \llbracket 1, k \rrbracket} \text{ess sup}_{\mathbb{P}_W} \|W_j\|.$$

Let \mathcal{S} be a positive real parameter. Assume that $|\mathcal{T}| \geq 2$ and that $n \geq 2\mathcal{S}K(\mathcal{T})$. For any $\delta \in]0, 1[$, with probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\begin{aligned} \mathcal{C}(\theta) \leq \bar{\mathcal{C}}(\theta) + & \left(\frac{\log(n\mathcal{S}^{-1}K(\mathcal{T})^{-1})}{\log(2)} \sqrt{\frac{8K(\mathcal{T})\log(|\mathcal{T}|)}{n}} + 2\sqrt{\frac{K(\mathcal{T})\log(|\mathcal{T}|)}{n}} \right) \|\Theta\| \|W\|_\infty \right. \\ & \left. + \sqrt{\frac{(\sqrt{2}+1)(\mathcal{S}(b-a)^2 + 2\log(e|\mathcal{T}|)\|W\|_\infty^2\|\Theta\|^2)K(\mathcal{T})}{n}} + \sqrt{\frac{\log(\delta^{-1})}{2n}}(b-a). \right) \end{aligned}$$

If $\theta^* \in \Theta$ is a non random value of the parameter, with probability at least $1 - \delta$, for any $\theta \in \Theta$,

$$\begin{aligned} \mathcal{C}(\theta) - \mathcal{C}(\theta^*) \leq & \bar{\mathcal{C}}(\theta) - \bar{\mathcal{C}}(\theta^*) \\ & + \left(\frac{\log(n\mathcal{S}^{-1}K(\mathcal{T})^{-1})}{\log(2)} \sqrt{\frac{8K(\mathcal{T})\log(|\mathcal{T}|)}{n}} + 2\sqrt{\frac{K(\mathcal{T})\log(|\mathcal{T}|)}{n}} \right) \|\Theta\| \|W\|_\infty \\ & + \sqrt{\frac{(\sqrt{2}+1)(\mathcal{S}(b-a)^2 + 2\log(e|\mathcal{T}|)\|W\|_\infty^2\|\Theta\|^2)K(\mathcal{T})}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}}(b-a). \end{aligned}$$

Consequently, if

$$\hat{\theta} \in \arg \min_{\theta \in \Theta} \bar{\mathcal{C}}(\theta),$$

with probability at least $1 - \delta$,

$$\begin{aligned} \mathcal{C}(\hat{\theta}) \leq \inf_{\theta \in \Theta} \mathcal{C}(\theta) + & \left(\frac{\log(n\mathcal{S}^{-1}K(\mathcal{T})^{-1})}{\log(2)} \sqrt{\frac{8K(\mathcal{T})\log(|\mathcal{T}|)}{n}} + 2\sqrt{\frac{K(\mathcal{T})\log(|\mathcal{T}|)}{n}} \right) \|\Theta\| \|W\|_\infty \\ & + \sqrt{\frac{(\sqrt{2}+1)(\mathcal{S}(b-a)^2 + 2\log(e|\mathcal{T}|)\|W\|_\infty^2\|\Theta\|^2)K(\mathcal{T})}{n}} + \sqrt{\frac{2\log(\delta^{-1})}{n}}(b-a) \end{aligned}$$

In expectation

$$\begin{aligned} \mathbb{P}_{W^{(1)}, \dots, W^{(n)}} \left(\mathcal{C}(\hat{\theta}) \right) \leq & \inf_{\theta \in \Theta} \mathcal{C}(\theta) + \left(\frac{\log(n\mathcal{S}^{-1}K(\mathcal{T})^{-1})}{\log(2)} \sqrt{\frac{8K(\mathcal{T})\log(|\mathcal{T}|)}{n}} + 2\sqrt{\frac{K(\mathcal{T})\log(|\mathcal{T}|)}{n}} \right) \|\Theta\| \|W\|_\infty \\ & + \sqrt{\frac{(\sqrt{2}+1)(\mathcal{S}(b-a)^2 + 2\log(e|\mathcal{T}|)\|W\|_\infty^2\|\Theta\|^2)K(\mathcal{T})}{n}}. \end{aligned}$$

PROOF. We follow here the same arguments as in the proof of Lemma 36 on page 90, so that we consider the same perturbation

$$\rho_{\theta'|\theta} = \mathbb{P}_{\theta_i + \beta^{-1/2}\varepsilon_i, i \in \mathbb{N}},$$

along with

$$\rho_{\theta'|\theta} = \bigotimes_{j=1}^k \rho_{\theta'_j|\theta_j},$$

in the case where θ and $\theta' \in \mathbb{R}^{\mathbb{N} \times k}$.

Taking a basis, we can assume without loss of generality that $W = (W_1, \dots, W_k)$ is a random vector in the separable Hilbert space $(\ell_2)^k \subset \mathbb{R}^{\mathbb{N} \times k}$. Define the loss function

$$f(\theta, w) = \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, w_j \rangle, \quad \theta \in \mathbb{R}^{\mathbb{N} \times k}, w = (w_1, \dots, w_k) \in \mathbb{R}^{\mathbb{N} \times k},$$

where the scalar product is extended beyond ℓ_2 as in equation (4.4) on page 61. Introduce

$$\bar{f}(\theta, w) = f(\theta, w) - \mathbb{P}_W(f(\theta, W)), \quad \theta \in \mathbb{R}^{\mathbb{N} \times k}, w \in \mathbb{R}^{\mathbb{N} \times k}.$$

Since

$$\min_A a_A - \min_A b_A \leq \max_A (a_A - b_A),$$

we see that

$$\begin{aligned} (\delta_{\theta'|\theta} - \rho_{\theta'|\theta}) f(\theta', W) &= \rho_{\theta'|\theta} \left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle - \min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta'_j, W_j \rangle \right) \\ &\leq \rho_{\theta'|\theta} \left(\max_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j - \theta'_j, W_j \rangle \right). \end{aligned}$$

To bound the right-hand side of the previous inequality, we need a maximal inequality, knowing that

$$\rho_{\theta'|\theta} \circ \left(\theta'_j \mapsto \sum_{j \in A} \langle \theta_j - \theta'_j, W_j \rangle \right)^{-1} = \mathcal{N}\left(0, \sum_{j \in A} \|W_j\|^2 / \beta\right).$$

LEMMA 43 *Let \mathcal{T} be a set of subsets of $\llbracket 1, k \rrbracket$ and let $(\varepsilon_A)_{A \in \mathcal{T}}$ be a random process with marginal distributions $\mathbb{P}_{\varepsilon_A} = \mathcal{N}(0, \sigma_A^2)$, $A \in \mathcal{T}$. The expectation of its maximum is bounded by*

$$\mathbb{P}_\varepsilon \left(\max_{A \in \mathcal{T}} \varepsilon_A \right) \leq \sqrt{2 \log(|\mathcal{T}|) \max_{A \in \mathcal{T}} \sigma_A^2}.$$

PROOF. By Jensen's inequality, for any $t > 0$

$$\begin{aligned} \mathbb{P}_\varepsilon \left(\max_{A \in \mathcal{T}} \varepsilon_A \right) &\leq \frac{1}{t} \log \mathbb{P}_\varepsilon \left(\max_{A \in \mathcal{T}} \exp(t \varepsilon_A) \right) \\ &\leq \frac{1}{t} \log \left(\sum_{A \in \mathcal{T}} \mathbb{P}_\varepsilon \left(\exp(t \varepsilon_A) \right) \right) \leq \frac{1}{t} \log \left(|\mathcal{T}| \exp \left(t^2 \max_{A \in \mathcal{T}} \sigma_A^2 / 2 \right) \right) \\ &\leq \frac{\log(|\mathcal{T}|)}{t} + \frac{t}{2} \max_{A \in \mathcal{T}} \sigma_A^2. \end{aligned}$$

Minimizing over t ends the proof. \square

Applying this lemma in the particular case where $\sigma_A^2 = \sum_{j \in A} \|W_j\|^2 / \beta$, and remarking that $\max_{A \in \mathcal{T}} \sigma_A^2 \leq K(\mathcal{T}) \|W\|_\infty^2 / \beta$, we get

$$(\delta_{\theta'|\theta} - \rho_{\theta'|\theta}) f(\theta', W) \leq \rho_{\theta'|\theta} \left(\max_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j - \theta'_j, W_j \rangle \right) \leq \sqrt{2K(\mathcal{T}) \log(|\mathcal{T}|) / \beta} \|W\|_\infty.$$

Accordingly, the same bound holding true for the opposite,

$$\left| \left(\delta_{\theta'|\theta} - \rho_{\theta'|\theta} \right) \bar{f}(\theta', W) \right| \leq 2\sqrt{2K(\mathcal{T}) \log(|\mathcal{T}|)/\beta} \|W\|_\infty.$$

Following the same line of proof as in Lemma 36 on page 90, we obtain

$$\mathbb{P}_{W_1, \dots, W_n} \left[\sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) (\rho_{\theta'|\theta} - \rho_{\theta'|\theta}^2) f(\theta', W) \right] \leq \frac{4\lambda}{\beta} K(\mathcal{T}) \log(|\mathcal{T}|) \|W\|_\infty^2 + \frac{\beta \|\Theta\|^2}{2n\lambda}.$$

In the same way, choose

$$\lambda = \frac{\beta \|\Theta\|}{\sqrt{8nK(\mathcal{T}) \log(|\mathcal{T}|)} \|W\|_\infty}$$

and define

$$F = \|W\|_\infty \|\Theta\| \sqrt{\frac{8K(\mathcal{T}) \log(|\mathcal{T}|)}{n}}. \quad (4.28)$$

Following the PAC-Bayesian chaining strategy introduced previously, let us consider a sequence of measures $(\rho_{\theta'|\theta}^{2^h})_{h \in \llbracket 0, H-1 \rrbracket}$. For any integer $h \in \llbracket 0, H-1 \rrbracket$,

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left[(\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\rho_{\theta'|\theta}^{2^h} - \rho_{\theta'|\theta}^{2^{h+1}} \right) f(\theta', W) \right] \right\} \leq F.$$

Summing over h , we get

$$\begin{aligned} & \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left[\sum_{h=0}^{H-1} (\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\rho_{\theta'|\theta}^{2^h} - \rho_{\theta'|\theta}^{2^{h+1}} \right) f(\theta', W) \right] \right\} \\ & \leq \mathbb{P}_{W_1, \dots, W_n} \left\{ \sum_{h=0}^{H-1} \sup_{\theta \in \Theta} \left[(\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\rho_{\theta'|\theta}^{2^h} - \rho_{\theta'|\theta}^{2^{h+1}} \right) f(\theta', W) \right] \right\} \leq HF. \end{aligned}$$

Hence, simplifying the telescoping sums on the left-hand side yields

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \left[(\mathbb{P}_W - \bar{\mathbb{P}}_W) \left(\rho_{\theta'|\theta} - \rho_{\theta'|\theta}^{2^H} \right) f(\theta', W) \right] \right\} \leq HF.$$

As in the proof of Lemma 36 on page 90, since we are dealing with $(\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W)$, it remains to bound the following two quantities

$$(\mathbb{P}_W - \bar{\mathbb{P}}_W) (\delta_{\theta'|\theta} - \rho_{\theta'|\theta}) f(\theta', W) \quad (4.29)$$

$$\text{and } (\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta}^{2^H} f(\theta', W). \quad (4.30)$$

For the sake of simplicity, we can rewrite this last quantity as

$$(\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta} f(\theta', W), \quad (4.31)$$

after a change of notation. Using the influence function ψ introduced in [Cat12] and defined by equation (4.23) on page 93 to decompose

$$(\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta} f(\theta', W),$$

into a sub-Gaussian part and an other part representing extreme values, we get

$$\begin{aligned} (\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta'|\theta} f(\theta', W) &= \\ \rho_{\theta'|\theta} \left[\mathbb{P}_W \tilde{f}(\theta', W) - \bar{\mathbb{P}}_W \left(\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] \right) \right] &\quad (4.32) \\ + \rho_{\theta'|\theta} \bar{\mathbb{P}}_W \left[\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] - \tilde{f}(\theta', W) \right], &\quad (4.33) \end{aligned}$$

where $\tilde{f}(\theta, W)$ is defined in the same way as in the proof of 36 on page 90 as

$$\tilde{f}(\theta, W) = f(\theta, W) - \frac{a+b}{2}.$$

From inequality (4.26) on page 93, the inequality $(a+b)^2 \leq 2a^2 + 2b^2$, and the properties of the variance, we get

$$\begin{aligned} \rho_{\theta'|\theta} \left[\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] - \tilde{f}(\theta', W) \right] &\leq \frac{\lambda}{4(1+\sqrt{2})} \rho_{\theta'|\theta} [\tilde{f}(\theta', W)^2] \\ &\leq \frac{\lambda}{2(1+\sqrt{2})} \left[\left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle - (a+b)/2 \right)^2 + \rho_{\theta'|\theta} \left(\max_{A \in \mathcal{T}} \left(\sum_{j \in A} \langle \theta'_j, W_j \rangle \right)^2 \right) \right]. \end{aligned}$$

Next, we need to bound the expectation of the maximum of the squares of Gaussian random variables. Consider the following adaptation of Lemma 37 on page 94.

LEMMA 44 *Let \mathcal{T} be a set of subsets of $\llbracket 1, k \rrbracket$ and let $(\varepsilon_A)_{A \in \mathcal{T}}$ be a random process with marginal distributions $\mathbb{P}_{\varepsilon_A} = \mathcal{N}(0, \sigma_A^2)$, $A \in \mathcal{T}$. It satisfies*

$$\mathbb{E} \left(\max_{A \in \mathcal{T}} \varepsilon_A^2 \right) \leq 2 \left(\max_{A \in \mathcal{T}} \sigma_A^2 \right) \log(|\mathcal{T}|e).$$

PROOF. It is similar to Lemma 37 on page 94. Indeed,

$$\begin{aligned} \mathbb{P}_\varepsilon \left(\max_{A \in \mathcal{T}} \varepsilon_A^2 \right) &\leq \int_{\mathbb{R}_+} \min \left\{ \sum_{A \in \mathcal{T}} \exp \left(-\frac{t}{2\sigma_A^2} \right), 1 \right\} dt \\ &\leq \int_{\mathbb{R}_+} \min \left\{ |\mathcal{T}| \exp \left(-\frac{t}{2 \max_{A \in \mathcal{T}} \sigma_A^2} \right), 1 \right\} dt = 2 \left(\max_{A \in \mathcal{T}} \sigma_A^2 \right) \log(|\mathcal{T}|e). \end{aligned}$$

□

From this lemma, we obtain, \mathbb{P}_W almost surely,

$$\begin{aligned} \rho_{\theta'|\theta} \left[\lambda^{-1} \psi[\lambda \tilde{f}(\theta', W)] - \tilde{f}(\theta', W) \right] &\leq \frac{\lambda}{2(1+\sqrt{2})} \left[\left(\min_{A \in \mathcal{T}} \sum_{j \in A} \langle \theta_j, W_j \rangle - (a+b)/2 \right)^2 + \rho_{\theta'|\theta} \left(\max_{A \in \mathcal{T}} \left(\sum_{j \in A} \langle \theta'_j, W_j \rangle \right)^2 \right) \right] \\ &\leq \frac{\lambda}{2(1+\sqrt{2})} \left[(b-a)^2/4 + 2K(\mathcal{T}) \log(e|\mathcal{T}|) \|W\|_\infty^2 / \beta \right]. \quad (4.34) \end{aligned}$$

This gives a bound for (4.33). It remains to derive an upper bound for (4.32). Here, we just have to follow the same line of reasoning as in the proof of the Lemma 36 on page 90,

which consists in applying a PAC-Bayesian inequality with influence function ψ combined with Jensen's inequality. Using the previous lemma, we are led to

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} \rho_{\theta' \mid \theta} \left[\mathbb{P}_W \left(\tilde{f}(\theta', W) \right) - \bar{\mathbb{P}}_W \left(\lambda^{-1} \psi [\lambda \tilde{f}(\theta', W)] \right) \right] \right\} \\ \leq \lambda \left[(b-a)^2/4 + 2K(\mathcal{T}) \log(e|\mathcal{T}|) \|W\|_\infty^2 / \beta \right] + \frac{\beta \|\Theta\|^2}{2n\lambda}. \end{aligned}$$

Then, combining this inequality with (4.34), we get

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta' \mid \theta} f(\theta', W) \right\} \\ \leq \frac{(\sqrt{2}+1)\lambda}{2} \left[(b-a)^2/4 + 2K(\mathcal{T}) \log(e|\mathcal{T}|) \|W\|_\infty^2 / \beta \right] + \frac{\beta \|\Theta\|^2}{2n\lambda}. \end{aligned}$$

At this point, we choose

$$\lambda = \sqrt{\frac{4\beta \|\Theta\|^2}{(\sqrt{2}+1) \left[(b-a)^2 + 8K(\mathcal{T}) \log(e|\mathcal{T}|) \|W\|_\infty^2 / \beta \right] n}},$$

which leads to

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) \rho_{\theta' \mid \theta} f(\theta', W) \right\} \\ \leq \tilde{F}(\beta) \stackrel{\text{def}}{=} \sqrt{\frac{(\sqrt{2}+1) \left(\beta(b-a)^2 + 8K(\mathcal{T}) \log(e|\mathcal{T}|) \|W\|_\infty^2 \right) \|\Theta\|^2}{4n}}. \end{aligned}$$

Therefore,

$$\mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W) \right\} \leq 2\sqrt{2K(\mathcal{T}) \log(|\mathcal{T}|) / \beta} \|W\|_\infty + \tilde{F}(2^{-H}\beta) + HF,$$

where F is defined in (4.28). Take

$$H = \left\lfloor \log \left(\frac{n}{K(\mathcal{T})\mathcal{S}} \right) \log(2)^{-1} \right\rfloor$$

and $\beta = 2n\|\Theta\|^{-2}$, so that

$$2^{-H}\beta \leq 4K(\mathcal{T})\mathcal{S}\|\Theta\|^{-2}.$$

We obtain

$$\begin{aligned} \mathbb{P}_{W_1, \dots, W_n} \left\{ \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W) \right\} \\ \leq \left(\frac{\log(n \mathcal{S}^{-1} K(\mathcal{T})^{-1})}{\log(2)} \sqrt{\frac{8K(\mathcal{T}) \log(|\mathcal{T}|)}{n}} + 2\sqrt{\frac{K(\mathcal{T}) \log(|\mathcal{T}|)}{n}} \right) \|\Theta\| \|W\|_\infty \\ + \sqrt{\frac{(\sqrt{2}+1) \left(\mathcal{S}(b-a)^2 + 2\log(e|\mathcal{T}|) \|W\|_\infty^2 \|\Theta\|^2 \right) K(\mathcal{T})}{n}}. \end{aligned}$$

Accordingly, from the bounded difference inequality, we get with probability at least $1 - \delta$

$$\begin{aligned} & \sup_{\theta \in \Theta} (\mathbb{P}_W - \bar{\mathbb{P}}_W) f(\theta, W) \\ & \leq \left(\frac{\log(n \mathbf{S}^{-1} K(\mathbf{T})^{-1})}{\log(2)} \sqrt{\frac{8K(\mathbf{T}) \log(|\mathbf{T}|)}{n}} + 2 \sqrt{\frac{K(\mathbf{T}) \log(|\mathbf{T}|)}{n}} \right) \|\Theta\| \|W\|_\infty \\ & \quad + \sqrt{\frac{(\sqrt{2} + 1) (\mathbf{S}(b-a)^2 + 2 \log(e|\mathbf{T}|) \|W\|_\infty^2 \|\Theta\|^2) K(\mathbf{T})}{n}} + \sqrt{\frac{\log(\delta^{-1})}{2n}} (b-a). \end{aligned}$$

In that way, we obtain the first part of the lemma. The other part follows in the same manner as in the proof of Lemma 36 on page 90. \square

At this point, we are in a position to apply previous Lemma 42 on page 101 and get faster speeds with respect to n than in Propositions 33 on page 86, 34 on page 88 and 35 on page 89 for the different risks $(\mathcal{C}_i)_{i \in \{1,3,4\}}$ defined in section 4.4 on page 78.

PROPOSITION 45 *Consider the same setting as in Proposition 33 on page 86. Consider the model*

$$\mathbf{M}(\mathbf{S}) = \left\{ \mu \in [-B, B]^{d \times k} : \sum_{j=1}^k \mathbb{P}_S(\text{supp}(\mu_j)) \leq \mathbf{S}, |\mathbf{T}_{\mu,K}| \geq 2 \right\}.$$

Consider any $k \geq 2$, any $K \geq 1$, any $\mathbf{S} \in [1, k]$, any $n \geq 2\mathbf{S}K$, and any $\delta \in]0, 1[$. With probability at least $1 - \delta$, for any $\mu \in \mathbf{M}(\mathbf{S})$,

$$\begin{aligned} \mathcal{C}_1(\mu) - \mathbb{P}_X[\mathbb{P}_S(X_S^2)] & \leq \bar{\mathcal{C}}_1(\mu) - \bar{\mathbb{P}}_X[\mathbb{P}_S(X_S^2)] \\ & \quad + \sqrt{10} B^2 \left(\frac{\log(n \mathbf{S}^{-1} K^{-1})}{\log(2)} \sqrt{\frac{8\mathbf{S}K \log(|\mathbf{T}_{\mu,K}|)}{n}} + 2 \sqrt{\frac{\mathbf{S}K \log(|\mathbf{T}_{\mu,K}|)}{n}} \right) \\ & \quad + B^2 \left(\sqrt{\frac{4(\sqrt{2} + 1)(9 + 5 \log(|\mathbf{T}_{\mu,K}|)) \mathbf{S}K}{n}} + 2 \sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} \right). \end{aligned}$$

For any non random set of fragments $\mu^ \in \mathbf{M}(\mathbf{S})$, with probability at least $1 - \delta$, for any $\mu \in \mathbf{M}(\mathbf{S})$,*

$$\begin{aligned} \mathcal{C}_1(\mu) - \mathcal{C}_1(\mu^*) - \bar{\mathcal{C}}_1(\mu) + \bar{\mathcal{C}}_1(\mu^*) & \leq \left(\frac{\log(n \mathbf{S}^{-1} K^{-1})}{\log(2)} \sqrt{\frac{8\mathbf{S}K \log(|\mathbf{T}_{\mu,K}|)}{n}} + 2 \sqrt{\frac{\mathbf{S}K \log(|\mathbf{T}_{\mu,K}|)}{n}} \right) \sqrt{10} B^2 \\ & \quad + \sqrt{\frac{4(\sqrt{2} + 1)(4 + 5 \log(e|\mathbf{T}_{\mu,K}|)) \mathbf{S}K}{n}} B^2 + 4 \sqrt{\frac{k(k-1) \log(2) + 2 \log(\delta^{-1})}{n}} B^2. \end{aligned}$$

Consequently, if

$$\hat{\mu} \in \arg \min_{\mu \in \mathbf{M}(\mathbf{S})} \bar{\mathcal{C}}_1(\mu) + \left(\frac{\log(n \mathbf{S}^{-1} K^{-1})}{\log(2)} \sqrt{\frac{8\mathbf{S}K \log(|\mathbf{T}_{\mu,K}|)}{n}} + 2 \sqrt{\frac{\mathbf{S}K \log(|\mathbf{T}_{\mu,K}|)}{n}} \right) \sqrt{10} B^2$$

$$+ \sqrt{\frac{4(\sqrt{2}+1)\left(4+5\log(e|\mathcal{T}_{\mu,K}|)\right)\mathcal{S}K}{n}B^2},$$

with probability at least $1 - \delta$,

$$\begin{aligned} \mathcal{C}_1(\hat{\mu}) \leq \inf_{\mu \in \mathcal{M}(\mathcal{S})} \mathcal{C}_1(\mu) &+ \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \sqrt{10}B^2 \\ &+ \sqrt{\frac{4(\sqrt{2}+1)\left(4+5\log(e|\mathcal{T}_{\mu,K}|)\right)\mathcal{S}K}{n}B^2} + 4\sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{n}B^2}. \end{aligned}$$

PROOF. In section 4.4 on page 78, the risk \mathcal{C}_1 has already been written in a suitable form. Therefore the proof follows from Lemma 42 on page 101 and the bounds $\|\Theta\| \leq B\sqrt{2\mathcal{S}}$, $\|W\|_\infty \leq B\sqrt{5}$ and $K(\mathcal{T}) \leq K$, along with $b - a \leq 4B^2$. \square

Let us state now the faster bounds related to the risk \mathcal{C}_3 .

PROPOSITION 46 Consider the model

$$\mathcal{M}(\mathcal{S}) = \left\{ \mu \in \mathbb{R}^{d \times k} : \sum_{j=1}^k \mathbb{P}_S(\text{supp}(\mu_j)) \leq \mathcal{S}, |\mathcal{T}_{\mu,K}| \geq 2 \right\},$$

where $\mathcal{S} \in [1, k]$. Assume that $n \geq 2\mathcal{S}K$. For any $\delta \in]0, 1[$, with probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathcal{S})$,

$$\begin{aligned} \mathcal{C}_3(\mu) \leq \bar{\mathcal{C}}_3(\mu) &+ 2\sigma^2 \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ &+ 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)\left(1+2\log(e|\mathcal{T}_{\mu,K}|)\right)\mathcal{S}K}{n}} + \sigma^2 \sqrt{\frac{2kd\log(2) + 2\log(\delta^{-1})}{n}}. \end{aligned}$$

Consider a non random set of fragments $\mu^* \in \mathcal{M}(\mathcal{S})$. With probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathcal{S})$,

$$\begin{aligned} \mathcal{C}_3(\mu) - \mathcal{C}_3(\mu^*) \leq \bar{\mathcal{C}}_3(\mu) - \bar{\mathcal{C}}_3(\mu^*) &+ 2\sigma^2 \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ &+ 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)\left(1+2\log(e|\mathcal{T}_{\mu,K}|)\right)\mathcal{S}K}{n}} + 2\sigma^2 \sqrt{\frac{2kd\log(2) + 2\log(\delta^{-1})}{n}}. \end{aligned}$$

Consequently, if

$$\hat{\mu} \in \arg \min_{\mu \in \mathcal{M}(\mathcal{S})} \bar{\mathcal{C}}_3(\mu) + 2\sigma^2 \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right)$$

$$+ 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)(1+2\log(e|\mathcal{T}_{\mu,K}|))\mathcal{S}K}{n}},$$

with probability at least $1 - \delta$,

$$\begin{aligned} \mathcal{C}_3(\widehat{\mu}) &\leq \inf_{\mu \in \mathcal{M}(\mathcal{S})} \mathcal{C}_3(\mu) \\ &\quad + 2\sigma^2 \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ &\quad + 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)(1+2\log(e|\mathcal{T}_{\mu,K}|))\mathcal{S}K}{n}} + 2\sigma^2 \sqrt{\frac{2kd \log(2) + 2\log(\delta^{-1})}{n}}. \end{aligned}$$

PROOF. Considering the formulation of the risk \mathcal{C}_3 in terms of θ and W given in section 4.4 on page 78 and using the fact that $\|W\|_\infty = 1$, $b - a = 1$ and $\|\Theta\| \leq \mathcal{S}^{1/2}$, we conclude from Lemma 42 on page 101. \square

As discussed in section 4.4, the generalization bounds obtained for the risk \mathcal{C}_3 depend on the (possibly high) dimension d of the signal. This is a motivation to consider rather the risk \mathcal{C}_4 , for which we know how to get dimension free bounds.

PROPOSITION 47 Define the model

$$\mathcal{M}(\mathcal{S}) = \left\{ \mu \in \mathbb{R}^{d \times k} : \sum_{j=1}^k \mathbb{P}_S(\text{supp}(\mu_j)) \leq \mathcal{S}, |\mathcal{T}_{\mu,K}| \geq 2 \right\},$$

where $\mathcal{S} \in [1, k]$ and assume that $n \geq 2\mathcal{S}K$. For any $\delta \in]0, 1[$, with probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathcal{S})$,

$$\begin{aligned} \mathcal{C}_4(\mu) &\leq \bar{\mathcal{C}}_4(\mu) + 2\sigma^2 \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ &\quad + 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)(1+2\log(e|\mathcal{T}_{\mu,K}|))\mathcal{S}K}{n}} + \sigma^2 \sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{n}}. \end{aligned}$$

Moreover, if $\mu^* \in \mathcal{M}(\mathcal{S})$ is a non random set of fragments, with probability at least $1 - \delta$, for any $\mu \in \mathcal{M}(\mathcal{S})$,

$$\begin{aligned} \mathcal{C}_4(\mu) - \mathcal{C}_4(\mu^*) - \bar{\mathcal{C}}_4(\mu) + \bar{\mathcal{C}}_4(\mu^*) &\leq 2\sigma^2 \left(\frac{\log(n\mathcal{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathcal{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ &\quad + 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)(1+2\log(e|\mathcal{T}_{\mu,K}|))\mathcal{S}K}{n}} + 2\sigma^2 \sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{n}}. \end{aligned}$$

Consequently, if

$$\begin{aligned}\widehat{\mu} \in \arg \min_{\mu \in \mathcal{M}(\mathbf{S})} \overline{\mathbf{C}}_4(\mu) + 2\sigma^2 \left(\frac{\log(n\mathbf{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathbf{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathbf{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ + 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)(1+2\log(e|\mathcal{T}_{\mu,K}|))\mathbf{S}K}{n}},\end{aligned}$$

with probability at least $1 - \delta$

$$\begin{aligned}\mathbf{C}_4(\widehat{\mu}) \leq \inf_{\mu \in \mathcal{M}(\mathbf{S})} \mathbf{C}_4(\mu) + 2\sigma^2 \left(\frac{\log(n\mathbf{S}^{-1}K^{-1})}{\log(2)} \sqrt{\frac{8\mathbf{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} + 2\sqrt{\frac{\mathbf{S}K \log(|\mathcal{T}_{\mu,K}|)}{n}} \right) \\ + 2\sigma^2 \sqrt{\frac{(\sqrt{2}+1)(1+2\log(e|\mathcal{T}_{\mu,K}|))\mathbf{S}K}{n}} + 2\sigma^2 \sqrt{\frac{k(k-1)\log(2) + 2\log(\delta^{-1})}{n}}.\end{aligned}$$

PROOF. We can apply Lemma 42 on page 101 to the risk \mathbf{C}_4 expressed in terms of θ and W . Let us recall that

$$\mathbf{C}_4(g, \mu) = 2\sigma^2 \left[1 + \mathbb{P}_X \left(\min_{A \in \mathcal{T}_{g,K}} \underbrace{\sum_{j \in A} \langle \theta_j, W_j \rangle}_{\in [-1,0]} \right) \right],$$

where

$$\begin{aligned}\theta_j(s) &= \mathbb{1}(s \in B_j) \Psi(\mu_{j,s}), \\ X_j(s) &= -\Psi(X_s), \quad 1 \leq s \leq d, \quad 1 \leq j \leq k.\end{aligned}$$

We conclude the proof remarking that $\|\theta_j\|^2 = \mathbb{P}_S(B_j)$, $\|W\|_\infty = 1$, and $\|\Theta\| \leq \mathbf{S}^{1/2}$. \square

CHAPTER 5


Experiment on digital images

In this section, we describe some very preliminary experiment showing what to expect and hope for. We wrote a program that computes the syntax tree from a training set $(X_1, \dots, X_n) \in \mathbb{R}^{d \times n}$. Each X_i is a 300×300 greyscale image extracted at random from two larger greyscale images (we only kept the green channel from the original jpeg images) and $n = 200$. In order to play with the different parameters and visualize the experimental results in an interactive way, we decided to create a graphical user interface with the help of the `rWidgets2` package of the R language. We created two control panes containing several buttons and sliders to monitor the experiment. A first pane allows to load images and extract samples from them as square windows of the desired size. The positions of the windows can be constrained to a grid and are otherwise random. The grey levels are transformed by

$$f(x) = \log(10^\varepsilon + x), \quad x \in [0, 1],$$

the default value for ε being -2 as seen on the screen shot. This transform is justified by the fact that we will use an additive error model whereas we are more sensitive to light intensity ratios than to light intensity differences. Hence the logarithmic transform.


This is the control pane allowing to pick-up the training set drawn at random from larger scenes.




We will show results on a sample drawn from two scenes featuring the same kitten.



The images are drawn from the yellow area of each scene. This is the beginning of the training sample



The fragmentation and syntax analysis are launched from a second control pane.



The parameter $\beta = \text{Beta}$ is used to compute the fragmentation criterion

$$2 \times 10^\beta |B_j| - |A_j|.$$

The parameter $\tau = \text{Threshold value}$ is used to compute the threshold

$$\alpha = \tau \bar{\mathbb{P}}_S \left[\text{Var}(\bar{\mathbb{P}}_{X_{I,S} | S}) \right]$$

appearing in equation (2.9) on page 26. Pressing the **Compute initial patches** button executes the sample fragmentation into a maximum number of patches prescribed in the text field **Number of patches**. Pressing the **Compute next level patches** button computes the next two levels of the syntax tree. This means that it computes pair labels and deduces from the compression of the pair label list a syntax classification f to be applied to pair labels. Pressing the button many times, we can grow the syntax tree ad libitum, until no new relabelling is observed. See the figure representing the syntax tree on page 32. The text fields **Number of merged labels** and **Number of syntax labels** specify the maximum number of labels to be created at each stage. Visualization is performed with the help of the sliders occupying the second half of the control pane. The first slider, named **Syntax level**, describes which level of the syntax tree is displayed. The second slider, titled **Show patches in image number**, displays all the patches in a given image (at the syntax level indicated by the previous slider). The next slider goes through all the patches present in a given image at a

given level. In other words, it displays one level of the syntax tree of a given image. The last slider, titled **Change image**, gathers all the images containing a given patch label at a given syntax level. The buttons **Show all** and **Show patch** toggle between displaying a particular patch on a green background and displaying all the patches present in one image using a rainbow of colors.

The experiment can be viewed as a crude vision model where images of two scenes are acquired by some sensor or retina through random eye saccades. In our experiment, the sample is shuffled by a random permutation and the locations in each scene of the acquired images are also random.

This is somehow a stress test for a vision model since there is no structure in the image acquisition process, by opposition to what we would get for instance if the image sample was extracted from a video stream.


In this situation, two questions come to mind.

1. Will our unsupervised classification algorithm be able to put the same label on images that come from the same scene and are overlapping, so that they have some content in common, but translated. Note that our syntax analysis does not include any explicit translation model. The algorithm only compares pixel values being at the same pixel location in different sample images. The algorithm indeed makes no use of pixel location, that is of the sensor geometry. It is invariant with respect to any arbitrary permutation of the pixels, as long as the same permutation is applied at the same time to all sample images. Thus labelling translated patterns is a challenge. What we hope for is that syntax analysis can solve the problem of invariant pattern classification in a novel way. Syntax analysis describes the world using a mix of logics and compression theory, building up relabelling schemes described by rewriting rules and chosen for their ability to compress information. Chaining rewriting rules can be seen as some sort of crude logical reasoning. So the question is whether this crude logical reasoning can produce a flexible solution to the problem of invariant pattern recognition ? The more classical approach is to model explicitly pattern transformations using geometry or differential geometry and to pose the question of invariant classification with respect to a family of transformations in mathematical terms. Unfortunately, this route leads to daunting mathematical challenges, even in the somehow simplest case of translation invariance.
2. Will syntax analysis be able to put the same label on images containing related patterns coming from different scenes ? For example, in our sample, the kitten's head is present in the two scenes, but slightly rotated and on different backgrounds. Will it be possible to identify those two views of the same head, giving them the same label ?


We will see that the answer to both questions is positive. Although more experiments would be needed, to see how stable things are and how the algorithm scales with larger data bases,

this is encouraging.


Let us show first the result of the fragmentation algorithm itself. Visually, when $\alpha = 1/10$, the coding distortion does not ruin the image content. We give two examples, on the left we see the patch reconstruction and on the right the original image




We now show two examples of fragment supports (when we use 1000 fragments for the whole sample). Colors do not match, we use a different rainbow in the two examples, each rainbow indexing only the fragments present in the corresponding image.




We will now choose an image, and show first its fragments



and then its syntax labels at the highest level (we computed a syntax tree with 2×5 levels, stopping when no new syntax levels are added). The levels are counted as in the diagram on page 32. Two levels are created each time we press the **Compute next level patches** button.



We show all the other images sharing with it a syntax label of the highest level. It turns out that each label is either specific to the reference image, or shared by two images, except for one label that is shared by three images. We start with a label shared by an image drawn from the other scene, since this is maybe the most exciting part of the result



On the left, we can see the patch corresponding to the common label. In this case it has the


same support on both images, but we will see that it is not always the case (as expected). Let us see now another patch




This time the patch has the same syntax label, but not the same support in both images. We can see that the support is somewhat shifted in the direction of the translation. This gives the impression that the syntax analysis performed on top of the fragmentation algorithm brings a positive contribution to the detection of translations. Let us see the other patches. Here again, the patch support changes




whereas in the next case, it is fixed




so that we do not repeat the original image. It is the same with the next two patches.



Finally, we have a syntax label indexing three images, whose support takes two different values, as shown in the next three pictures.





In conclusion, this small experiment let us hope for much. We have a pattern matching principle that produces visually interesting results. Moreover, it can be applied directly on high dimensional data, here we work with a sample of size 200 in \mathbb{R}^{90000} without being plagued by the curse of dimension. This was predicted at least for the fragmentation part by our dimension free generalization bounds, and it seems to be also the case in practice. That does not mean that, if we were to build a more serious vision algorithm, we could not combine the approach with some preprocessing. Indeed, our algorithm can take as input any vector of measurements, so we could try to extract first a vector of real valued features, like for instance wavelet coefficients. A more thoughtful vision model would also presumably include some sort of multi-scale processing, as well as the use of a retina with a space varying pixel density. Anyhow we preferred to apply a single general purpose treatment to raw data, to show that it can manage to deliver results by itself.

CHAPTER 6

Conclusion

In this thesis, we proposed new algorithms to perform unsupervised clustering and suggested new ideas for signal modeling, with a special focus on digital images. To this aim, we first introduced the information k -means algorithm and developed its mathematical analysis. We showed that it is a generalization of the classical Euclidean k -means criterion and demonstrated its benefits for the clustering of bag of words. Then, based on the interpretation of the information k -means setting as a density estimation framework, we introduced alternative bounded loss functions. Those criteria do not require any integrability assumptions on the sample, and can be regarded as robust variants of the usual k -means loss function.

We put the information k -means algorithm into a broader context that we called information fragmentation. On top of proving generalisation bounds, we justified this algorithm from a data compression perspective, as it computes shorter indices for large and frequent data blocks, similarly to the Lempel Ziv algorithm. We described fragmentation for signals belonging to \mathbb{R}^d . Although it covers the case of digital images, it is in fact much more general, since we do not use pixel geometry through any kind of neighboring relations. More precisely, the same fragments will be computed if we apply any given permutation to the pixel indices of all the images of a given training sample. Thus, the fragmentation algorithm can be applied to any kind of signal, color images, 3D images, video streams, speech and so on.

The fragmentation algorithm is a lossy compression scheme that maps any signal consisting in a vector of real valued measurements to a discrete representation consisting in a set of fragment labels. This mapping is chosen to optimize the compression of a training sample. Compression is then pushed further using our next proposal: a syntax analysis algorithm.

It is made of two stages: grouping and context analysis. The grouping stage performs a lossless compression by merging pairs of labels that appear frequently. This relabeling scheme produces new non terminal symbols and binary context free rewriting rules that we can gather into a context free grammar. The signal representation is changed to a new set of labels, containing non terminal *super fragment* labels that rewrite each into a unique set of fragment labels. The context analysis stage consists in compressing the representation of the

grammar through factoring and grouping of the rewriting rules. It produces a grammar of the grammar made of new rewriting rules for new syntax labels. These syntax labels perform some kind of context analysis drawn by a compression criterion. They induce a classification of the *super fragment* labels into syntax categories. We get a new signal representation using syntax categories. Repeating this scheme as long as we can increase the training sample compression rate, we build a syntax tree producing a hierarchical unsupervised classification of the content of each signal of the training set. Note that the whole scheme is guided by a single criterion, the compression rate and uses essentially two ingredients, grouping and factorization.

This is different from contextual modeling based on conditional probability measures, whose estimation is a difficult statistical inference issue. The choice of a compression scheme rather than a statistical model may be a way to avoid the curse of dimension as suggested by Lemma 4 on page 33. This lemma relates our compression approach to a statistical estimator whose generalization properties depend on the compression rate of the compression scheme.

The rest of our dissertation is devoted to the mathematical justification of the fragmentation algorithm through generalization bounds. These generalization bounds characterize the stability of the fragmentation, assessing that fragments computed from one training set would work almost as well in expectation, and consequently would also work almost as well to represent another independent training set.

Combining PAC-Bayesian lemmas with the kernel trick, we established dimension-free non asymptotic bounds on the excess risk of both k -means, information k -means and information fragmentation. These bounds show that the fragmentation algorithm does not overfit the data as long as

$$\frac{K^2 \mathbf{S} \log(k/K)}{n}$$

goes to zero, where k is the number of fragments K the maximum number of fragments used to represent a single signal, and \mathbf{S} is the total number of pixels of the fragments divided by the number of pixels in one image.

We used the same line of proof to obtain generalization bounds of order

$$\mathcal{O}\left(\frac{k \log(k)}{n}\right)^{1/4}$$

for the classical k -means risk in a separable Hilbert space. In order to improve on the $1/4$ exponent, we blended the chaining method and the PAC-Bayesian technology to get bounds of order

$$\mathcal{O}\left(\log\left(\frac{n}{k}\right)\left(\frac{k \log(k)}{n}\right)^{1/2}\right)$$

for our various flavours of the k -means criterion, including the classical one and improving in this case on the bound of order

$$\mathcal{O}\left(\frac{k}{\sqrt{n}}\right)$$

proved in [BDL08].

Finally, we produced experimental results to show what the fragmentation algorithm combined with syntax analysis is capable of, when applied to digital images. The experiment, although preliminary, was very encouraging. First, we worked with 300×300 images, randomly drawn from two larger scenes, without suffering from the curse of dimension. Second, we worked with a sample of size 200, showing that interesting things can be learnt even from small training sets. Third, the algorithm was able to recognize (that is give the same label to) patterns present in the two scenes (a kitten's head) as well as translations of the observation window. This fosters the hope to bring a generic solution to the problem of invariant pattern recognition, where invariance with respect to transformations is understood in a loose sense, without the need to build specific methods dealing with a mathematically well defined subset of pattern transformations.

All this of course is very preliminary. We wanted to transpose syntax analysis from texts to signals, making a realistic proposal for unsupervised signal classification that could be demonstrated on real data. To complete this, we had somehow to stop as soon as we got some success at each step of our research, in order to keep some time to explore the next step. So we are pretty sure everything can be improved. For instance, the optimization strategy both in the fragmentation algorithm and in the syntax analysis is a one pass scheme: we build smaller and smaller fragments, never coming back on past choices, and then build larger and larger super fragments in the syntax phase, whereas it could be worth going up and down more than one time. Also, we presented a batch algorithm, letting open the question of its sequential pendent. Experimenting how the method scales on bigger data sets would also be necessary. The relationship between compression and estimation that Lemma 4 on page 33 hints at would deserve a more precise study. The syntax analysis scheme we propose is a crude first attempt, a more systematic investigation of the use of context free grammars to build compression schemes should be possible.

The use of chaining in PAC-Bayesian bounds is another technical subject that calls for further studies.


Further research may also include testing the method on other types of data, such as speech, or combinations of different types of data, like video streams with sound.

Our agenda also includes coming back to natural language processing to test the pendent of our syntax analysis algorithm for natural language processing. This would pave the way to combined data analysis, such as images with textual comments.

APPENDIX A

Code highlights

The syntax analysis is coded in R and C++, using the interface provided by the Rcpp package, see [EF11] and [Edd13]. We used the `Rcpp.package.skeleton()` function, to create our own R package. The advantage of this approach is that the communication between R and C++ is made easier and the final code represents a self-contained R package, thus easy to distribute, see [EF13] for more details concerning the creation of R packages with Rcpp. Moreover, the C++ code is parallelized using OpenMP parallelization directives and makes good use of an eight core processor as shown by the result of the linux htop command.



We give the code of the main C++ functions implementing the fragmentation algorithm, to show how things can be done in practice, when the goal is only to compute the syntax tree. The implementation of the rest of the syntax tree, that is the computation of the pair labels and syntax labels, is quite similar. The main loop `computeLoop()` (line 5) runs the `split()` function (line 25), that is defined on line 32. In order to take less memory, variables and objects are passed to C++ via pointers, and all functions are of type `void`. Notice before the definition of the `computeLoop` function, the typical Rcpp instruction `// [[`

`Rcpp :: export]]` that is required to make any compiled C++ function accessible into the R environment. Besides, we adopt an object oriented programming approach creating 2 classes (similar to the `struct` data type in C) named `Patches` and `mergeParam`. For the sake of clarity, we omit the description of the corresponding header file `initialization.h` containing the declaration of the classes. The class `Patches` is associated with the fragmentation step whereas the `mergeParam` is related to the merge and syntax analysis step. For the purpose of brevity, we will only talk about the class `Patches`. This class is composed of several attributes but more importantly contains the method `split()` that is called in the critical loop. The first method is called after the instantiation of a `Patches` object named `parameters` in the `computeLoop` function below. This method is used to initialize the quantity α , mentioned in the introduction of the experiment. Then, comes the call of the `split()` method which tries to find the pair $\{j_{k,1}, j_{k,2}\}$ (noted `max_i` and `max_j` in the code) maximizing the fragmentation criterion, as it is described in section 2.3 on page 25. This is done essentially by computing the fragmentation criterion for each iteration but in a parallel manner due to the compiler directive `#pragma omp parallel for` before the C++ `for` loop. As we already mentioned, this will distribute the computation of the criterion over the different processors. We perform first the computation of the index $j_{k,1}$ (`max_i`) in a first loop and then $j_{k,2}$ (`max_j`) in a second loop. As we discussed in section 2.3 on page 25, this approach avoids the computation of the criterion over all the possible pairs J_k . It is important to notice that we make use of the compiler directive `#pragma omp critical` twice. This compiler directive allows us to execute sequentially a block of code over the different processors. This means that the block of code can only be executed by one processor at a time. In our particular case, our critical block of code will be designed to retrieve the maximum fragmentation criterion (noted `max_crit` in the code), as well as the corresponding index j_k each processor had to compute in his own set of iterations. This provides us a way to reduce all the computations, performed separately over the processors, into one single result. Let us also point out that `max_crit` as well as `max_i` and `max_j` are global variables whereas `my_max_crit`, `my_max_i` and `my_max_j` are local variables private to each processor. Then, the remaining of the code is a matter of updating the storage of the criterion and the pair $\{j_{k,1}, j_{k,2}\}$, along with the matrices A and B representing the sequences of sets $A_{k,j}$ and $B_{k,j}$, $j \in \llbracket 1, k \rrbracket$. For an in-depth look at the code, the reader may look at the GitHub repository <https://github.com/GautierAppert/PatchProcess>.

```

2 #include "initialization.h"
3
4 // [[Rcpp::export]]
5
6 void computeLoop (
7     NumericVector &m,
8     NumericVector &v,
9     NumericVector &w,
10    NumericVector &C,
11
12    NumericVector &alpha,
13    NumericVector &beta,
14    NumericVector &gamma,
15    NumericVector &delta,
16    NumericVector &epsilon,
17    NumericVector &phi,
18    NumericVector &psi,
19    NumericVector &theta,
20    NumericVector &rho,
21    NumericVector &sigma,
22    NumericVector &tau,
23    NumericVector &lambda,
24    NumericVector &mu,
25    NumericVector &nu,
26    NumericVector &xi,
27    NumericVector &eta,
28    NumericVector &zeta,
29    NumericVector &xi2,
30    NumericVector &eta2,
31    NumericVector &zeta2,
32    NumericVector &phi2,
33    NumericVector &psi2,
34    NumericVector &theta2,
35    NumericVector &rho2,
36    NumericVector &sigma2,
37    NumericVector &tau2,
38    NumericVector &lambda2,
39    NumericVector &mu2,
40    NumericVector &nu2,
41    NumericVector &xi3,
42    NumericVector &eta3,
43    NumericVector &zeta3,
44    NumericVector &phi3,
45    NumericVector &psi3,
46    NumericVector &theta3,
47    NumericVector &rho3,
48    NumericVector &sigma3,
49    NumericVector &tau3,
50    NumericVector &lambda3,
51    NumericVector &mu3,
52    NumericVector &nu3,
53    NumericVector &xi4,
54    NumericVector &eta4,
55    NumericVector &zeta4,
56    NumericVector &phi4,
57    NumericVector &psi4,
58    NumericVector &theta4,
59    NumericVector &rho4,
60    NumericVector &sigma4,
61    NumericVector &tau4,
62    NumericVector &lambda4,
63    NumericVector &mu4,
64    NumericVector &nu4,
65    NumericVector &xi5,
66    NumericVector &eta5,
67    NumericVector &zeta5,
68    NumericVector &phi5,
69    NumericVector &psi5,
70    NumericVector &theta5,
71    NumericVector &rho5,
72    NumericVector &sigma5,
73    NumericVector &tau5,
74    NumericVector &lambda5,
75    NumericVector &mu5,
76    NumericVector &nu5,
77    NumericVector &xi6,
78    NumericVector &eta6,
79    NumericVector &zeta6,
80    NumericVector &phi6,
81    NumericVector &psi6,
82    NumericVector &theta6,
83    NumericVector &rho6,
84    NumericVector &sigma6,
85    NumericVector &tau6,
86    NumericVector &lambda6,
87    NumericVector &mu6,
88    NumericVector &nu6,
89    NumericVector &xi7,
90    NumericVector &eta7,
91    NumericVector &zeta7,
92    NumericVector &phi7,
93    NumericVector &psi7,
94    NumericVector &theta7,
95    NumericVector &rho7,
96    NumericVector &sigma7,
97    NumericVector &tau7,
98    NumericVector &lambda7,
99    NumericVector &mu7,
100   NumericVector &nu7,
101   NumericVector &xi8,
102   NumericVector &eta8,
103   NumericVector &zeta8,
104   NumericVector &phi8,
105   NumericVector &psi8,
106   NumericVector &theta8,
107   NumericVector &rho8,
108   NumericVector &sigma8,
109   NumericVector &tau8,
110   NumericVector &lambda8,
111   NumericVector &mu8,
112   NumericVector &nu8,
113   NumericVector &xi9,
114   NumericVector &eta9,
115   NumericVector &zeta9,
116   NumericVector &phi9,
117   NumericVector &psi9,
118   NumericVector &theta9,
119   NumericVector &rho9,
120   NumericVector &sigma9,
121   NumericVector &tau9,
122   NumericVector &lambda9,
123   NumericVector &mu9,
124   NumericVector &nu9,
125   NumericVector &xi10,
126   NumericVector &eta10,
127   NumericVector &zeta10,
128   NumericVector &phi10,
129   NumericVector &psi10,
130   NumericVector &theta10,
131   NumericVector &rho10,
132   NumericVector &sigma10,
133   NumericVector &tau10,
134   NumericVector &lambda10,
135   NumericVector &mu10,
136   NumericVector &nu10,
137   NumericVector &xi11,
138   NumericVector &eta11,
139   NumericVector &zeta11,
140   NumericVector &phi11,
141   NumericVector &psi11,
142   NumericVector &theta11,
143   NumericVector &rho11,
144   NumericVector &sigma11,
145   NumericVector &tau11,
146   NumericVector &lambda11,
147   NumericVector &mu11,
148   NumericVector &nu11,
149   NumericVector &xi12,
150   NumericVector &eta12,
151   NumericVector &zeta12,
152   NumericVector &phi12,
153   NumericVector &psi12,
154   NumericVector &theta12,
155   NumericVector &rho12,
156   NumericVector &sigma12,
157   NumericVector &tau12,
158   NumericVector &lambda12,
159   NumericVector &mu12,
160   NumericVector &nu12,
161   NumericVector &xi13,
162   NumericVector &eta13,
163   NumericVector &zeta13,
164   NumericVector &phi13,
165   NumericVector &psi13,
166   NumericVector &theta13,
167   NumericVector &rho13,
168   NumericVector &sigma13,
169   NumericVector &tau13,
170   NumericVector &lambda13,
171   NumericVector &mu13,
172   NumericVector &nu13,
173   NumericVector &xi14,
174   NumericVector &eta14,
175   NumericVector &zeta14,
176   NumericVector &phi14,
177   NumericVector &psi14,
178   NumericVector &theta14,
179   NumericVector &rho14,
180   NumericVector &sigma14,
181   NumericVector &tau14,
182   NumericVector &lambda14,
183   NumericVector &mu14,
184   NumericVector &nu14,
185   NumericVector &xi15,
186   NumericVector &eta15,
187   NumericVector &zeta15,
188   NumericVector &phi15,
189   NumericVector &psi15,
190   NumericVector &theta15,
191   NumericVector &rho15,
192   NumericVector &sigma15,
193   NumericVector &tau15,
194   NumericVector &lambda15,
195   NumericVector &mu15,
196   NumericVector &nu15,
197   NumericVector &xi16,
198   NumericVector &eta16,
199   NumericVector &zeta16,
200   NumericVector &phi16,
201   NumericVector &psi16,
202   NumericVector &theta16,
203   NumericVector &rho16,
204   NumericVector &sigma16,
205   NumericVector &tau16,
206   NumericVector &lambda16,
207   NumericVector &mu16,
208   NumericVector &nu16,
209   NumericVector &xi17,
210   NumericVector &eta17,
211   NumericVector &zeta17,
212   NumericVector &phi17,
213   NumericVector &psi17,
214   NumericVector &theta17,
215   NumericVector &rho17,
216   NumericVector &sigma17,
217   NumericVector &tau17,
218   NumericVector &lambda17,
219   NumericVector &mu17,
220   NumericVector &nu17,
221   NumericVector &xi18,
222   NumericVector &eta18,
223   NumericVector &zeta18,
224   NumericVector &phi18,
225   NumericVector &psi18,
226   NumericVector &theta18,
227   NumericVector &rho18,
228   NumericVector &sigma18,
229   NumericVector &tau18,
230   NumericVector &lambda18,
231   NumericVector &mu18,
232   NumericVector &nu18,
233   NumericVector &xi19,
234   NumericVector &eta19,
235   NumericVector &zeta19,
236   NumericVector &phi19,
237   NumericVector &psi19,
238   NumericVector &theta19,
239   NumericVector &rho19,
240   NumericVector &sigma19,
241   NumericVector &tau19,
242   NumericVector &lambda19,
243   NumericVector &mu19,
244   NumericVector &nu19,
245   NumericVector &xi20,
246   NumericVector &eta20,
247   NumericVector &zeta20,
248   NumericVector &phi20,
249   NumericVector &psi20,
250   NumericVector &theta20,
251   NumericVector &rho20,
252   NumericVector &sigma20,
253   NumericVector &tau20,
254   NumericVector &lambda20,
255   NumericVector &mu20,
256   NumericVector &nu20,
257   NumericVector &xi21,
258   NumericVector &eta21,
259   NumericVector &zeta21,
260   NumericVector &phi21,
261   NumericVector &psi21,
262   NumericVector &theta21,
263   NumericVector &rho21,
264   NumericVector &sigma21,
265   NumericVector &tau21,
266   NumericVector &lambda21,
267   NumericVector &mu21,
268   NumericVector &nu21,
269   NumericVector &xi22,
270   NumericVector &eta22,
271   NumericVector &zeta22,
272   NumericVector &phi22,
273   NumericVector &psi22,
274   NumericVector &theta22,
275   NumericVector &rho22,
276   NumericVector &sigma22,
277   NumericVector &tau22,
278   NumericVector &lambda22,
279   NumericVector &mu22,
280   NumericVector &nu22,
281   NumericVector &xi23,
282   NumericVector &eta23,
283   NumericVector &zeta23,
284   NumericVector &phi23,
285   NumericVector &psi23,
286   NumericVector &theta23,
287   NumericVector &rho23,
288   NumericVector &sigma23,
289   NumericVector &tau23,
290   NumericVector &lambda23,
291   NumericVector &mu23,
292   NumericVector &nu23,
293   NumericVector &xi24,
294   NumericVector &eta24,
295   NumericVector &zeta24,
296   NumericVector &phi24,
297   NumericVector &psi24,
298   NumericVector &theta24,
299   NumericVector &rho24,
300   NumericVector &sigma24,
301   NumericVector &tau24,
302   NumericVector &lambda24,
303   NumericVector &mu24,
304   NumericVector &nu24,
305   NumericVector &xi25,
306   NumericVector &eta25,
307   NumericVector &zeta25,
308   NumericVector &phi25,
309   NumericVector &psi25,
310   NumericVector &theta25,
311   NumericVector &rho25,
312   NumericVector &sigma25,
313   NumericVector &tau25,
314   NumericVector &lambda25,
315   NumericVector &mu25,
316   NumericVector &nu25,
317   NumericVector &xi26,
318   NumericVector &eta26,
319   NumericVector &zeta26,
320   NumericVector &phi26,
321   NumericVector &psi26,
322   NumericVector &theta26,
323   NumericVector &rho26,
324   NumericVector &sigma26,
325   NumericVector &tau26,
326   NumericVector &lambda26,
327   NumericVector &mu26,
328   NumericVector &nu26,
329   NumericVector &xi27,
330   NumericVector &eta27,
331   NumericVector &zeta27,
332   NumericVector &phi27,
333   NumericVector &psi27,
334   NumericVector &theta27,
335   NumericVector &rho27,
336   NumericVector &sigma27,
337   NumericVector &tau27,
338   NumericVector &lambda27,
339   NumericVector &mu27,
340   NumericVector &nu27,
341   NumericVector &xi28,
342   NumericVector &eta28,
343   NumericVector &zeta28,
344   NumericVector &phi28,
345   NumericVector &psi28,
346   NumericVector &theta28,
347   NumericVector &rho28,
348   NumericVector &sigma28,
349   NumericVector &tau28,
350   NumericVector &lambda28,
351   NumericVector &mu28,
352   NumericVector &nu28,
353   NumericVector &xi29,
354   NumericVector &eta29,
355   NumericVector &zeta29,
356   NumericVector &phi29,
357   NumericVector &psi29,
358   NumericVector &theta29,
359   NumericVector &rho29,
360   NumericVector &sigma29,
361   NumericVector &tau29,
362   NumericVector &lambda29,
363   NumericVector &mu29,
364   NumericVector &nu29,
365   NumericVector &xi30,
366   NumericVector &eta30,
367   NumericVector &zeta30,
368   NumericVector &phi30,
369   NumericVector &psi30,
370   NumericVector &theta30,
371   NumericVector &rho30,
372   NumericVector &sigma30,
373   NumericVector &tau30,
374   NumericVector &lambda30,
375   NumericVector &mu30,
376   NumericVector &nu30,
377   NumericVector &xi31,
378   NumericVector &eta31,
379   NumericVector &zeta31,
380   NumericVector &phi31,
381   NumericVector &psi31,
382   NumericVector &theta31,
383   NumericVector &rho31,
384   NumericVector &sigma31,
385   NumericVector &tau31,
386   NumericVector &lambda31,
387   NumericVector &mu31,
388   NumericVector &nu31,
389   NumericVector &xi32,
390   NumericVector &eta32,
391   NumericVector &zeta32,
392   NumericVector &phi32,
393   NumericVector &psi32,
394   NumericVector &theta32,
395   NumericVector &rho32,
396   NumericVector &sigma32,
397   NumericVector &tau32,
398   NumericVector &lambda32,
399   NumericVector &mu32,
400   NumericVector &nu32,
401   NumericVector &xi33,
402   NumericVector &eta33,
403   NumericVector &zeta33,
404   NumericVector &phi33,
405   NumericVector &psi33,
406   NumericVector &theta33,
407   NumericVector &rho33,
408   NumericVector &sigma33,
409   NumericVector &tau33,
410   NumericVector &lambda33,
411   NumericVector &mu33,
412   NumericVector &nu33,
413   NumericVector &xi34,
414   NumericVector &eta34,
415   NumericVector &zeta34,
416   NumericVector &phi34,
417   NumericVector &psi34,
418   NumericVector &theta34,
419   NumericVector &rho34,
420   NumericVector &sigma34,
421   NumericVector &tau34,
422   NumericVector &lambda34,
423   NumericVector &mu34,
424   NumericVector &nu34,
425   NumericVector &xi35,
426   NumericVector &eta35,
427   NumericVector &zeta35,
428   NumericVector &phi35,
429   NumericVector &psi35,
430   NumericVector &theta35,
431   NumericVector &rho35,
432   NumericVector &sigma35,
433   NumericVector &tau35,
434   NumericVector &lambda35,
435   NumericVector &mu35,
436   NumericVector &nu35,
437   NumericVector &xi36,
438   NumericVector &eta36,
439   NumericVector &zeta36,
440   NumericVector &phi36,
441   NumericVector &psi36,
442   NumericVector &theta36,
443   NumericVector &rho36,
444   NumericVector &sigma36,
445   NumericVector &tau36,
446   NumericVector &lambda36,
447   NumericVector &mu36,
448   NumericVector &nu36,
449   NumericVector &xi37,
450   NumericVector &eta37,
451   NumericVector &zeta37,
452   NumericVector &phi37,
453   NumericVector &psi37,
454   NumericVector &theta37,
455   NumericVector &rho37,
456   NumericVector &sigma37,
457   NumericVector &tau37,
458   NumericVector &lambda37,
459   NumericVector &mu37,
460   NumericVector &nu37,
461   NumericVector &xi38,
462   NumericVector &eta38,
463   NumericVector &zeta38,
464   NumericVector &phi38,
465   NumericVector &psi38,
466   NumericVector &theta38,
467   NumericVector &rho38,
468   NumericVector &sigma38,
469   NumericVector &tau38,
470   NumericVector &lambda38,
471   NumericVector &mu38,
472   NumericVector &nu38,
473   NumericVector &xi39,
474   NumericVector &eta39,
475   NumericVector &zeta39,
476   NumericVector &phi39,
477   NumericVector &psi39,
478   NumericVector &theta39,
479   NumericVector &rho39,
480   NumericVector &sigma39,
481   NumericVector &tau39,
482   NumericVector &lambda39,
483   NumericVector &mu39,
484   NumericVector &nu39,
485   NumericVector &xi40,
486   NumericVector &eta40,
487   NumericVector &zeta40,
488   NumericVector &phi40,
489   NumericVector &psi40,
490   NumericVector &theta40,
491   NumericVector &rho40,
492   NumericVector &sigma40,
493   NumericVector &tau40,
494   NumericVector &lambda40,
495   NumericVector &mu40,
496   NumericVector &nu40,
497   NumericVector &xi41,
498   NumericVector &eta41,
499   NumericVector &zeta41,
500   NumericVector &phi41,
501   NumericVector &psi41,
502   NumericVector &theta41,
503   NumericVector &rho41,
504   NumericVector &sigma41,
505   NumericVector &tau41,
506   NumericVector &lambda41,
507   NumericVector &mu41,
508   NumericVector &nu41,
509   NumericVector &xi42,
510   NumericVector &eta42,
511   NumericVector &zeta42,
512   NumericVector &phi42,
513   NumericVector &psi42,
514   NumericVector &theta42,
515   NumericVector &rho42,
516   NumericVector &sigma42,
517   NumericVector &tau42,
518   NumericVector &lambda42,
519   NumericVector &mu42,
520   NumericVector &nu42,
521   NumericVector &xi43,
522   NumericVector &eta43,
523   NumericVector &zeta43,
524   NumericVector &phi43,
525   NumericVector &psi43,
526   NumericVector &theta43,
527   NumericVector &rho43,
528   NumericVector &sigma43,
529   NumericVector &tau43,
530   NumericVector &lambda43,
531   NumericVector &mu43,
532   NumericVector &nu43,
533   NumericVector &xi44,
534   NumericVector &eta44,
535   NumericVector &zeta44,
536   NumericVector &phi44,
537   NumericVector &psi44,
538   NumericVector &theta44,
539   NumericVector &rho44,
540   NumericVector &sigma44,
541   NumericVector &tau44,
542   NumericVector &lambda44,
543   NumericVector &mu44,
544   NumericVector &nu44,
545   NumericVector &xi45,
546   NumericVector &eta45,
547   NumericVector &zeta45,
548   NumericVector &phi45,
549   NumericVector &psi45,
550   NumericVector &theta45,
551   NumericVector &rho45,
552   NumericVector &sigma45,
553   NumericVector &tau45,
554   NumericVector &lambda45,
555   NumericVector &mu45,
556   NumericVector &nu45,
557   NumericVector &xi46,
558   NumericVector &eta46,
559   NumericVector &zeta46,
560   NumericVector &phi46,
561   NumericVector &psi46,
562   NumericVector &theta46,
563   NumericVector &rho46,
564   NumericVector &sigma46,
565   NumericVector &tau46,
566   NumericVector &lambda46,
567   NumericVector &mu46,
568   NumericVector &nu46,
569   NumericVector &xi47,
570   NumericVector &eta47,
571   NumericVector &zeta47,
572   NumericVector &phi47,
573   NumericVector &psi47,
574   NumericVector &theta47,
575   NumericVector &rho47,
576   NumericVector &sigma47,
577   NumericVector &tau47,
578   NumericVector &lambda47,
579   NumericVector &mu47,
580   NumericVector &nu47,
581   NumericVector &xi48,
582   NumericVector &eta48,
583   NumericVector &zeta48,
584   NumericVector &phi48,
585   NumericVector &psi48,
586   NumericVector &theta48,
587   NumericVector &rho48,
588   NumericVector &sigma48,
589   NumericVector &tau48,
590   NumericVector &lambda48,
591   NumericVector &mu48,
592   NumericVector &nu48,
593   NumericVector &xi49,
594   NumericVector &eta49,
595   NumericVector &zeta49,
596   NumericVector &phi49,
597   NumericVector &psi49,
598   NumericVector &theta49,
599   NumericVector &rho49,
600   NumericVector &sigma49,
601   NumericVector &tau49,
602   NumericVector &lambda49,
603   NumericVector &mu49,
604   NumericVector &nu49,
605   NumericVector &xi50,
606   NumericVector &eta50,
607   NumericVector &zeta50,
608   NumericVector &phi50,
609   NumericVector &psi50,
610   NumericVector &theta50,
611   NumericVector &rho50,
612   NumericVector &sigma50,
613   NumericVector &tau50,
614   NumericVector &lambda50,
615   NumericVector &mu50,
616   NumericVector &nu50,
617   NumericVector &xi51,
618   NumericVector &eta51,
619   NumericVector &zeta51,
620   NumericVector &phi51,
621   NumericVector &psi51,
622   NumericVector &theta51,
623   NumericVector &rho51,
624   NumericVector &sigma51,
625   NumericVector &tau51,
626   NumericVector &lambda51,
627   NumericVector &mu51,
628   NumericVector &nu51,
629   NumericVector &xi52,
630   NumericVector &eta52,
631   NumericVector &zeta52,
632   NumericVector &phi52,
633   NumericVector &psi52,
634   NumericVector &theta52,
635   NumericVector &rho52,
636   NumericVector &sigma52,
637   NumericVector &tau52,
638   NumericVector &lambda52,
639   NumericVector &mu52,
640   NumericVector &nu52,
641   NumericVector &xi53,
642   NumericVector &eta53,
643   NumericVector &zeta53,
644   NumericVector &phi53,
645   NumericVector &psi53,
646   NumericVector &theta53,
647   NumericVector &rho53,
648   NumericVector &sigma53,
649   NumericVector &tau53,
650   NumericVector &lambda53,
651   NumericVector &mu53,
652   NumericVector &nu53,
653   NumericVector &xi54,
654   NumericVector &eta54,
655   NumericVector &zeta54,
656   NumericVector &phi54,
657   NumericVector &psi54,
658   NumericVector &theta54,
659   NumericVector &rho54,
660   NumericVector &sigma54,
661   NumericVector &tau54,
662   NumericVector &lambda54,
663   NumericVector &mu54,
664   NumericVector &nu54,
665   NumericVector &xi55,
666   NumericVector &eta55,
667   NumericVector &zeta55,
668   NumericVector &phi55,
669   NumericVector &psi55,
670   NumericVector &theta55,
671   NumericVector &rho55,
672   NumericVector &sigma55,
673   NumericVector &tau55,
674   NumericVector &lambda55,
675   NumericVector &mu55,
676   NumericVector &nu55,
677   NumericVector &xi56,
678   NumericVector &eta56,
679   NumericVector &zeta56,
680   NumericVector &phi56,
681   NumericVector &psi56,
682   NumericVector &theta56,
683   NumericVector &rho56,
684   NumericVector &sigma56,
685   NumericVector &tau56,
686   NumericVector &lambda56,
687   NumericVector &mu56,
688   NumericVector &nu56,
689   NumericVector &xi57,
690   NumericVector &eta57,
691   NumericVector &zeta57,
692   NumericVector &phi57,
693   NumericVector &psi57,
694   NumericVector &theta57,
695   NumericVector &rho57,
696   NumericVector &sigma57,
697   NumericVector &tau57,
698   NumericVector &lambda57,
699   NumericVector &mu57,
700   NumericVector &nu57,
701   NumericVector &xi58,
702   NumericVector &eta58,
703   NumericVector &zeta58,
704   NumericVector &phi58,
705   NumericVector &psi58,
706   NumericVector &theta58,
707   NumericVector &rho58,
708   NumericVector &sigma58,
709   NumericVector &tau58,
710   NumericVector &lambda58,
711   NumericVector &mu58,
712   NumericVector &nu58,
713   NumericVector &xi59,
714   NumericVector &eta59,
715   NumericVector &zeta59,
716   NumericVector &phi59,
717   NumericVector &psi59,
718   NumericVector &theta59,
719   NumericVector &rho59,
720   NumericVector &sigma59,
721   NumericVector &tau59,
722   NumericVector &lambda59,
723   NumericVector &mu59,
724   NumericVector &nu59,
725   NumericVector &xi60,
726   NumericVector &eta60,
727   NumericVector &zeta60,
728   NumericVector &phi60,
729   NumericVector &psi60,
730   NumericVector &theta60,
731   NumericVector &rho60,
732   NumericVector &sigma60,
733   NumericVector &tau60,
734   NumericVector &lambda60,
735   NumericVector &mu60,
736   NumericVector &nu60,
737   NumericVector &xi61,
738   NumericVector &eta61,
739   NumericVector &zeta61,
740   NumericVector &phi61,
741   NumericVector &psi61,
742   NumericVector &theta61,
743   NumericVector &rho61,
744   NumericVector &sigma61,
745   NumericVector &tau61,
746   NumericVector &lambda61,
747   NumericVector &mu61,
748   NumericVector &nu61,
749   NumericVector &xi62,
750   NumericVector &eta62,
751   NumericVector &zeta62,
752   NumericVector &phi62,
753   NumericVector &psi62,
754   NumericVector &theta62,
755   NumericVector &rho62,
756   NumericVector &sigma62,
757   NumericVector &tau62,
758   NumericVector &lambda62,
759   NumericVector &mu62,
760   NumericVector &nu62,
761   NumericVector &xi63,
762   NumericVector &eta63,
763   NumericVector &zeta63,
764   NumericVector &phi63,
765   NumericVector &psi63,
766   NumericVector &theta63,
767   NumericVector &rho63,
768   NumericVector &sigma63,
769   NumericVector &tau63,
770   NumericVector &lambda63,
771   NumericVector &mu63,
772   NumericVector &nu63,
773   NumericVector &xi64,
774   NumericVector &eta64,
775   NumericVector &zeta64,
776   NumericVector &phi64,
777   NumericVector &psi64,
778   NumericVector &theta64,
779   NumericVector &rho64,
780   NumericVector &sigma64,
781   NumericVector &tau64,
782   NumericVector &lambda64,
783   NumericVector &mu64,
784   NumericVector &nu64,
785   NumericVector &xi65,
786   NumericVector &eta65,
787   NumericVector &zeta65,
788   NumericVector &phi65,
789   NumericVector &psi65,
790   NumericVector &theta65,
791   NumericVector &rho65,
792   NumericVector &sigma65,
793   NumericVector &tau65,
794   NumericVector &lambda65,
795   NumericVector &mu65,
796   NumericVector &nu65,
797   NumericVector &xi66,
798   NumericVector &eta66,
799   NumericVector &zeta66,
800   NumericVector &phi66,
801   NumericVector &psi66,
802   NumericVector &theta66,
803   NumericVector &rho66,
804   NumericVector &sigma66,
805   NumericVector &tau66,
806   NumericVector &lambda66,
807   NumericVector &mu66,
808   NumericVector &nu66,
809   NumericVector &xi67,
810   NumericVector &eta67,
811   NumericVector &zeta67,
812   NumericVector &phi67,
813   NumericVector &psi67,

```

```

10     IntegerVector &A,
11     IntegerVector &B,
12     IntegerVector &Wsx,
13     List &paramList)
14 {
15     int i, j;
16
17     // creates a Patches object to pass parameters from R
18     Patches parameters(m, v, w, C, A, B, Wsx, paramList);
19
20     parameters.computeBeta();
21     paramList["thresholdValue"] = parameters.thresholdValue;
22
23     // loop on patch index
24     for ( i = parameters.firstNewLabel - 1; i < parameters.
25           lastNewLabel; ++i )
26     {
27         parameters.split(i);
28         Rcout << "patch number " << i+1-parameters.shift << " computed
29             \n";
30         Rcout << "at iteration number " << i+1 << "\n";
31     }
32     parameters.output();
33     Rcout << "betaCoeff = " << parameters.betaCoeff << "\n";
34 }
35
36 void Patches::split(int iter) {
37     int i,j,s;
38     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
39     double crit, max_crit, Cmax_i, Cmax_j;
40     int max_i, count, max_j, int_buf;
41
42     // computes arg max C
43     max_crit = -1;
44     #pragma omp parallel private(i)
45     {
46         double my_max_crit = -1;
47         int my_max_i;
48         #pragma omp for
49         for (i=shift; i<iter-shift; ++i)
50         {
51             if ( my_max_crit < C[i] ) {
52                 my_max_crit = C[i]; // selection according to the criterion
53
54                 .
55                 my_max_i = i;
56
57             }
58         }
59     }
60 }
61
62 void Patches::output() {
63     int i,j,s;
64     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
65     double crit, max_crit, Cmax_i, Cmax_j;
66     int max_i, count, max_j, int_buf;
67
68     #pragma omp parallel private(i)
69     {
70         double my_max_crit = -1;
71         int my_max_i;
72         #pragma omp for
73         for (i=shift; i<iter-shift; ++i)
74         {
75             if ( my_max_crit < C[i] ) {
76                 my_max_crit = C[i]; // selection according to the criterion
77
78                 .
79                 my_max_i = i;
80
81             }
82         }
83     }
84 }
85
86 void Patches::computeBeta() {
87     int i,j,s;
88     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
89     double crit, max_crit, Cmax_i, Cmax_j;
90     int max_i, count, max_j, int_buf;
91
92     #pragma omp parallel private(i)
93     {
94         double my_max_crit = -1;
95         int my_max_i;
96         #pragma omp for
97         for (i=shift; i<iter-shift; ++i)
98         {
99             if ( my_max_crit < C[i] ) {
100                 my_max_crit = C[i]; // selection according to the criterion
101
102                 .
103                 my_max_i = i;
104
105             }
106         }
107     }
108 }
109
110 void Patches::split(int i) {
111     int j;
112     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
113     double crit, max_crit, Cmax_i, Cmax_j;
114     int max_i, count, max_j, int_buf;
115
116     #pragma omp parallel private(j)
117     {
118         double my_max_crit = -1;
119         int my_max_i;
120         #pragma omp for
121         for (j=shift; j<i-shift; ++j)
122         {
123             if ( my_max_crit < C[j] ) {
124                 my_max_crit = C[j]; // selection according to the criterion
125
126                 .
127                 my_max_i = j;
128
129             }
130         }
131     }
132 }
133
134 void Patches::computeBeta() {
135     int i,j,s;
136     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
137     double crit, max_crit, Cmax_i, Cmax_j;
138     int max_i, count, max_j, int_buf;
139
140     #pragma omp parallel private(i)
141     {
142         double my_max_crit = -1;
143         int my_max_i;
144         #pragma omp for
145         for (i=shift; i<iter-shift; ++i)
146         {
147             if ( my_max_crit < C[i] ) {
148                 my_max_crit = C[i]; // selection according to the criterion
149
150                 .
151                 my_max_i = i;
152
153             }
154         }
155     }
156 }
157
158 void Patches::output() {
159     int i,j,s;
160     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
161     double crit, max_crit, Cmax_i, Cmax_j;
162     int max_i, count, max_j, int_buf;
163
164     #pragma omp parallel private(i)
165     {
166         double my_max_crit = -1;
167         int my_max_i;
168         #pragma omp for
169         for (i=shift; i<iter-shift; ++i)
170         {
171             if ( my_max_crit < C[i] ) {
172                 my_max_crit = C[i]; // selection according to the criterion
173
174                 .
175                 my_max_i = i;
176
177             }
178         }
179     }
180 }
181
182 void Patches::computeBeta() {
183     int i,j,s;
184     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
185     double crit, max_crit, Cmax_i, Cmax_j;
186     int max_i, count, max_j, int_buf;
187
188     #pragma omp parallel private(i)
189     {
190         double my_max_crit = -1;
191         int my_max_i;
192         #pragma omp for
193         for (i=shift; i<iter-shift; ++i)
194         {
195             if ( my_max_crit < C[i] ) {
196                 my_max_crit = C[i]; // selection according to the criterion
197
198                 .
199                 my_max_i = i;
200
201             }
202         }
203     }
204 }
205
206 void Patches::output() {
207     int i,j,s;
208     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
209     double crit, max_crit, Cmax_i, Cmax_j;
210     int max_i, count, max_j, int_buf;
211
212     #pragma omp parallel private(i)
213     {
214         double my_max_crit = -1;
215         int my_max_i;
216         #pragma omp for
217         for (i=shift; i<iter-shift; ++i)
218         {
219             if ( my_max_crit < C[i] ) {
220                 my_max_crit = C[i]; // selection according to the criterion
221
222                 .
223                 my_max_i = i;
224
225             }
226         }
227     }
228 }
229
230 void Patches::computeBeta() {
231     int i,j,s;
232     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
233     double crit, max_crit, Cmax_i, Cmax_j;
234     int max_i, count, max_j, int_buf;
235
236     #pragma omp parallel private(i)
237     {
238         double my_max_crit = -1;
239         int my_max_i;
240         #pragma omp for
241         for (i=shift; i<iter-shift; ++i)
242         {
243             if ( my_max_crit < C[i] ) {
244                 my_max_crit = C[i]; // selection according to the criterion
245
246                 .
247                 my_max_i = i;
248
249             }
250         }
251     }
252 }
253
254 void Patches::output() {
255     int i,j,s;
256     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
257     double crit, max_crit, Cmax_i, Cmax_j;
258     int max_i, count, max_j, int_buf;
259
260     #pragma omp parallel private(i)
261     {
262         double my_max_crit = -1;
263         int my_max_i;
264         #pragma omp for
265         for (i=shift; i<iter-shift; ++i)
266         {
267             if ( my_max_crit < C[i] ) {
268                 my_max_crit = C[i]; // selection according to the criterion
269
270                 .
271                 my_max_i = i;
272
273             }
274         }
275     }
276 }
277
278 void Patches::computeBeta() {
279     int i,j,s;
280     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
281     double crit, max_crit, Cmax_i, Cmax_j;
282     int max_i, count, max_j, int_buf;
283
284     #pragma omp parallel private(i)
285     {
286         double my_max_crit = -1;
287         int my_max_i;
288         #pragma omp for
289         for (i=shift; i<iter-shift; ++i)
290         {
291             if ( my_max_crit < C[i] ) {
292                 my_max_crit = C[i]; // selection according to the criterion
293
294                 .
295                 my_max_i = i;
296
297             }
298         }
299     }
300 }
301
302 void Patches::output() {
303     int i,j,s;
304     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
305     double crit, max_crit, Cmax_i, Cmax_j;
306     int max_i, count, max_j, int_buf;
307
308     #pragma omp parallel private(i)
309     {
310         double my_max_crit = -1;
311         int my_max_i;
312         #pragma omp for
313         for (i=shift; i<iter-shift; ++i)
314         {
315             if ( my_max_crit < C[i] ) {
316                 my_max_crit = C[i]; // selection according to the criterion
317
318                 .
319                 my_max_i = i;
320
321             }
322         }
323     }
324 }
325
326 void Patches::computeBeta() {
327     int i,j,s;
328     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
329     double crit, max_crit, Cmax_i, Cmax_j;
330     int max_i, count, max_j, int_buf;
331
332     #pragma omp parallel private(i)
333     {
334         double my_max_crit = -1;
335         int my_max_i;
336         #pragma omp for
337         for (i=shift; i<iter-shift; ++i)
338         {
339             if ( my_max_crit < C[i] ) {
340                 my_max_crit = C[i]; // selection according to the criterion
341
342                 .
343                 my_max_i = i;
344
345             }
346         }
347     }
348 }
349
350 void Patches::output() {
351     int i,j,s;
352     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
353     double crit, max_crit, Cmax_i, Cmax_j;
354     int max_i, count, max_j, int_buf;
355
356     #pragma omp parallel private(i)
357     {
358         double my_max_crit = -1;
359         int my_max_i;
360         #pragma omp for
361         for (i=shift; i<iter-shift; ++i)
362         {
363             if ( my_max_crit < C[i] ) {
364                 my_max_crit = C[i]; // selection according to the criterion
365
366                 .
367                 my_max_i = i;
368
369             }
370         }
371     }
372 }
373
374 void Patches::computeBeta() {
375     int i,j,s;
376     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
377     double crit, max_crit, Cmax_i, Cmax_j;
378     int max_i, count, max_j, int_buf;
379
380     #pragma omp parallel private(i)
381     {
382         double my_max_crit = -1;
383         int my_max_i;
384         #pragma omp for
385         for (i=shift; i<iter-shift; ++i)
386         {
387             if ( my_max_crit < C[i] ) {
388                 my_max_crit = C[i]; // selection according to the criterion
389
390                 .
391                 my_max_i = i;
392
393             }
394         }
395     }
396 }
397
398 void Patches::output() {
399     int i,j,s;
400     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
401     double crit, max_crit, Cmax_i, Cmax_j;
402     int max_i, count, max_j, int_buf;
403
404     #pragma omp parallel private(i)
405     {
406         double my_max_crit = -1;
407         int my_max_i;
408         #pragma omp for
409         for (i=shift; i<iter-shift; ++i)
410         {
411             if ( my_max_crit < C[i] ) {
412                 my_max_crit = C[i]; // selection according to the criterion
413
414                 .
415                 my_max_i = i;
416
417             }
418         }
419     }
420 }
421
422 void Patches::computeBeta() {
423     int i,j,s;
424     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
425     double crit, max_crit, Cmax_i, Cmax_j;
426     int max_i, count, max_j, int_buf;
427
428     #pragma omp parallel private(i)
429     {
430         double my_max_crit = -1;
431         int my_max_i;
432         #pragma omp for
433         for (i=shift; i<iter-shift; ++i)
434         {
435             if ( my_max_crit < C[i] ) {
436                 my_max_crit = C[i]; // selection according to the criterion
437
438                 .
439                 my_max_i = i;
440
441             }
442         }
443     }
444 }
445
446 void Patches::output() {
447     int i,j,s;
448     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
449     double crit, max_crit, Cmax_i, Cmax_j;
450     int max_i, count, max_j, int_buf;
451
452     #pragma omp parallel private(i)
453     {
454         double my_max_crit = -1;
455         int my_max_i;
456         #pragma omp for
457         for (i=shift; i<iter-shift; ++i)
458         {
459             if ( my_max_crit < C[i] ) {
460                 my_max_crit = C[i]; // selection according to the criterion
461
462                 .
463                 my_max_i = i;
464
465             }
466         }
467     }
468 }
469
470 void Patches::computeBeta() {
471     int i,j,s;
472     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
473     double crit, max_crit, Cmax_i, Cmax_j;
474     int max_i, count, max_j, int_buf;
475
476     #pragma omp parallel private(i)
477     {
478         double my_max_crit = -1;
479         int my_max_i;
480         #pragma omp for
481         for (i=shift; i<iter-shift; ++i)
482         {
483             if ( my_max_crit < C[i] ) {
484                 my_max_crit = C[i]; // selection according to the criterion
485
486                 .
487                 my_max_i = i;
488
489             }
490         }
491     }
492 }
493
494 void Patches::output() {
495     int i,j,s;
496     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
497     double crit, max_crit, Cmax_i, Cmax_j;
498     int max_i, count, max_j, int_buf;
499
500     #pragma omp parallel private(i)
501     {
502         double my_max_crit = -1;
503         int my_max_i;
504         #pragma omp for
505         for (i=shift; i<iter-shift; ++i)
506         {
507             if ( my_max_crit < C[i] ) {
508                 my_max_crit = C[i]; // selection according to the criterion
509
510                 .
511                 my_max_i = i;
512
513             }
514         }
515     }
516 }
517
518 void Patches::computeBeta() {
519     int i,j,s;
520     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
521     double crit, max_crit, Cmax_i, Cmax_j;
522     int max_i, count, max_j, int_buf;
523
524     #pragma omp parallel private(i)
525     {
526         double my_max_crit = -1;
527         int my_max_i;
528         #pragma omp for
529         for (i=shift; i<iter-shift; ++i)
530         {
531             if ( my_max_crit < C[i] ) {
532                 my_max_crit = C[i]; // selection according to the criterion
533
534                 .
535                 my_max_i = i;
536
537             }
538         }
539     }
540 }
541
542 void Patches::output() {
543     int i,j,s;
544     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
545     double crit, max_crit, Cmax_i, Cmax_j;
546     int max_i, count, max_j, int_buf;
547
548     #pragma omp parallel private(i)
549     {
550         double my_max_crit = -1;
551         int my_max_i;
552         #pragma omp for
553         for (i=shift; i<iter-shift; ++i)
554         {
555             if ( my_max_crit < C[i] ) {
556                 my_max_crit = C[i]; // selection according to the criterion
557
558                 .
559                 my_max_i = i;
560
561             }
562         }
563     }
564 }
565
566 void Patches::computeBeta() {
567     int i,j,s;
568     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
569     double crit, max_crit, Cmax_i, Cmax_j;
570     int max_i, count, max_j, int_buf;
571
572     #pragma omp parallel private(i)
573     {
574         double my_max_crit = -1;
575         int my_max_i;
576         #pragma omp for
577         for (i=shift; i<iter-shift; ++i)
578         {
579             if ( my_max_crit < C[i] ) {
580                 my_max_crit = C[i]; // selection according to the criterion
581
582                 .
583                 my_max_i = i;
584
585             }
586         }
587     }
588 }
589
590 void Patches::output() {
591     int i,j,s;
592     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
593     double crit, max_crit, Cmax_i, Cmax_j;
594     int max_i, count, max_j, int_buf;
595
596     #pragma omp parallel private(i)
597     {
598         double my_max_crit = -1;
599         int my_max_i;
600         #pragma omp for
601         for (i=shift; i<iter-shift; ++i)
602         {
603             if ( my_max_crit < C[i] ) {
604                 my_max_crit = C[i]; // selection according to the criterion
605
606                 .
607                 my_max_i = i;
608
609             }
610         }
611     }
612 }
613
614 void Patches::computeBeta() {
615     int i,j,s;
616     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
617     double crit, max_crit, Cmax_i, Cmax_j;
618     int max_i, count, max_j, int_buf;
619
620     #pragma omp parallel private(i)
621     {
622         double my_max_crit = -1;
623         int my_max_i;
624         #pragma omp for
625         for (i=shift; i<iter-shift; ++i)
626         {
627             if ( my_max_crit < C[i] ) {
628                 my_max_crit = C[i]; // selection according to the criterion
629
630                 .
631                 my_max_i = i;
632
633             }
634         }
635     }
636 }
637
638 void Patches::output() {
639     int i,j,s;
640     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
641     double crit, max_crit, Cmax_i, Cmax_j;
642     int max_i, count, max_j, int_buf;
643
644     #pragma omp parallel private(i)
645     {
646         double my_max_crit = -1;
647         int my_max_i;
648         #pragma omp for
649         for (i=shift; i<iter-shift; ++i)
650         {
651             if ( my_max_crit < C[i] ) {
652                 my_max_crit = C[i]; // selection according to the criterion
653
654                 .
655                 my_max_i = i;
656
657             }
658         }
659     }
660 }
661
662 void Patches::computeBeta() {
663     int i,j,s;
664     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
665     double crit, max_crit, Cmax_i, Cmax_j;
666     int max_i, count, max_j, int_buf;
667
668     #pragma omp parallel private(i)
669     {
670         double my_max_crit = -1;
671         int my_max_i;
672         #pragma omp for
673         for (i=shift; i<iter-shift; ++i)
674         {
675             if ( my_max_crit < C[i] ) {
676                 my_max_crit = C[i]; // selection according to the criterion
677
678                 .
679                 my_max_i = i;
680
681             }
682         }
683     }
684 }
685
686 void Patches::output() {
687     int i,j,s;
688     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
689     double crit, max_crit, Cmax_i, Cmax_j;
690     int max_i, count, max_j, int_buf;
691
692     #pragma omp parallel private(i)
693     {
694         double my_max_crit = -1;
695         int my_max_i;
696         #pragma omp for
697         for (i=shift; i<iter-shift; ++i)
698         {
699             if ( my_max_crit < C[i] ) {
700                 my_max_crit = C[i]; // selection according to the criterion
701
702                 .
703                 my_max_i = i;
704
705             }
706         }
707     }
708 }
709
710 void Patches::computeBeta() {
711     int i,j,s;
712     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
713     double crit, max_crit, Cmax_i, Cmax_j;
714     int max_i, count, max_j, int_buf;
715
716     #pragma omp parallel private(i)
717     {
718         double my_max_crit = -1;
719         int my_max_i;
720         #pragma omp for
721         for (i=shift; i<iter-shift; ++i)
722         {
723             if ( my_max_crit < C[i] ) {
724                 my_max_crit = C[i]; // selection according to the criterion
725
726                 .
727                 my_max_i = i;
728
729             }
730         }
731     }
732 }
733
734 void Patches::output() {
735     int i,j,s;
736     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
737     double crit, max_crit, Cmax_i, Cmax_j;
738     int max_i, count, max_j, int_buf;
739
740     #pragma omp parallel private(i)
741     {
742         double my_max_crit = -1;
743         int my_max_i;
744         #pragma omp for
745         for (i=shift; i<iter-shift; ++i)
746         {
747             if ( my_max_crit < C[i] ) {
748                 my_max_crit = C[i]; // selection according to the criterion
749
750                 .
751                 my_max_i = i;
752
753             }
754         }
755     }
756 }
757
758 void Patches::computeBeta() {
759     int i,j,s;
760     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
761     double crit, max_crit, Cmax_i, Cmax_j;
762     int max_i, count, max_j, int_buf;
763
764     #pragma omp parallel private(i)
765     {
766         double my_max_crit = -1;
767         int my_max_i;
768         #pragma omp for
769         for (i=shift; i<iter-shift; ++i)
770         {
771             if ( my_max_crit < C[i] ) {
772                 my_max_crit = C[i]; // selection according to the criterion
773
774                 .
775                 my_max_i = i;
776
777             }
778         }
779     }
780 }
781
782 void Patches::output() {
783     int i,j,s;
784     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
785     double crit, max_crit, Cmax_i, Cmax_j;
786     int max_i, count, max_j, int_buf;
787
788     #pragma omp parallel private(i)
789     {
789         double my_max_crit = -1;
790         int my_max_i;
791         #pragma omp for
792         for (i=shift; i<iter-shift; ++i)
793         {
794             if ( my_max_crit < C[i] ) {
795                 my_max_crit = C[i]; // selection according to the criterion
796
797                 .
798                 my_max_i = i;
799
800             }
801         }
802     }
803 }
804
805 void Patches::computeBeta() {
806     int i,j,s;
807     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
808     double crit, max_crit, Cmax_i, Cmax_j;
809     int max_i, count, max_j, int_buf;
810
811     #pragma omp parallel private(i)
812     {
813         double my_max_crit = -1;
814         int my_max_i;
815         #pragma omp for
816         for (i=shift; i<iter-shift; ++i)
817         {
818             if ( my_max_crit < C[i] ) {
819                 my_max_crit = C[i]; // selection according to the criterion
820
821                 .
822                 my_max_i = i;
823
824             }
825         }
826     }
827 }
828
829 void Patches::output() {
830     int i,j,s;
831     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
832     double crit, max_crit, Cmax_i, Cmax_j;
833     int max_i, count, max_j, int_buf;
834
835     #pragma omp parallel private(i)
836     {
837         double my_max_crit = -1;
838         int my_max_i;
839         #pragma omp for
840         for (i=shift; i<iter-shift; ++i)
841         {
842             if ( my_max_crit < C[i] ) {
843                 my_max_crit = C[i]; // selection according to the criterion
844
845                 .
846                 my_max_i = i;
847
848             }
849         }
850     }
851 }
852
853 void Patches::computeBeta() {
854     int i,j,s;
855     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
856     double crit, max_crit, Cmax_i, Cmax_j;
857     int max_i, count, max_j, int_buf;
858
859     #pragma omp parallel private(i)
860     {
860         double my_max_crit = -1;
861         int my_max_i;
862         #pragma omp for
863         for (i=shift; i<iter-shift; ++i)
864         {
865             if ( my_max_crit < C[i] ) {
866                 my_max_crit = C[i]; // selection according to the criterion
867
868                 .
869                 my_max_i = i;
870
871             }
872         }
873     }
874 }
875
876 void Patches::output() {
877     int i,j,s;
878     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
879     double crit, max_crit, Cmax_i, Cmax_j;
880     int max_i, count, max_j, int_buf;
881
882     #pragma omp parallel private(i)
883     {
883         double my_max_crit = -1;
884         int my_max_i;
885         #pragma omp for
886         for (i=shift; i<iter-shift; ++i)
887         {
888             if ( my_max_crit < C[i] ) {
889                 my_max_crit = C[i]; // selection according to the criterion
890
891                 .
892                 my_max_i = i;
893
894             }
895         }
896     }
897 }
898
899 void Patches::computeBeta() {
900     int i,j,s;
901     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
902     double crit, max_crit, Cmax_i, Cmax_j;
903     int max_i, count, max_j, int_buf;
904
905     #pragma omp parallel private(i)
906     {
906         double my_max_crit = -1;
907         int my_max_i;
908         #pragma omp for
909         for (i=shift; i<iter-shift; ++i)
910         {
911             if ( my_max_crit < C[i] ) {
912                 my_max_crit = C[i]; // selection according to the criterion
913
914                 .
915                 my_max_i = i;
916
917             }
918         }
919     }
920 }
921
922 void Patches::output() {
923     int i,j,s;
924     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
925     double crit, max_crit, Cmax_i, Cmax_j;
926     int max_i, count, max_j, int_buf;
927
928     #pragma omp parallel private(i)
929     {
929         double my_max_crit = -1;
930         int my_max_i;
931         #pragma omp for
932         for (i=shift; i<iter-shift; ++i)
933         {
934             if ( my_max_crit < C[i] ) {
935                 my_max_crit = C[i]; // selection according to the criterion
936
937                 .
938                 my_max_i = i;
939
940             }
941         }
942     }
943 }
944
945 void Patches::computeBeta() {
946     int i,j,s;
947     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
948     double crit, max_crit, Cmax_i, Cmax_j;
949     int max_i, count, max_j, int_buf;
950
951     #pragma omp parallel private(i)
952     {
952         double my_max_crit = -1;
953         int my_max_i;
954         #pragma omp for
955         for (i=shift; i<iter-shift; ++i)
956         {
957             if ( my_max_crit < C[i] ) {
958                 my_max_crit = C[i]; // selection according to the criterion
959
960                 .
961                 my_max_i = i;
962
963             }
964         }
965     }
966 }
967
968 void Patches::output() {
969     int i,j,s;
970     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
971     double crit, max_crit, Cmax_i, Cmax_j;
972     int max_i, count, max_j, int_buf;
973
974     #pragma omp parallel private(i)
975     {
975         double my_max_crit = -1;
976         int my_max_i;
977         #pragma omp for
978         for (i=shift; i<iter-shift; ++i)
979         {
980             if ( my_max_crit < C[i] ) {
981                 my_max_crit = C[i]; // selection according to the criterion
982
983                 .
984                 my_max_i = i;
985
986             }
987         }
988     }
989 }
990
991 void Patches::computeBeta() {
992     int i,j,s;
993     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
994     double crit, max_crit, Cmax_i, Cmax_j;
995     int max_i, count, max_j, int_buf;
996
997     #pragma omp parallel private(i)
998     {
998         double my_max_crit = -1;
999         int my_max_i;
1000         #pragma omp for
1001         for (i=shift; i<iter-shift; ++i)
1002         {
1003             if ( my_max_crit < C[i] ) {
1004                 my_max_crit = C[i]; // selection according to the criterion
1005
1006                 .
1007                 my_max_i = i;
1008
1009             }
1010         }
1011     }
1012 }
1013
1014 void Patches::output() {
1015     int i,j,s;
1016     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
1017     double crit, max_crit, Cmax_i, Cmax_j;
1018     int max_i, count, max_j, int_buf;
1019
1020     #pragma omp parallel private(i)
1021     {
1021         double my_max_crit = -1;
1022         int my_max_i;
1023         #pragma omp for
1024         for (i=shift; i<iter-shift; ++i)
1025         {
1026             if ( my_max_crit < C[i] ) {
1027                 my_max_crit = C[i]; // selection according to the criterion
1028
1029                 .
1030                 my_max_i = i;
1031
1032             }
1033         }
1034     }
1035 }
1036
1037 void Patches::computeBeta() {
1038     int i,j,s;
1039     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
1040     double crit, max_crit, Cmax_i, Cmax_j;
1041     int max_i, count, max_j, int_buf;
1042
1043     #pragma omp parallel private(i)
1044     {
1044         double my_max_crit = -1;
1045         int my_max_i;
1046         #pragma omp for
1047         for (i=shift; i<iter-shift; ++i)
1048         {
1049             if ( my_max_crit < C[i] ) {
1050                 my_max_crit = C[i]; // selection according to the criterion
1051
1052                 .
1053                 my_max_i = i;
1054
1055             }
1056         }
1057     }
1058 }
1059
1060 void Patches::output() {
1061     int i,j,s;
1062     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
1063     double crit, max_crit, Cmax_i, Cmax_j;
1064     int max_i, count, max_j, int_buf;
1065
1066     #pragma omp parallel private(i)
1067     {
1067         double my_max_crit = -1;
1068         int my_max_i;
1069         #pragma omp for
1070         for (i=shift; i<iter-shift; ++i)
1071         {
1072             if ( my_max_crit < C[i] ) {
1073                 my_max_crit = C[i]; // selection according to the criterion
1074
1075                 .
1076                 my_max_i = i;
1077
1078             }
1079         }
1080     }
1081 }
1082
1083 void Patches::computeBeta() {
1084     int i,j,s;
1085     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
1086     double crit, max_crit, Cmax_i, Cmax_j;
1087     int max_i, count, max_j, int_buf;
1088
1089     #pragma omp parallel private(i)
1090     {
1090         double my_max_crit = -1;
1091         int my_max_i;
1092         #pragma omp for
1093         for (i=shift; i<iter-shift; ++i)
1094         {
1095             if ( my_max_crit < C[i] ) {
1096                 my_max_crit = C[i]; // selection according to the criterion
1097
1098                 .
1099                 my_max_i = i;
1100
1101             }
1102         }
1103     }
1104 }
1105
1106 void Patches::output() {
1107     int i,j,s;
1108     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
1109     double crit, max_crit, Cmax_i, Cmax_j;
1110     int max_i, count, max_j, int_buf;
1111
1112     #pragma omp parallel private(i)
1113     {
1113         double my_max_crit = -1;
1114         int my_max_i;
1115         #pragma omp for
1116         for (i=shift; i<iter-shift; ++i)
1117         {
1118             if ( my_max_crit < C[i] ) {
1119                 my_max_crit = C[i]; // selection according to the criterion
1120
1121                 .
1122                 my_max_i = i;
1123
1124             }
1125         }
1126     }
1127 }
1128
1129 void Patches::computeBeta() {
1130     int i,j,s;
1131     double m_buf, v_buf, weight_buf, square_buf, a, double_buf;
1132     double crit, max_crit, Cmax_i, Cmax_j;
1133     int max_i, count, max_j, int_buf;
1134
1135     #pragma omp parallel private(i)
1136     {
1136         double my_max_crit = -1;
1137         int my_max_i;
1138         #pragma omp for
1139         for (i=shift; i<iter-shift; ++i)
1140         {
1141             if ( my_max_crit <
```

```

50         }
51     }
52     #pragma omp critical
53     {
54         if ( max_crit < my_max_crit ) {
55             max_crit = my_max_crit;
56             max_i = my_max_i;
57         }
58     }
59 }
60
61 if (max_crit < 0) {
62     Rcout << "Nothing to split any more.\nLast split = " << iter-
63         shift << "\n";
64         ++shift;
65     return;
66 }
67
68 Rcout << "max_i = " << max_i + 1 << "\n";
69 // computes C for each j != max_i
70 max_crit = -1;
71 #pragma omp parallel private(j, count, s, weight_buf, crit, m_buf
72     ,
73     square_buf, v_buf)
74 {
75     double my_max_crit = -1;
76     int my_max_j;
77     #pragma omp for
78     for (j=shift; j<iter-shift; ++j)
79     {
80         if (j == max_i) continue; // we are looking for j != max_i
81         count = 0;
82         // compute the intersection between B_max_i and B_j.
83         for (s=0; s<d; ++s)
84         {
85             count += B[s+d*max_i] * B[s+d*j];
86         }
87         if (count == 0) continue;
88         // compute the sum of weights.
89         weight_buf = w[max_i] + w[j];
90
91         // compute the criterion for j.
92         crit = 0;

```

```

92
93     // for each pixel do
94     // compute the criterion using the law of total variance.
95     for (s=0; s<d; ++s)
96     {
97         // Test if there is an intersection.
98         if (B[s+d*max_i]*B[s+d*j] == 0) continue;
99
100        // compute the mean of the mean.
101        m_buf = (w[max_i]*m[s+d*max_i]
102            + w[j]*m[s+d*j])/weight_buf;
103        square_buf = m[s+d*max_i] - m_buf;
104
105        // compute the variance of the means.
106        square_buf *= square_buf;
107
108        // compute the means of the variance.
109        v_buf = w[max_i] * ( v[s+d*max_i] + square_buf );
110        square_buf = m_buf - m[s+d*j];
111        square_buf *= square_buf;
112        v_buf += w[j] * ( v[s+d*j] + square_buf );
113        v_buf /= weight_buf;
114        if (v_buf < thresholdValue) {
115            crit += 1;
116        }
117    }
118    crit *= betaCoeff;
119    crit -= weight_buf;
120    if (my_max_crit < crit) {
121        my_max_crit = crit;
122        my_max_j = j;
123    }
124}
125
126    // get our max_j.
127    #pragma omp critical
128    {
129        if (max_crit < my_max_crit) {
130            max_crit = my_max_crit;
131            max_j = my_max_j;
132        }
133    }
134}

```

```

136
137     if (max_crit < 0) {
138         #pragma omp parallel for private(s,double_buf,int_buf)
139         for (s=0;s<d;++s)
140     {
141         double_buf = m[s+d*shift];
142         m[s+d*shift] = m[s+d*max_i];
143         m[s+d*max_i] = double_buf;
144         double_buf = v[s+d*shift];
145         v[s+d*shift] = v[s+d*max_i];
146         v[s+d*max_i] = double_buf;
147         int_buf = B[s+d*shift];
148         B[s+d*shift] = B[s+d*max_i];
149         B[s+d*max_i] = int_buf;
150     }
151     double_buf = w[shift];
152     w[shift] = w[max_i];
153     w[max_i] = double_buf;
154     double_buf = C[shift];
155     C[shift] = C[max_i];
156     C[max_i] = double_buf;
157     #pragma omp parallel for private(i,int_buf)
158     for (i=0;i<n;++i)
159     {
160         int_buf = A[i+n*shift];
161         A[i+n*shift] = A[i+n*max_i];
162         A[i+n*max_i] = int_buf;
163     }
164     ++shift;
165     return;
166 }

167     Rcout << "max_j = " << max_j + 1 << "\n";
168     C[iter-shift] = max_crit;
169     Cmax_i = Cmax_j = 0;
170     weight_buf = w[max_i] + w[max_j];
171
172     #pragma omp parallel for private(s, square_buf) reduction(+: Cmax_i, Cmax_j)
173     for (s=0; s<d; ++s) // computes patch number iter
174     {
175         // compute patch mean using mean of m[s,i] and m[s,j].
176         m[s + d*(iter-shift)] = (w[max_i]*m[s+d*max_i]

```

```

    + w[max_j]*m[s+d*max_j])/weight_buf;

180
    square_buf = m[s + d*max_i] - m[s + d*(iter-shift)];
182    square_buf *= square_buf;
    v[s+d*(iter-shift)] = w[max_i] * ( v[s+d*max_i] + square_buf );
184    square_buf = m[s+d*max_j] - m[s+d*(iter-shift)];
    square_buf *= square_buf;
186    v[s+d*(iter-shift)] += w[max_j] * ( v[s+d*max_j] + square_buf );
    v[s+d*(iter-shift)] /= weight_buf;

188
    // Update A and B
189    // we need to compute the intersection
190    // and check that the variance is under thresholdValue.
191    if ( (B[s+d*max_i]*B[s+d*max_j] > 0) && (v[s+d*(iter-shift)]
192        < thresholdValue) )
193    {
194        B[s+d*(iter-shift)] = 1;
195        B[s+d*max_i] = 0;
196        B[s+d*max_j] = 0;
197    } else {
198        B[s+d*(iter-shift)] = 0;
199    }
200    if (B[s+d*max_i] > 0) {
201        Cmax_i += 1;
202    }
203    if (B[s+d*max_j] > 0) {
204        Cmax_j += 1;
205    }
206}
207
208    C[max_i] = Cmax_i*betaCoeff - w[max_i];
209    C[max_j] = Cmax_j*betaCoeff - w[max_j];
210    w[iter-shift] = w[max_i] + w[max_j];
211    #pragma omp parallel for private(i)
212    for (i=0; i<n; ++i)
213    {
214        A[i+n*(iter-shift)] = A[i+n*max_i] + A[i+n*max_j];
215    }
216    return;
217}
218

```

References

- [AR20] Pierre Alquier and James Ridgway. “Concentration of tempered posteriors and of their variational approximations.” In: *Ann. Statist.* 48.3 (June 2020), pp. 1475–1497. DOI: [10.1214/19-AOS1855](https://doi.org/10.1214/19-AOS1855). URL: <https://doi.org/10.1214/19-AOS1855>.
- [ARC16] Pierre Alquier, James Ridgway, and Nicolas Chopin. “On the properties of variational approximations of Gibbs posteriors.” In: *Journal of Machine Learning Research* 17.236 (2016), pp. 1–41. URL: <http://jmlr.org/papers/v17/15-290.html>.
- [Aro50] Nachman Aronszajn. “Theory of reproducing kernels.” In: *Transactions of the American mathematical society* 68.3 (1950), pp. 337–404.
- [BDG04] Arindam Banerjee, Inderjit Dhillon, and Joydeep Ghosh. “Clustering with Bregman Divergences.” In: *Journal of Machine Learning Research* 6 (June 2004). DOI: [10.1137/1.9781611972740.22](https://doi.org/10.1137/1.9781611972740.22).
- [BDL08] Gérard Biau, L. Devroye, and G. Lugosi. “On the performance of clustering in Hilbert spaces.” In: *IEEE Transactions on Information Theory* 54.2 (2008), pp. 781–790. URL: <https://hal.archives-ouvertes.fr/hal-00290855>.
- [Bis06] Christopher M. Bishop. *Pattern Recognition and Machine Learning (Information Science and Statistics)*. Secaucus, NJ, USA: Springer-Verlag New York, Inc., 2006. ISBN: 0387310738.
- [BKM17] David M. Blei, Alp Kucukelbir, and Jon D. McAuliffe. “Variational Inference: A Review for Statisticians.” In: *Journal of the American Statistical Association* 112.518 (2017), pp. 859–877. DOI: [10.1080/01621459.2017.1285773](https://doi.org/10.1080/01621459.2017.1285773). eprint: <https://doi.org/10.1080/01621459.2017.1285773>. URL: <https://doi.org/10.1080/01621459.2017.1285773>.
- [BLM13] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.
- [BFL20] Claire Brécheteau, Aurélie Fischer, and Clément Levraud. “Robust Bregman Clustering.” working paper or preprint. Apr. 2020. URL: <https://hal.archives-ouvertes.fr/hal-01948051>.

- [Cao+13] Jie Cao, Zhiang Wu, Junjie Wu, and Wenjie Liu. “Towards information-theoretic K-means clustering for image indexing.” In: *Signal Processing* 93.7 (2013), pp. 2026–2037.
- [Cat04] O. Catoni. *Statistical learning theory and stochastic optimization. Ecole d’été de probabilités de Saint-Flour XXXI-2001*. Collection : Lecture notes in mathematics n°1851. Springer, 2004, pp. viii–272. URL: <https://hal.archives-ouvertes.fr/hal-00104952>.
- [Cat12] Olivier Catoni. “Challenging the empirical mean and empirical variance: A deviation study.” In: *Ann. Inst. H. Poincaré Probab. Statist.* 48.4 (Nov. 2012), pp. 1148–1185. DOI: [10.1214/11-AIHP454](https://doi.org/10.1214/11-AIHP454). URL: <https://doi.org/10.1214/11-AIHP454>.
- [Cat14] Olivier Catoni. *Lecture notes for the IFCAM Summer School on Applied Mathematics, Indian Institute of Science*. Bangalore, 2014.
- [CG17] Olivier Catoni and Ilaria Giulini. “Dimension-free PAC-Bayesian bounds for matrices, vectors, and linear least squares regression.” In: (Dec. 2017).
- [CG18] Olivier Catoni and Ilaria Giulini. “Dimension-free PAC-Bayesian bounds for the estimation of the mean of a random vector.” In: (Feb. 2018).
- [CS08] Andreas Christmann and Ingo Steinwart. “Support vector machines.” In: (2008).
- [Coh] Donald L Cohn. *Measure theory*. Springer.
- [Csi75] I. Csiszar. “ I -Divergence Geometry of Probability Distributions and Minimization Problems.” In: *Ann. Probab.* 3.1 (Feb. 1975), pp. 146–158. DOI: [10.1214/aop/1176996454](https://doi.org/10.1214/aop/1176996454). URL: <https://doi.org/10.1214/aop/1176996454>.
- [Csi84] Imre Csiszar. “Sanov Property, Generalized I -Projection and a Conditional Limit Theorem.” In: *Ann. Probab.* 12.3 (Aug. 1984), pp. 768–793. DOI: [10.1214/aop/1176993227](https://doi.org/10.1214/aop/1176993227). URL: <https://doi.org/10.1214/aop/1176993227>.
- [Dhi+03] Dhillon, Inderjit S., Mallela, Subramanyam, Kumar, and Rahul Kumar. “A divisive information theoretic feature clustering algorithm for text classification.” In: *J. Mach. Learn. Res.* 3 (Jan. 2003), pp. 1265–.
- [Doe16] Carl Doersch. “Tutorial on Variational Autoencoders.” In: *ArXiv* abs/1606.05908 (2016).
- [Edd13] Dirk Eddelbuettel. *Seamless R and C++ Integration with Rcpp*. ISBN 978-1-4614-6867-7. New York: Springer, 2013. DOI: [10.1007/978-1-4614-6868-4](https://doi.org/10.1007/978-1-4614-6868-4).
- [EF11] Dirk Eddelbuettel and Romain François. “Rcpp: Seamless R and C++ Integration.” In: *Journal of Statistical Software* 40.8 (2011), pp. 1–18. DOI: [10.18637/jss.v040.i08](https://doi.org/10.18637/jss.v040.i08). URL: <http://www.jstatsoft.org/v40/i08/>.
- [EF13] Dirk Eddelbuettel and Romain François. “Writing a package that uses Rcpp.” In: (2013). URL: <https://cran.r-project.org/web/packages/Rcpp/vignettes/Rcpp-package.pdf>.

- [Fis10] Aurélie Fischer. “Quantization and clustering with Bregman divergences.” In: *Journal of Multivariate Analysis* 101 (Oct. 2010), pp. 2207–2221. DOI: [10.1016/j.jmva.2010.05.008](https://doi.org/10.1016/j.jmva.2010.05.008).
- [Giu15] Ilaria Giulini. “Generalization bounds for random samples in Hilbert spaces.” PhD thesis. Paris, Ecole normale supérieure, 2015.
- [Jia+11] Bin Jiang, Jian Pei, Yufei Tao, and Xuemin Lin. “Clustering uncertain data based on probability distribution similarity.” In: *IEEE Transactions on Knowledge and Data Engineering* 25.4 (2011), pp. 751–763.
- [Lev13] Clément Levraud. “Non Asymptotic Bounds for Vector Quantization in Hilbert Spaces.” 30 pages, technical proofs are omitted and can be found in the related unpublished paper ”Margin conditions for vector quantization”. Oct. 2013. URL: <https://hal.archives-ouvertes.fr/hal-00877564>.
- [Lev15] Clément Levraud. “Nonasymptotic bounds for vector quantization in Hilbert spaces.” In: *Ann. Statist.* 43.2 (Apr. 2015), pp. 592–619. DOI: [10.1214/14-AOS1293](https://doi.org/10.1214/14-AOS1293). URL: <https://doi.org/10.1214/14-AOS1293>.
- [Li+11] Teng Li, Tao Mei, Soo-Ok Kweon, and Xian-Sheng Hua. “Contextual Bag-of-Words for Visual Categorization.” In: *IEEE Trans. Circuits Syst. Video Techn.* 21 (Apr. 2011), pp. 381–392. DOI: [10.1109/TCSVT.2010.2041828](https://doi.org/10.1109/TCSVT.2010.2041828).
- [Llo06] S. Lloyd. “Least Squares Quantization in PCM.” In: *IEEE Trans. Inf. Theor.* 28.2 (Sept. 2006), pp. 129–137. ISSN: 0018-9448. DOI: [10.1109/TIT.1982.1056489](https://doi.org/10.1109/TIT.1982.1056489). URL: [http://dx.doi.org/10.1109/TIT.1982.1056489](https://doi.org/10.1109/TIT.1982.1056489).
- [Mai14] Thomas Mainguy. “Markov Substitute Processes : a statistical model for linguistics.” Theses. Université Pierre et Marie Curie - Paris VI, Dec. 2014. URL: <https://tel.archives-ouvertes.fr/tel-01127344>.
- [MPS07] Pascal Massart, Jean Picard, and École d’été de probabilités de Saint-Flour. “Concentration inequalities and model selection.” In: 2007.
- [PTL02] Fernando Pereira, Naftali Tishby, and Lillian Lee. “Distributional Clustering Of English Words.” In: *Proceedings of the 31st Annual Meeting on Association for Computational Linguistics* (May 2002). DOI: [10.3115/981574.981598](https://doi.org/10.3115/981574.981598).
- [Sal+19a] Guillaume Salha, Romain Hennequin, Viet Anh Tran, and Michalis Vazirgiannis. “A Degeneracy Framework for Scalable Graph Autoencoders.” In: *28th International Joint Conference on Artificial Intelligence (IJCAI)*. 2019.
- [Sal+19b] Guillaume Salha, Stratis Limnios, Romain Hennequin, Viet Anh Tran, and Michalis Vazirgiannis. “Gravity-Inspired Graph Autoencoders for Directed Link Prediction.” In: *ACM International Conference on Information and Knowledge Management (CIKM)*. 2019.

- [TPB01] Naftali Tishby, Fernando Pereira, and William Bialek. “The Information Bottleneck Method.” In: *Proceedings of the 37th Allerton Conference on Communication, Control and Computation* 49 (July 2001).
- [Tri16] Bertrand Clarke Tri Le. *Using the Bayesian Shtarkov solution for predictions*. Computational Statistics and Data Analysis, 2016.
- [Tsa12] Chih-Fong Tsai. “Bag-of-Words Representation in Image Annotation: A Review.” In: *International Scholarly Research Notices* 2012 (2012), pp. 1–19.
- [Wu12] Junjie Wu. *Advances in K-means clustering: a data mining thinking*. Springer Science & Business Media, 2012.