

## Original article

# Short-term associations of PM<sub>10</sub> attributed to biomass burning with respiratory and cardiovascular hospital admissions in Peninsular Malaysia

Muhammad Abdul Basit Ahmad Tajudin<sup>1,2</sup>, Lina Madaniyazi<sup>1</sup>, Xerxes Seposo<sup>1,2</sup>, Mazrura Sahani<sup>3</sup>, Aurelio Tobias<sup>4</sup>, Mohd Talib Latif<sup>5</sup>, Wan Rozita Wan Mahiyuddin<sup>6</sup>, Mohd Faiz Ibrahim<sup>7</sup>, Shingo Tamaki<sup>1</sup>, Kazuhiko Moji<sup>1</sup>, Masahiro Hashizume<sup>1,8</sup> and Chris Fook Sheng Ng<sup>1,8</sup>

<sup>1</sup>School of Tropical Medicine and Global Health (TMGH), Nagasaki University, Nagasaki, Japan, <sup>2</sup>Department of Hygiene, Department of Social Medicine, Graduate School of Medicine, Hokkaido University, Sapporo, Hokkaido, Japan, <sup>3</sup>Center for Toxicology and Health Risk Studies (CORE), Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia, <sup>4</sup>Institute of Environmental Assessment and Water Research (IDAEA), Spanish Council for Scientific Research (CSIC), Barcelona, Spain, <sup>5</sup>Department of Earth Sciences and Environment, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Selangor, Malaysia, <sup>6</sup>SEAMEO TROPED Malaysia, Institute for Medical Research, National Institutes of Health, Ministry of Health Malaysia, Shah Alam, Selangor, Malaysia, <sup>7</sup>Environmental Health Research Center, Institute for Medical Research, National Institutes of Health, Ministry of Health Malaysia, Shah Alam, Selangor, Malaysia and <sup>8</sup>Department of Global Health Policy, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo, Japan

Corresponding author. Department of Hygiene, Department of Social Medicine, Graduate School of Medicine, Hokkaido University, Kita 15 Nishi 7, Kita ward, Sapporo 060-8638, Japan. E-mail: [abdulbasit@pop.med.hokudai.ac.jp](mailto:abdulbasit@pop.med.hokudai.ac.jp)

## Abstract

**Background:** Biomass burning (BB) is a major source of air pollution and particulate matter (PM) in Southeast Asia. However, the health effects of PM smaller than 10 µm (PM<sub>10</sub>) originating from BB may differ from those of other sources. This study aimed to estimate the short-term association of PM<sub>10</sub> from BB with respiratory and cardiovascular hospital admissions in Peninsular Malaysia, a region often exposed to BB events.

**Methods:** We obtained and analyzed daily data on hospital admissions, PM<sub>10</sub> levels and BB days from five districts from 2005 to 2015. We identified BB days by evaluating the BB hotspots and backward wind trajectories. We estimated PM<sub>10</sub> attributable to BB from the excess of the moving average of PM<sub>10</sub> during days without BB hotspots. We fitted time-series quasi-Poisson regression models for each district and pooled them using meta-analyses. We adjusted for potential confounders and examined the lagged effects up to 3 days, and potential effect modification by age and sex.

**Results:** We analyzed 210 960 respiratory and 178 952 cardiovascular admissions. Almost 50% of days were identified as BB days, with a mean PM<sub>10</sub> level of 53.1 µg/m<sup>3</sup> during BB days and 40.1 µg/m<sup>3</sup> during normal days. A 10 µg/m<sup>3</sup> increment in PM<sub>10</sub> from BB was associated with a 0.44% (95% CI: 0.06, 0.82%) increase in respiratory admissions at lag 0–1, with a stronger association in adults aged 15–64 years and females. We did not see any significant associations for cardiovascular admissions.

**Conclusions:** Our findings suggest that short-term exposure to PM<sub>10</sub> from BB increased the risk of respiratory hospitalizations in Peninsular Malaysia.

**Keywords:** Biomass burning, particulate matter, respiratory diseases, Southeast Asia.

## Key Messages

We estimated daily proportions of PM<sub>10</sub> attributable to biomass burning using fire hotspots and wind trajectory information.  
PM<sub>10</sub> from biomass and non-biomass burning sources increased the risk of respiratory hospital admission.  
The effects were more immediate for PM<sub>10</sub> from biomass-burning compared to those from non-burning sources.

## Introduction

Biomass Burning (BB) is a major source of air pollution that affects human health worldwide.<sup>1</sup> BB emits large amounts of particulate matter (PM) and gaseous pollutants into the

atmosphere, causing poor air quality and health hazards.<sup>1,2</sup> Exposure to BB emissions could increase the risks of respiratory and cardiovascular diseases, hospitalizations and deaths.<sup>3–6</sup> Previous literatures have stated that BB events in

Malaysia stems from both local and transboundary sources such as land use change, agricultural expansion and forest managements.<sup>7–10</sup> These days are known as ‘haze days’ by communities in the region to describe the obscurity of the atmosphere due to BB.

Most studies have used simple methods to define BB events and estimate PM levels from BB, which may not capture the complexity and variability of BB exposure.<sup>11–15</sup> Some studies have used a fixed PM threshold ( $PM_{10} > 100 \mu\text{g}/\text{m}^3$  or  $PM_{2.5} > 35 \mu\text{g}/\text{m}^3$ ) to define a BB day,<sup>4–6</sup> or the Pollutant Index above a specific threshold,<sup>11,15</sup> while others have used a percentile threshold to identify bushfire events, such as the 90th percentile,<sup>14</sup> the 95th percentile<sup>12</sup> or the 99th percentile<sup>13</sup> of the observed PM distribution. These methods, however, may overlook smaller events with PM concentrations below the threshold or miss the variability of PM levels within BB events. They may also be influenced by other sources of PM that are not related to BB, such as traffic or industry.

As previous studies on BB/haze days in Southeast Asia have analyzed their impact on health outcomes using binary indicators,<sup>5,14</sup> they might not account for the source-specific contribution of  $PM_{10}$  from BB events, which may vary depending on the location, season and meteorology. Hence, since  $PM_{10}$  is a major pollutant during BB events and can have adverse health effects,<sup>16,17</sup> there is public health relevance in examining the health effects of  $PM_{10}$  by distinguishing between BB  $PM_{10}$  and non-BB  $PM_{10}$  while also considering the modifying effect of the events.

In this study, we examined the short-term association between source-specific  $PM_{10}$  (based on the occurrence of BB) and hospital admissions for respiratory and cardiovascular diseases in Peninsular Malaysia. To our knowledge, this investigation is the first to use a continuous measurement for BB-related  $PM_{10}$  and assess its association with hospital admissions in Southeast Asia.

## Methods

### Study location and period

We selected five major districts spanning north to south of Peninsular Malaysia (Figure 1). They include Kinta, Kuantan, Kuala Lumpur, Klang and Johor Bahru, with populations ranging from slightly below half a million in Kuantan to over 1.6 million in Kuala Lumpur (Table 1).<sup>18</sup> These districts were chosen to represent different geographical regions and urbanization levels in Peninsular Malaysia. The study period ranges from 2005 to 2015 depending on the availability of hospital admission data.

### Environmental exposure data

We obtained daily air pollution and meteorological measurements from the Continuous Air Quality Monitoring (CAQM) stations managed by the Department of Environment Malaysia. Besides  $PM_{10}$  and temperature, other gaseous pollutants such as carbon monoxide (CO), sulphur dioxide ( $\text{SO}_2$ ), nitrogen dioxide ( $\text{NO}_2$ ), ozone ( $\text{O}_3$ ), as well as relative humidity were collected for sensitivity analyses. We downloaded daily fire hotspot data from the Fire Information for Resource Management System (FIRMS) archive covering an area from 4° S to 10° N and 95° E to 120° E. The fire hotspot data were measured by the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite by the National Aeronautics and Space Administration (NASA). We selected observations with high confidence from the MCD14DL

fire identification product. We filtered for hotspots recorded with a confidence level of 80% to minimize false alarms<sup>19</sup> and as type ‘presumed vegetation fires’<sup>20</sup> to focus on BB. We included all Fire Radiative Power values to account for smaller hotspots likely associated with agricultural residue burning.<sup>21</sup>

### Health outcome data

We collected daily count of hospital admissions for respiratory diseases [International Classification of Diseases 10th revision (ICD-10): J00–J99] and cardiovascular diseases (ICD-10: I00–I99) from the Health Information Centre, Ministry of Health Malaysia. At least one major hospital in each city was included (seven in total, six state hospitals and one respiratory medical institute). The data were stratified by gender and age groups: 0–14 (child), 15–64 (adult) and 65 years and above (elderly). We paired each hospital with the closest CAQM station (within 10 km) for exposure matching.

### Identification of BB days

We used the Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPPLIT) Model by the Air Resources Laboratory (ARL), National Oceanic and Atmospheric Administration (NOAA) to estimate the backward wind trajectories from each CAQM station with a resolution of 1° latitude and longitude. The trajectories were estimated at 500, 1000 and 1500 m above ground, twice a day at 12-hours interval at 00:00 and 12:00 hours, with a duration of 72-hours to cover the distance of both local and transboundary sources within the area as most peat fires in Kalimantan burned for more than 24 hours.<sup>22</sup> The procedure was accomplished using the *splitr* package in R. To identify BB days, we merged the daily information from the MODIS fire hotspots and the HYSPPLIT backward wind trajectories. We created a buffer of 0.5° radius for all trajectory coordinates. Days with at least one buffer containing a fire hotspot were defined as ‘BB days’, while others as ‘normal days’ (Supplementary Figure S1, available as Supplementary data at IJE online). The process is automated using Python script.

### $PM_{10}$ attributable to BB

We adapted the European Union reference method<sup>23,24</sup> to estimate the daily levels of  $PM_{10}$  attributable to BB. This method had been proven to estimate source-specific PM levels in areas with natural or anthropogenic sources (i.e. dust storms or wildfires).<sup>24,25</sup> We removed the days with BB identified in the previous step, and calculated the 90-day moving average using the  $PM_{10}$  levels observed on the remaining normal days without BB. The moving average was based a 3-month period to accommodate locations that documented BB events continuously for 60 days. This provides a smoothed daily concentration to represent background  $PM_{10}$  unrelated to BB (essentially the non-BB  $PM_{10}$  on BB days).  $PM_{10}$  level attributable to BB (BB  $PM_{10}$ ) was obtained by subtracting the moving average from the total concentration observed at a given station during a BB Day.

### Statistical analysis

We conducted a two-stage analysis. We used a time series model to estimate the source-specific associations (BB  $PM_{10}$  and non-BB  $PM_{10}$ ) during BB days or normal days represented by an interaction term with the BB indicator. We chose this model to account for the potential differences in the effects of  $PM_{10}$  from BB and non-BB sources during BB days and normal days.  $PM_{10}$  with a lag duration of up to

**Figure 1.** Study districts in Peninsular Malaysia**Table 1.** Study location, period, population characteristics and hospital admissions

City <sup>a</sup>	State	Region	Study period	Population in 2010	No. of hospital beds	Respiratory admissions		Cardiovascular admissions	
						Total (n)	Daily mean (SD)	Total (n)	Daily mean (SD)
Kinta	Perak	North	2007–2014	767 794	990	43 971	15.1 (5.3)	43 670	14.9 (5.2)
Kuantan	Pahang	East coast	2009–2011	461 906	793	12 172	11.1 (4.1)	8132	7.4 (3.2)
Kuala Lumpur	Kuala Lumpur	Central	2005–2010	1 674 621	2131	62 501	28.5 (9.3)	47 656	21.7 (6.2)
Klang	Selangor	Central	2008–2010	861 189	1154	23 902	21.8 (8.7)	19 558	17.5 (5.5)
Johor Bharu	Johor	South	2009–2015	1 386 569	1206	68 414	26.8 (9.4)	60 027	23.5 (6.8)
Overall						210 960	22.2 (17.2)	178 952	18.9 (12.3)

<sup>a</sup> Ordered by latitude, north to south.  
SD, standard deviation.

four previous days was examined based on previous studies that found significant associations between PM<sub>10</sub> exposure and respiratory or cardiovascular outcomes at these

lags.<sup>16,26,27</sup> In the second stage, we performed a random-effects meta-analysis to pool effect estimates across cities.<sup>28,29</sup> Heterogeneity was described using  $I^2$ , Qstatistic and  $\tau^2$ .<sup>30</sup>

The following quasi-Poisson regression model was fitted for each district:

$$E \log(y_t) = \beta_0 + \beta_1 \text{BBPM}_{10,t} + \beta_2 \text{nonBBPM}_{10,t} + \beta_3 \text{BBday}_t + \beta_4 \text{BBday}_t \text{nonBBPM}_{10,t} + \beta_5 \text{time; df} = 3 + \beta_6 \text{number of years} + \beta_7 \text{bs temp}_0 + \beta_8 I(\text{temp} > \text{median}) \text{knots}_{66\text{pct}} + \beta_9 \text{bs temp}_{ma3} + \beta_{10} I(\text{temp}_{ma3} < \text{median}) + \beta_{11} \text{knots}_{33\text{pct}} + \beta_{12} \text{holiday}_t + \beta_{13} \text{H1N1}_t + \beta_{14} \text{myhdw}_t + \beta_{15} \text{DOW}_t^k \quad (1)$$

where:  $y_t$  is the daily counts of admission on day  $t$ ;  $\text{BBPM}_{10}$  is the BB-related  $\text{PM}_{10}$  exposure variable on lag day  $l$ , representing the number of cumulative lags (i.e. lag 0–1, 0–2, 0–3 and 0–4);  $\text{nonBBPM}_{10}$  is the non-BB-related  $\text{PM}_{10}$  exposure variable;  $\text{BBday}$  is the binary indicator variable for days with BB events;  $\text{time}$  is calendar day smoothed using natural cubic spline ( $ns$ ) with three degrees of freedom ( $df$ ) per year to account for seasonality and long-term trend;  $\text{temp}_0$  and  $\text{temp}_{ma3}$  represent the current day (lag 0) and the average of the previous 3 days temperature (lag 1–3), respectively; and  $I(x)$  is an indicator function with a value of 1 when condition  $x$  is true; a quadratic B-spline ( $bs$ ) with one internal knot placed at the 33rd (for low temperature) or 66th percentile (for high temperature) was used to smooth the temperature terms, separately for high (above median value) and low temperatures (below median value) to allow possible delayed and nonlinear pattern;<sup>23–25</sup>  $\text{holiday}_t$  is an indicator variable for public holidays;  $\text{H1N1}_t$  is an indicator variable for influenza A (H1N1) outbreak in 2009;  $\text{myhdw}_t$  is an indicator variable for a change in reporting system by the Ministry of Health Malaysia after 2012; and  $\text{DOW}_t^k$  is an indicator variable for  $k$  days of the week. We also examined possible effect modification by age and gender.<sup>4–6,31</sup> We used R version 4.0.4<sup>32</sup> and *meta* package.<sup>33</sup> All results are reported as percent increases in risk (%IR) of admissions relative to a 10-  $\mu\text{g}/\text{m}^3$  increment in  $\text{PM}_{10}$  with 95% confidence intervals (CI).

### Additional analysis

Additional sensitivity analysis methods can be found in the [Supplementary Analyses](#) (available as [Supplementary data](#) at *IJE* online). We also used Bing Chat<sup>34</sup> to assist in revising our

article, specifically employing the prompt ‘improve the flow of this paragraph as a native speaker’ to generate suggestions for enhancing the English language. We acknowledge that the final responsibility for the content and quality of our article rests with us, the authors. Bing chat can be accessed at <https://www.bing.com/chat>.

## Results

### Descriptive results

During the study, we observed 210 960 respiratory and 178 952 cardiovascular hospital admissions, averaging 22.2 and 18.9 daily cases ([Table 1](#)). Children (49.9%), adults (33.8%) and the elderly (16.8%) were the primary groups for respiratory admissions, while for cardiovascular cases, the majority were adults (64.4%), followed by the elderly (33.1%) and children (2.5%). Males represented a higher percentage of cases in both categories ([Supplementary Table S1](#), available as [Supplementary data](#) at *IJE* online). BB events occurred on 4868 days, nearly half the study period, with  $\text{PM}_{10}$  levels averaging 53.1  $\mu\text{g}/\text{m}^3$  on BB days vs 40.1  $\mu\text{g}/\text{m}^3$  on non-BB days. Alternative BB day definitions based on fixed ( $\text{PM}_{10} = 100 \mu\text{g}/\text{m}^3$ ) or percentile thresholds ( $\text{PM}_{10}$  95th percentile) significantly reduced the BB day count to 1.9% and 5% ([Table 2](#)).

### Association between PM exposures and hospital admissions

[Table 3](#) presents the estimated risks of hospital admissions for exposure to source-specific  $\text{PM}_{10}$ . For respiratory admissions, exposure to BB  $\text{PM}_{10}$  was associated with 0.44% (95% CI: 0.06%, 0.80%) higher for cumulative lag 0–1 days. We did not find any associations for non-BB  $\text{PM}_{10}$  on BB days. On normal days, the estimated risk increase for exposure to non-BB  $\text{PM}_{10}$  was larger and more delayed up to lag 0–4 days (%IR 3.14%; 95% CI: 2.13%, 4.15%). For cardiovascular admissions, we did not find any significant risk increases, except for exposure to non-BB  $\text{PM}_{10}$  on BB days where risk decreases were observed for lag 0–2 and 0–3. The location-specific risk estimates appeared homogeneous for both admission causes (all  $P$ -values for  $Q > 0.05$ ).

We also examined possible effect modification by age and gender subgroups. [Figure 2](#) shows the results stratified by age and gender subgroups. Exposure to BB  $\text{PM}_{10}$  was associated with increased respiratory admissions in the adult and female subgroups with %IR of 0.76% (95% CI: 0.09%, 1.43%) and 0.57% (95% CI: 0.39%, 0.75%), respectively. On normal

**Table 2.** Summary of biomass burning occurrences and the corresponding daily  $\text{PM}_{10}$  concentration in the five study locations in Peninsular Malaysia

Location <sup>a</sup>	Total days	Days with biomass burning <sup>b</sup>						Normal days (no biomass burning)					
		No. of days identified, $n$ (%)	$\text{PM}_{10}$ concentration ( $\mu\text{g}/\text{m}^3$ )					No. of days identified, $n$ (%)	$\text{PM}_{10}$ concentration ( $\mu\text{g}/\text{m}^3$ )				
			Mean (SD)	Min	P25	P75	Max		Mean (SD)	Min	P25	P75	Max
Kinta	2922	1244 (42.6)	50.1 (18.2)	21.7	38.2	57.2	205.5	1678 (57.4)	37.9 (8.5)	13.3	32.0	42.8	85.6
Kuantan	1095	518 (47.3)	40.7 (11.8)	14.7	32.6	46.9	108.4	577 (52.7)	31.3 (7.6)	12.8	26.2	36.1	66.9
Kuala Lumpur	2191	1112 (50.8)	56.3 (28.6)	18.3	42.5	62.7	465.4	1079 (49.3)	41.7 (11.4)	13.1	34.0	48.5	130.2
Klang	1096	530 (48.4)	71.1 (26.4)	21.5	54.5	82.2	194.0	566 (51.6)	55.7 (15.5)	19.3	46.3	63.8	157.1
Johor Bahru	2556	1467 (57.4)	51.1 (25.9)	18.7	37.1	56.9	348.6	1089 (42.6)	38.5 (11.2)	12.4	31.0	44.4	99.6
Overall	9860	4868 (49.4)	53.1 (24.9)	14.7	38.8	60.3	465.4	4992 (50.6)	40.1 (12.4)	12.4	32.0	46.1	157.1

<sup>a</sup> Ordered by latitude, north to south.

<sup>b</sup> Days with biomass burning are defined by the presence of Moderate Resolution Imaging Spectroradiometer (MODIS) satellite fire hotspots and supported by the Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPLIT) Model wind trajectory. P25, 25th percentile; P75, 75th percentile; PM, particulate matter; SD, standard deviation.

**Table 3.** Estimated percent increase in risk (IR) (95% CI) for respiratory and cardiovascular hospital admissions associated with biomass burning PM<sub>10</sub> and non-biomass burning PM<sub>10</sub> during biomass burning days<sup>a</sup> and non-biomass burning PM<sub>10</sub> during normal days

Hospital admission	Lag days	BB PM <sub>10</sub> <sup>a</sup>					Non-BB PM <sub>10</sub> during BB day <sup>a</sup>					Non-BB PM <sub>10</sub> during normal day <sup>a</sup>				
		% IR (95% CI) <sup>b</sup>	I <sup>2</sup> (%)	<sup>2</sup>	Q P		% IR (95% CI) <sup>b</sup>	I <sup>2</sup> (%)	<sup>2</sup>	Q P		% IR (95% CI) <sup>b</sup>	I <sup>2</sup> (%)	<sup>2</sup>	Q P	
Respiratory	0–1	0.44 (0.06, 0.80)	7	<0.01	0.37		–0.04 (–0.73, 0.65)	0	0	0.82		0.86 (0.31, 1.41)	0	0	0.88	
	0–2	0.39 (–0.18, 0.97)	37	<0.01	0.18		0.25 (–0.83, 1.33)	0	<0.01	0.41		1.34 (0.25, 2.44)	0	0	0.57	
	0–3	0.32 (–0.37, 1.02)	49	<0.01	0.10		0.33 (–1.20, 1.87)	47	<0.01	0.11		2.35 (1.44, 3.25)	0	0	0.85	
	0–4	0.29 (–0.54, 1.13)	55	<0.01	0.07		0.3 (–1.11, 1.72)	38	<0.01	0.17		3.14 (2.13, 4.15)	0	0	0.87	
Cardiovascular	0–1	0.05 (–0.50, 0.60)	40	<0.01	0.16		–0.49 (–1.75, 0.79)	17	<0.01	0.31		0.10 (–1.10, 1.30)	40	<0.01	0.15	
	0–2	–0.02 (–0.70, 0.70)	46	<0.01	0.12		–0.86 (–1.61, –0.11)	0	0	0.72		0.07 (–1.60, 1.80)	47	<0.01	0.11	
	0–3	–0.10 (–0.88, 0.68)	45	<0.01	0.12		–0.64 (–1.19, –0.09)	0	0	0.89		0.13 (–1.78, 2.08)	39	<0.01	0.16	
	0–4	–0.20 (–1.06, 0.68)	51	<0.01	0.09		–0.71 (–1.43, 0.02)	0	0	0.77		0.69 (–1.20, 2.62)	21	<0.01	0.28	

<sup>a</sup> The estimates for non-biomass burning and biomass burning PM<sub>10</sub> were estimated from three-sources model adjusted for the other PM source in turn, and with interaction of biomass burning days with non-biomass burning PM<sub>10</sub>.

<sup>b</sup> Percent increase in risk (% IR) per 10 µg/m<sup>3</sup> increase in exposure.  
P < 0.05.

BB, biomass burning; CI, confidence interval; PM, particulate matter; I<sup>2</sup>, heterogeneity among location-specific estimates; <sup>2</sup>, variance of the true effect sizes; Q P value, Cochran's Q Test.

**Figure 2.** Estimated percent increase in risk (IR) (95% CI) of respiratory admissions (white symbol) and cardiovascular admissions (black symbol) associated with 10 µg/m<sup>3</sup> increase in biomass burning PM<sub>10</sub> (a), non-biomass burning PM<sub>10</sub> during biomass burning days (b), and non-biomass burning PM<sub>10</sub> during normal days (c) for lag 0–1 days, stratified by age and gender

days, respiratory admissions showed a positive association with exposure to non-BB PM<sub>10</sub> amongst the females at %IR of 1.35% (95% CI: 0.13%, 2.58%). This association was not observed during BB days. For cardiovascular admissions, we did not find any associations across the different subgroups.

Our study found that different degrees of freedom didn't significantly affect quasi-AIC values (Supplementary Table S3, available as Supplementary data at IJE online), and buffer size changes mostly yielded consistent results, except for non-BB PM<sub>10</sub> on BB days which varied by admission cause (Supplementary Figures S2 and S3, available as Supplementary data at IJE online). Larger buffers increased respiratory risks but not cardiovascular ones. Additional adjustments for gases and humidity had minimal impact (Supplementary Figures S4 and S5, available as Supplementary data at IJE online). Different methods for defining BB days, like fixed or percentile

thresholds, altered results compared to the HYSPLIT-MODIS method, with fixed thresholds indicating increased risks for BB and non-BB PM<sub>10</sub> (Supplementary Figures S6 and S7, available as Supplementary data at IJE online).

## Discussion

In this study, we examined the associations between PM<sub>10</sub> attributable to BB and hospital admissions in Peninsular Malaysia. We used a novel exposure definition that allowed us to disentangle the associations between PM<sub>10</sub> and hospital admissions attributable to BB and non-BB sources during BB days and normal days. We found that respiratory admissions were associated with BB PM<sub>10</sub> and non-BB PM<sub>10</sub> during normal days. The lag time was relatively shorter for BB PM<sub>10</sub> (0–1 days) and longer for non-BB PM<sub>10</sub> during normal days

(0–4 days). Subgroup analysis showed evidence of associations in adults aged 15–64 years old and females. We did not observe consistent associations for cardiovascular disease admissions.

Our results suggest that both BB PM<sub>10</sub> and non-BB PM<sub>10</sub> can have an adverse effect on respiratory admissions, with BB PM<sub>10</sub> having a more immediate impact while non-BB PM<sub>10</sub> has a longer lag association. The difference in the size of the estimated effects and the lag structure between BB PM<sub>10</sub> and non-BB PM<sub>10</sub> may be related to the BB/haze warning systems implemented in Malaysia. The Department of Environment and mass media provide alerts for BB/haze days in Malaysia by updates its website with API and BB/haze warnings<sup>35</sup> and news highlights during a BB/haze day. These BB/haze warning systems could lead to changes in behavior, such as staying indoors and reducing exposure, during BB/haze days. However, even though API calculations from monitoring stations are published daily, PM<sub>10</sub> concentrations during normal days may be neglected due to media focus on BB/haze days, resulting in delayed associations in our models. Another possible explanation for the associations with respiratory admissions could be the toxicity of source specific PM<sub>10</sub>. A study by de Oliveira Alves *et al.* (2017) also suggests that exposure to PM<sub>10</sub> which is also abundant during non-bb days can cause significant damage to human lung cells.<sup>36</sup>

We did not find a consistent association between PM<sub>10</sub> exposure and cardiovascular disease admissions. Negative associations were observed for lag 0–2 and 0–3 with non-BB PM<sub>10</sub> during BB days. One possible reason for the association is that people who have existing risk factors for CVD (i.e. hypertension, high cholesterol) may have other underlying conditions and are aware of their risks thus reducing their time outside even more. The National Health and Morbidity Survey 2019 in Malaysia showed that approximately 1.7 million (8.1%) people currently live with three major risk factors (High cholesterol, Hypertension and Diabetes) and approximately 3.4 million (16%) live with two major risk factors.<sup>37</sup> Compared to respiratory disease, the results for cardiovascular disease admissions were inconsistent when we used different buffer sizes in the sensitivity analysis to identify BB days. Previous studies have also reported inconsistent results regarding the association between BB events or PM<sub>10</sub> exposure and cardiovascular outcomes.<sup>13,38–40</sup> This lack of association between cardiovascular hospital admissions and PM<sub>10</sub> exposure could be due to the immediate onset of cardiovascular diseases. The association between PM<sub>10</sub> and cardiovascular outcomes might be better reflected using emergency room visitation data rather than hospital admission data as the severity of a cardiovascular disease onset might cause it to be treated immediately by emergency personnel,<sup>41–43</sup> with only severe cases being admitted to the hospital.

Our subgroup analysis found a significantly positive association between BB PM<sub>10</sub> and the adult subgroup. The Department of Statistics Malaysia, (2010) recorded that the highest percentage of occupations for adults in Malaysia are in service and sales (19.3%), elementary occupations (12.2%) and plant and machine operation (11.7%).<sup>18</sup> These occupations often involve outdoor labour, travel, construction or manual labour and have higher effect estimates of air pollutants compared to white-collar jobs.<sup>44</sup> This prolonged outdoor exposure, and pre-existing medical conditions may increase the risk of disease. We also found significantly positive risk estimates among females, which is consistent with

some literature that found higher risks for respiratory/asthma diseases in females.<sup>40,42,45,46</sup> This may be due to hormonal factors unique to females,<sup>47</sup> such as the pre-menstrual or menstrual phase of their monthly cycle which could increase the breathing rate,<sup>48</sup> exacerbate asthma symptoms,<sup>49</sup> and reduce lung function.<sup>50</sup>

Our study has several strengths. Firstly, we used a multi-criteria method to improve the identification of BB days. This process is computationally intensive as it involves calculating many back trajectories to provide daily coordinates for detecting fire hotspots at three different altitudes. Secondly, to our knowledge, this is the first study in Malaysia to distinguish the effects of source specific PM<sub>10</sub> from BB and examine their short-term associations with respiratory and cardiovascular hospital admissions. Our results suggest that the effects of BB PM<sub>10</sub> and non-BB PM<sub>10</sub> were consistent across Peninsular Malaysia and that the pooled results can be used to represent the region. Our study also has several limitations. Firstly, the definition of BB PM<sub>10</sub> we used is an adaptation from desert dust definitions by Escudero *et al.*<sup>51</sup> and Tobias and Stafoggia,<sup>24</sup> and there is no existing knowledge on how to assess PM<sub>10</sub> using chemical source apportionment data during BB events in Southeast Asia. Secondly, we used ground monitor PM<sub>10</sub> concentrations to reflect exposure, but we cannot accurately determine if patients were exposed to the same daily PM<sub>10</sub> concentration throughout the study period. Thirdly, we were unable to control for pre-existing health conditions in our model due to a lack of data. Fourthly, while we filtered for 'presumed vegetation fires', some non-vegetation fires may have been included in our analysis. Finally, our results should not be generalized to other locations with different topographical, meteorological and fire sources.

## Conclusion

Our study suggests that exposure to PM<sub>10</sub> from BB is associated with increased respiratory hospital admissions in Peninsular Malaysia, especially among adults and females. The effect of BB PM<sub>10</sub> is more immediate than that of non-BB PM<sub>10</sub>, which has a longer lag association. We did not find consistent associations between PM<sub>10</sub> exposure and cardiovascular hospital admissions. Our findings have implications for public health policy and intervention, as they highlight the need to monitor and reduce the exposure to PM<sub>10</sub> from BB, especially during BB events, and to raise the awareness of the health risks of PM<sub>10</sub> from non-BB sources during normal days. Future research could include using Chemical Transport Model to identify and measure BB PM<sub>10</sub> more accurately, as well as the potential interactions with other air pollutants or risk factors.

## Ethics approval

This study was registered with the National Medical Research Register (NMRR-21-71-57986) and received approval from the Ministry of Health Research Ethics Committee (KKM/NIHSEC/P21-229(2)) prior to data collection.

## Data availability

Restrictions apply to the availability of the health and air pollution data where access requires permission from the Ministry of Health Malaysia and the Department of

Environment Malaysia of the Ministry of Natural Resources while the NASA—FIRMS MODIS fire archive is available from the following website: <http://firms.modaps.eosdis.nasa.gov/download/>. The NOAA—ARL HYSPLIT wind trajectory is available from the following website: <https://www.ready.noaa.gov/HYSPLIT.php>.

## Supplementary data

Supplementary data are available at *IJE* online.

## Author contributions

M.A.B.A.T. contributed to data management, drafted the manuscript and final revision. M.A.B.A.T. and C.F.S.N. contributed to the study design, data analysis and manuscript writing. S.T. wrote the algorithm in Python to automatically identify biomass burning days. L.M., X.S., M.S., A.T., M.T. L., K.M., M.F.I., W.R.W.M. and M.H. contributed to the refinement of the protocol and provided edits to the manuscript for intellectual content.

## Funding

This work was supported by JSPS KAKENHI [grant numbers JP19K10644].

## Acknowledgements

We extend our sincere appreciation to the Director General of Health Malaysia for granting permission to publish this article, and to the Malaysia Department of Environment for providing the air quality data. Special thanks are also due to the Health Informatics Centre, Ministry of Health, for providing hospital admissions data.

## Conflict of interest

None declared.

## References

- Latif MT, Othman M, Idris N *et al*. Impact of regional haze towards air quality in Malaysia: a review. *Atmos Environ* 2018; **177**:28–44.
- Vadrevu KP, Lasko K, Giglio L, Schroeder W, Biswas S, Justice C. Trends in vegetation fires in South and Southeast Asian countries. *Sci Rep* 2019; **9**:7422.
- American Lung Association. *The Health Dangers of Wildfires Are Well Known, Why Not Biomass?* 2016. <https://www.lung.org/blog/dangers-wildfire-known-biomass> (22 June 2023, date last accessed).
- Phung VLH, Ueda K, Sahani M *et al*. Investigation of association between smoke haze and under-five mortality in Malaysia, accounting for time lag, duration and intensity. *Int J Epidemiol* 2022; **51**:155–65.
- Sahani M, Zainon NA, Wan Mahiyuddin WR *et al*. A case-crossover analysis of forest fire haze events and mortality in Malaysia. *Atmos Environ* 2014; **96**:257–65.
- Sulong NA, Latif MT, Khan MF *et al*. Source apportionment and health risk assessment among specific age groups during haze and non-haze episodes in Kuala Lumpur, Malaysia. *Sci Total Environ* 2017; **601–602**:556–70.
- Biswas S, Vadrevu KP, Lwin ZM, Lasko K, Justice CO. Factors controlling vegetation fires in protected and non-protected areas of Myanmar. *PLoS One* 2015; **10**:e0124346.
- Dung NV, Vien TD, Lam NT, Tuong TM, Cadisch G. Analysis of the sustainability within the composite Swidden agroecosystem in northern Vietnam - 1. Partial nutrient balances and recovery times of upland fields. *Agr Ecosyst Environ* 2008; **128**:37–51.
- Inoue Y. Ecosystem carbon stock, atmosphere, and food security in slash-and-burn land use: a geospatial study in mountainous region of Laos. In: Vadrevu K, Ohara T, Justice, C (eds). *Land-Atmospheric Research Applications in South and Southeast Asia*, Springer Remote Sensing/Photogrammetry. Springer, Cham, 2018, pp. 641–65.
- Tacconi L, Vayda AP. Slash and burn and fires in Indonesia: a comment. *Ecol Econ* 2006; **56**:1–4.
- Ho AFW, Wah W, Earnest A *et al*. Singapore PAROS Investigator. Health impacts of the Southeast Asian haze problem—a time-stratified case crossover study of the relationship between ambient air pollution and sudden cardiac deaths in Singapore. *Int J Cardiol* 2018; **271**:352–58.
- Johnston F, Hanigan I, Henderson S, Morgan G, Bowman D. Extreme air pollution events from bushfires and dust storms and their association with mortality in Sydney, Australia 1994–2007. *Environ Res* 2011; **111**:811–16.
- Johnston FH, Purdie S, Jalaludin B, Martin KL, Henderson SB, Morgan GG. Air pollution events from forest fires and emergency department attendances in Sydney, Australia 1996–2007: a case-crossover analysis. *Environ Health* 2014; **13**:105.
- Mueller W, Loh M, Vardoulakis S *et al*. Ambient particulate matter and biomass burning: an ecological time series study of respiratory and cardiovascular hospital visits in northern Thailand. *Environ Health* 2020; **19**:77.
- Othman J, Sahani M, Mahmud M, Ahmad MK. Transboundary smoke haze pollution in Malaysia: inpatient health impacts and economic valuation. *Environ Pollut* 2014; **189**:194–201.
- Phosri A, Ueda K, Phung VLH, Tawatsupa B, Honda A, Takano H. Effects of ambient air pollution on daily hospital admissions for respiratory and cardiovascular diseases in Bangkok, Thailand. *Sci Total Environ* 2019; **651**:1144–53.
- Uttajug A, Ueda K, Oyoshi K, Honda A, Takano H. Association between PM<sub>10</sub> from vegetation fire events and hospital visits by children in upper northern Thailand. *Sci Total Environ* 2021; **764**:142923.
- Department of Statistics, Malaysia. Population Distribution and Basic Demographic Characteristics. 2010. <https://www.mycensus.gov.my/index.php/ms/produk-banci/penerbitan/banci-2010/664-taburan-penduduk-dan-ciri-ciri-asas-demografi-2010>. (28 June 2023, date last accessed).
- L. G. MODIS Collection 6 Active Fire Product User's Guide Revision B. 2018. [http://modis-fire.umd.edu/files/MODIS\\_C6\\_Fire\\_User\\_Guide\\_B.pdf](http://modis-fire.umd.edu/files/MODIS_C6_Fire_User_Guide_B.pdf) (19 August 2020, date last accessed).
- Giglio LSW, Hall JV, Justice CO. MODIS Collection 6 Active Fire Product User's Guide Revision C. 2020. [https://modis-fire.umd.edu/files/MODIS\\_C6\\_Fire\\_User\\_Guide\\_C.pdf](https://modis-fire.umd.edu/files/MODIS_C6_Fire_User_Guide_C.pdf). (December 2020, date last accessed).
- Xu WD, Wooster MJ, Kaneko T, He JP, Zhang TR, Fisher D. Major advances in geostationary fire radiative power (FRP) retrieval over Asia and Australia stemming from use of Himawari-8 AHI. *Remote Sens Environ* 2017; **193**:138–49.
- Uda SK, Hein L, Atmoko D. Assessing the health impacts of peatland fires: a case study for Central Kalimantan, Indonesia. *Environ Sci Pollut Res Int* 2019; **26**:31315–27.
- Stafoggia M, Zauli-Sajani S, Pey J *et al*. MED-PARTICLES Study Group. Desert dust outbreaks in southern Europe: contribution to daily PM<sub>10</sub> concentrations and short-term associations with mortality and hospital admissions. *Environ Health Perspect* 2016; **124**:413–19.
- Tobias A, Stafoggia M. Modeling desert dust exposures in epidemiologic short-term health effects studies. *Epidemiology* 2020; **31**:788–95.

25. Querol X, Tobias A, Perez N *et al.* Monitoring the impact of desert dust outbreaks for air quality for health studies. *Environ Int* 2019; **130**:104867.
26. Ma Y, Zhao Y, Yang S *et al.* Short-term effects of ambient air pollution on emergency room admissions due to cardiovascular causes in Beijing, China. *Environ Pollut* 2017; **230**:974–80.
27. Phung D, Hien TT, Linh HN *et al.* Air pollution and risk of respiratory and cardiovascular hospitalizations in the most populous city in Vietnam. *Sci Total Environ* 2016; **557–558**:322–30.
28. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 2010; **1**:97–111.
29. Viechtbauer W. Bias and efficiency of meta-analytic variance estimators in the random-effects model. *J Educ Behav Stat* 2005; **30**:261–93.
30. Moss RH, Edmonds JA, Hibbard KA *et al.* The next generation of scenarios for climate change research and assessment. *Nature* 2010; **463**:747–56.
31. Ibrahim MF, Hod R, Ahmad Tajudin MAB, Wan Mahiyuddin WR, Mohammed Nawi A, Sahani M. Children's exposure to air pollution in a natural gas industrial area and their risk of hospital admission for respiratory diseases. *Environ Res* 2022; **210**:112966.
32. RC Team. *R: A Language and Environment for Statistical Computing: R Foundation for Statistical Computing*. 2021. <https://www.r-project.org/> (28 June 2023, date last accessed).
33. Harrer M, Cuijpers P, Furukawa T, Ebert D. *Doing Meta-Analysis with R: A Hands-on Guide*, 1st ed. New York: Chapman and Hall/CRC, 2021.
34. Microsoft. *Bing Chat (June 28 Version) [Large Language Model]*. 2023. <https://www.bing.com/chat> (28 June 2023, date last accessed).
35. Department of Environment, Malaysia. *Air Pollutant Index Management System: Department of Environment, Ministry of Environment & Water*. 2022. [http://apims.doe.gov.my/public\\_v2/home.html](http://apims.doe.gov.my/public_v2/home.html) (28 June 2023, date last accessed).
36. de Oliveira Alves N, Vessoni AT, Quinet A *et al.* Biomass burning in the Amazon region causes DNA damage and cell death in human lung cells. *Sci Rep* 2017; **7**:10937.
37. Institute for Public Health. *National Health and Morbidity Survey (NHMS) 2019: Non-Communicable Diseases, Healthcare Demand, and Health Literacy—Key Findings*. 2020. <https://iku.moh.gov.my/nhms-2019> (28 June 2023, date last accessed).
38. Hutchinson JA, Vargo J, Milet M *et al.* The San Diego 2007 wildfires and Medi-Cal emergency department presentations, inpatient hospitalizations, and outpatient visits: an observational study of smoke exposure periods and a bidirectional case-crossover analysis. *PLoS Med* 2018; **15**:e1002601.
39. Martin KL, Hanigan IC, Morgan GG, Henderson SB, Johnston FH. Air pollution from bushfires and their association with hospital admissions in Sydney, Newcastle and Wollongong, Australia 1994–2007. *Aust N Z J Public Health* 2013; **37**:238–43.
40. Reid CE, Jerrett M, Tager IB, Petersen ML, Mann JK, Balmes JR. Differential respiratory health effects from the 2008 northern California wildfires: a spatiotemporal approach. *Environ Res* 2016; **150**:227–35.
41. Dennekamp M, Straney LD, Erbas B *et al.* Forest fire smoke exposures and out-of-hospital cardiac arrests in Melbourne, Australia: a case-crossover study. *Environ Health Perspect* 2015; **123**:959–64.
42. Rappold AG, Stone SL, Cascio WE *et al.* Peat bog wildfire smoke exposure in rural North Carolina is associated with cardiopulmonary emergency department visits assessed through syndromic surveillance. *Environ Health Perspect* 2011; **119**:1415–20.
43. Wettstein ZS, Hoshiko S, Fahimi J, Harrison RJ, Cascio WE, Rappold AG. Cardiovascular and cerebrovascular emergency department visits associated with wildfire smoke exposure in California in 2015. *J Am Heart Assoc* 2018; **7**:e007492.
44. Ou C-Q, Hedley AJ, Chung RY *et al.* Socioeconomic disparities in air pollution-associated mortality. *Environ Res* 2008; **107**:237–44.
45. Haikerwal A, Akram M, Sim MR, Meyer M, Abramson MJ, Dennekamp M. Fine particulate matter PM2.5 exposure during a prolonged wildfire period and emergency department visits for asthma. *Respirology* 2016; **21**:88–94.
46. Stowell JD, Geng G, Saikawa E *et al.* Associations of wildfire smoke PM2.5 exposure with cardiorespiratory events in Colorado 2011–2014. *Environ Int* 2019; **133**:105151.
47. Troisi RJ, Speizer FE, Willett WC, Trichopoulos D, Rosner B. Menopause, postmenopausal estrogen preparations, and the risk of adult-onset asthma. A prospective cohort study. *Am J Respir Crit Care Med* 1995; **152**:1183–88.
48. Gargaglioni LH, Marques DA, Patrone LGA. Sex differences in breathing. *Comp Biochem Physiol A Mol Integr Physiol* 2019; **238**:110543.
49. Pignataro FS, Bonini M, Forgione A, Melandri S, Usmani OS. Asthma and gender: the female lung. *Pharmacol Res* 2017; **119**:384–90.
50. Real FG, Svanes C, Omenaas ER *et al.* Menstrual irregularity and asthma and lung function. *J Allergy Clin Immunol* 2007; **120**:557–64.
51. Escudero M, Querol X, Pey J *et al.* A methodology for the quantification of the net African dust load in air quality monitoring networks. *Atmos Environ* 2007; **41**:5516–24.