


Solving Linear Systems

(Linear Algebra)

Randil Pushpananda, PhD
rpn@ucsc.cmb.ac.lk



Column Way, Row Way, Matrix Way

Column way
Linear combination

$$c \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} + d \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

Row way
Two equations for c and d

$$v_1 c + w_1 d = b_1$$

$$v_2 c + w_2 d = b_2$$

Matrix way
2 by 2 matrix

$$\begin{bmatrix} v_1 & w_1 \\ v_2 & w_2 \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

Question 1

$$2x + y = 3$$

$$x - 2y = -1$$

- Find
- (i) Row Picture
 - (ii) Column Picture
 - (iii) Matrix Form

Question 2

$$2x - y = 0$$

$$-x + 2y = 3$$

- Find
- (i) Row Picture
 - (ii) Column Picture
 - (iii) Matrix Form

$$c \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} + d \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

If the points v and w and the zero vector 0 are not on the same line, there is exactly one solution c, d . Then the linear combinations of v and w exactly fill the xy plane. The vectors v and w are “linearly independent”. The 2 by 2 matrix $A = \begin{bmatrix} v & w \end{bmatrix}$ is “invertible”.

Zero Vector

Linearly Independent

Invertible

Zero Vector

- Vector that has a zero magnitude and no direction.
- Example:
 - Suppose two people are pulling a rope from its two ends with equal force but in opposite directions. So, the net force applied to the rope will be a zero vector (null vector) as the two equal forces balance each other out because they are in opposite directions

Linearly Independent

- Two or more vectors are said to be linearly independent if none of them can be written as a linear combination of the others.

$$\begin{bmatrix} 1 \\ 1 \\ 3 \\ 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ -1 \\ -3 \end{bmatrix} \begin{bmatrix} 2 \\ 5 \\ 9 \\ 6 \end{bmatrix} \xrightarrow{\text{linear combination}} c_1 \begin{bmatrix} 1 \\ 1 \\ 3 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} -1 \\ 1 \\ -1 \\ -3 \end{bmatrix} + c_3 \begin{bmatrix} 2 \\ 5 \\ 9 \\ 6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Invertible

- A square matrix such that the product of the matrix and its inverse generates the identity matrix.

$$AB = BA = I_n$$

$$\Rightarrow B = A^{-1}$$

Where,

A is $(n \times n)$ invertible matrix.

B is $(n \times n)$ matrix called inverse of A .

I_n is $(n \times n)$ identity matrix.

Vectors in 3 - Dimensions


$$\begin{array}{rcl} 2x & - & y & = & 0 \\ -x & + & 2y & - & z & = & -1 \\ & & -3y & + & 4z & = & 4 \end{array}$$

Find x, y and z

Vectors in 3 - Dimensions


$$\begin{array}{rcl} 2x & - & y & & = & 0 \\ -x & + & 2y & - & z & = & -1 \\ & & -3y & + & 4z & = & 4 \end{array}$$

Row Picture



Try: <https://www.desmos.com/3d>


$$-x + 2y - z = -1$$



$$-x + 2y - z = -1$$

$$2x - y = 0$$

☒ Extend to 3D




$$-x + 2y - z = -1$$

$$2x - y = 0$$

☒ Extend to 3D


$$-3y + 4z = 4$$



Vectors in 3 - Dimensions

$$\begin{array}{rcl} 2x & - & y & & = & 0 \\ -x & + & 2y & - & z & = & -1 \\ & & -3y & + & 4z & = & 4 \end{array}$$

Column Picture






Vectors in 3 - Dimensions

$$\begin{array}{rcl} 2x & - & y & = & 1 \\ -x & + & 2y & - & z & = & 1 \\ & & -3y & + & 4z & = & -3 \end{array}$$

Column Picture

$x = ?$, $y = ?$, $z = ?$





Matrix Form

$$\begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 4 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -3 \end{bmatrix}$$

$$A.x = b$$

Matrices and Their Column Spaces

1 $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$ is a **3 by 2 matrix**: $m = 3$ rows and $n = 2$ columns. Rank 2.

2 The 3 components of $A\mathbf{x}$ are dot products of the 3 rows of A with the vector \mathbf{x} :

$$\begin{array}{l} \text{Row at a time} \\ A \text{ times } \mathbf{x} \end{array} \quad \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \begin{bmatrix} 7 \\ 8 \end{bmatrix} = \begin{bmatrix} 1 \cdot 7 + 2 \cdot 8 \\ 3 \cdot 7 + 4 \cdot 8 \\ 5 \cdot 7 + 6 \cdot 8 \end{bmatrix} = \begin{bmatrix} 23 \\ 53 \\ 83 \end{bmatrix}.$$

3 $A\mathbf{x}$ is also a combination of the columns of A $\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \begin{bmatrix} 7 \\ 8 \end{bmatrix} = 7 \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + 8 \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}.$

4 The column space of A contains all combinations $A\mathbf{x} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2$ of the columns.

5 Rank one matrices: All columns of A (and all combinations $A\mathbf{x}$) are on **one line**.

Square Matrices and Types

- A typical matrix A is a rectangle of m times n numbers— m rows and n columns.
- If m equals n then A is a “square matrix”.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

**Identity
matrix**

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

**Diagonal
matrix**

$$\begin{bmatrix} 2 & 1 & -3 \\ 0 & 4 & 7 \\ 0 & 0 & 5 \end{bmatrix}$$

**Triangular
matrix**

$$\begin{bmatrix} 2 & 1 & -3 \\ 1 & 4 & 7 \\ -3 & 7 & 5 \end{bmatrix}$$

**Symmetric
matrix**

Two ways to the same answer

The row picture of $A\mathbf{x}$ will come from dot products of \mathbf{x} with the rows of A .

The column picture will come from linear combinations of the columns of A .

We often think of the columns of A as vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$. Each of those n vectors is in m -dimensional space. In this example the \mathbf{a} 's have $m = 3$ components each:

$m = 3$ rows
 $n = 4$ columns
3 by 4 matrix A

$$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 & \mathbf{a}_4 \end{bmatrix} = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix}$$

Row picture of $A\mathbf{x}$ Each row of A multiplies the column vector \mathbf{x} . Those multiplications *row times column* are dot products ! The first dot product comes from row 1 of A :

$$(\text{row } 1) \cdot \mathbf{x} = (-1, 1, 0, 0) \cdot (x_1, x_2, x_3, x_4) = \mathbf{x}_2 - \mathbf{x}_1.$$

It takes m times n small multiplications to find the $m = 3$ dot products that go into $A\mathbf{x}$.

Three rows
Three dot
products

$$A\mathbf{x} = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \text{row } 1 \cdot \mathbf{x} \\ \text{row } 2 \cdot \mathbf{x} \\ \text{row } 3 \cdot \mathbf{x} \end{bmatrix} = \begin{bmatrix} \mathbf{x}_2 - \mathbf{x}_1 \\ \mathbf{x}_3 - \mathbf{x}_2 \\ \mathbf{x}_4 - \mathbf{x}_3 \end{bmatrix} \quad (1)$$

Notice well that each row of A has the same number of components as the vector \mathbf{x} . Four columns of A multiply x_1 to x_4 . Otherwise multiplying $A\mathbf{x}$ would be impossible.

Column picture of $A\mathbf{x}$ The matrix A times the vector \mathbf{x} is a **combination of the columns of A** . The n columns are multiplied by the n numbers in \mathbf{x} . Then add those column vectors $x_1\mathbf{a}_1, \dots, x_n\mathbf{a}_n$ to find the same vector $A\mathbf{x}$ as in equation (1):

$$A\mathbf{x} = x_1(\text{column } \mathbf{a}_1) + x_2(\text{column } \mathbf{a}_2) + x_3(\text{column } \mathbf{a}_3) + x_4(\text{column } \mathbf{a}_4) \quad (2)$$

This combination of n columns involves exactly the same multiplications as dot products of \mathbf{x} with the m rows. But it is higher level! We have a vector equation instead of three dot products. You see the same result $A\mathbf{x}$ in equation (1) above and equation (3) below.

Combination of columns

$$A\mathbf{x} = x_1 \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} + x_4 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \mathbf{x}_2 - \mathbf{x}_1 \\ \mathbf{x}_3 - \mathbf{x}_2 \\ \mathbf{x}_4 - \mathbf{x}_3 \end{bmatrix} \quad (3)$$

Independence, Dependence, and Column space

That plane is the column space of this matrix : Plane = all combinations of the columns.

Dependent columns in Example 2

column 1 + column 2 − column 3 is $(0, 0, 0)$.

Example 3 $A_3 = \begin{bmatrix} 1 & 3 & 4 \\ 2 & 6 & 8 \\ 5 & 15 & 20 \end{bmatrix}$

Now \mathbf{a}_2 is 3 times \mathbf{a}_1 . And \mathbf{a}_3 is 4 times \mathbf{a}_1 .

Every pair of columns is dependent.