

JEE MAIN

MODERN PHYSICS - PART 5

FORMULAE

NUCLEAR PHYSICS

Now that's how you REVISE

-Mohit Goenka, IIT Kharagpur

List of Content on Eduniti YouTube Channel:

1. PYQs Video Solution Topic Wise:
(a) JEE Main 2018/2020/2021 Feb & March
2. Rank Booster Problems for JEE Main
3. Part Test Series for JEE Main
4. JEE Advanced Problem Solving Series
5. Short Concept Videos
6. Tips and Tricks Videos
7. JEE Advanced PYQs

.....and many more to come



Eduniti for Physics

MODERN PHYSICS

ATOMIC
PHYSICS
PART 1

PHOTOELECTRIC
EFFECT
PART 2

DUAL
NATURE
OF
LIGHT
PART 3



X-RAYS

RADIOACTIVITY
PART 4


NUCLEAR
PHYSICS
PART 5



1. HOW TO WRITE A NUCLEAR REACTION



2. NUCLEAR FORCE



Strong attractive short range force.

$$F_{nn} = F_{np} = F_{pp}$$

(i) F_e : repulsive electrostatic force

F_n : net attractive nuclear force

$$F_e = F_n \Rightarrow \text{Stable Nucleus}$$



3. NUCLEUS SIZE AND STABILITY OF HEAVY NUCLEUS

↳ Size of Nucleus \propto Atomic mass

$$\Rightarrow \frac{4}{3}\pi R^3 \propto A \Rightarrow \boxed{R = R_0 A^{1/3}}$$

fermi-const.
 $R_0 \sim 10^{-15} \text{ m}$

↳ If $R \uparrow \Rightarrow F_n \downarrow$

So, Nucleus gets unstable
 \Rightarrow Decay starts



NOTE: When reactants combine to form stable product, THERE IS MASS LOSS called "MASS DEFECT"

$$\boxed{\Delta m = Zm_p + (A-Z)m_n - M_X}$$


and, $E_b = \Delta m c^2 \begin{cases} \Delta m \text{ if in AMU,} \\ 1 \text{ AMU} = 1.66 \times 10^{-27} \text{ kg} \end{cases}$

or

$$\boxed{E_b = \Delta m (\text{in AMU}) \times 931.5 \text{ MeV}}$$

4. NUCLEAR BINDING ENERGY

the **energy required** to separate an atomic nucleus completely into its constituent protons and neutrons, or, equivalently, the **energy that would be liberated** by combining individual protons and neutrons into a single nucleus.




5. BINDING ENERGY PER NUCLEON


\hookrightarrow $\begin{matrix} A \\ Z \end{matrix} X$ A is no of nucleons
 NOTE: E_b is lower for $A < 30$ and $A > 170$

$$BE/\text{nucleon} = \frac{E_b}{A}$$

\downarrow
 tells how stable is a nucleus



6. NUCLEAR FISSION AND FUSION




- (i) In both Energy released
- (ii) $\Delta E_1 = (m_A + m_B - m_C) c^2$
- $\Delta E_2 = (m_X - m_Y - m_Z) c^2$

* NOTE: Energy released or even supplied is called Q-value
 (+VE) (-VE)

$$Q = \Delta m c^2$$

7. Alpha, BETA AND GAMMA DECAY



NOTE: This energy is released in form of Kinetic energy.



$$M_Y v_Y = M_\alpha v_\alpha \quad \text{--- (i)}, \quad Q = \frac{1}{2} M_Y v_Y^2 + \underbrace{\frac{1}{2} M_\alpha v_\alpha^2}_{K_\alpha} \quad \text{--- (ii)}$$

from (i) and (ii) :



$$K_\alpha = \frac{Q M_Y}{M_\alpha + M_Y} = \boxed{\frac{Q(A-4)}{A}}$$



(b) Beta Decay (e^- or e^+)
 e^- → electron
 e^+ → positron



ν → neutrino
 $\bar{\nu}$ → antineutrino

β^+ Decay




$$\Delta m = M_X - M_Y - 2m_e$$

β^- Decay



$$\Delta m = M_X - M_Y$$

K capture



$$\Delta m = M_X - M_Y$$

(In K capture, e^- is captured from K-shell)

(c) Gamma Decay (EM Radiation)



↓
Excited state



NOTE: It can happen even for β -decay