

JEE MAIN

MODERN PHYSICS – PART 6

FORMULAE

X - RAYS

Now that's how you REVISE

-Mohit Goenka, IIT Kharagpur

List of Content on Eduniti YouTube Channel:

1. PYQs Video Solution Topic Wise:
(a) JEE Main 2018/2020/2021 Feb & March
2. Rank Booster Problems for JEE Main
3. Part Test Series for JEE Main
4. JEE Advanced Problem Solving Series
5. Short Concept Videos
6. Tips and Tricks Videos
7. JEE Advanced PYQs
8. Formulae Revision Series

.....and many more to come



Eduniti for Physics

MODERN PHYSICS

ATOMIC
PHYSICS
PART 1

PHOTOELECTRIC
EFFECT
PART 2

DUAL
NATURE
OF
LIGHT
PART 3

X-RAYS
PART 6

RADIOACTIVITY
PART 4

NUCLEAR
PHYSICS
PART 5



TOPICS COVERED

1. Soft and Hard X – Rays
2. Production of X – Rays : Coolidge Tube
3. Continuous X – Ray Production
4. Characteristic X – Ray Production
5. Complete Spectrum
6. Moseley's Law



1. X-RAYS ($\approx 1 \text{ \AA}$)


Soft X-RAYS

- High wavelength
- Low energy

Hard X RAYS

- Low wavelengths
- High Energy

2. PRODUCTION OF X-RAYS (X-ray tubes)



X-Rays are produced by incidence of accelerated e^- on target material.



Continuous X-Rays
(Bremsstrahlung)

Characteristic X-Rays

3. **CONTINUOUS X-RAY** This Phenomena is called Bremsstrahlung
 ↳ Deceleration of e^- when deflected by atomic nucleus causes production of X-Rays.

(a) Energy of X-rays, $E = E_1 - E_2$
 $E_{\max} = E_1 = eV$ ($E_2 = 0$)

∴ Cutoff Wavelength of X-Ray,
 $\lambda_{\min} = \frac{hc}{E_{\max}} = \frac{hc}{eV} = \frac{12431}{V} \text{ \AA}$



→ X-Rays Continuum Radiation Spectra




4. CHARACTERISTIC X-RAY

- (i) some incident e^- knocks off orbital e^- of $K, L, M \dots$ shell.
- (ii) If $eV > \text{Binding Energy of "K shell } e^-"$, only then it is removed.
- (iii) e^- from $L, M, N \dots$ can jump to K and during this **photon (x-ray)** is emitted.

$$\lambda = \frac{hc}{\Delta E}$$

- (a) K_α x-Ray \rightarrow If e^- jumps from $L \rightarrow K$
- (b) K_β \rightarrow e^- jumps from $M \rightarrow K$
- (c) K_γ \rightarrow e^- jumps from $N \rightarrow K$

} K-series




NOTE:


(i) If you compare

K_β and K_α

$\lambda_{K_\beta} < \lambda_{K_\alpha} (\because \Delta E_{MK} > \Delta E_{LK})$



5. COMPLETE SPECTRUM



$K_{\beta} : M \rightarrow K$
 $K_{\alpha} : N \rightarrow K$
 $L_{\beta} : N \rightarrow L$
 $L_{\alpha} : M \rightarrow L$

6. MOSELEY'S LAW ($\sqrt{\nu} = a(z - \sigma)$)

λ of characteristic X-Rays :

$$\frac{1}{\lambda} = R(z - \sigma)^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$\underbrace{z - \sigma}_{\text{effective atomic no}}$

$$\therefore \nu = \frac{c}{\lambda}$$

$$\Rightarrow \nu = RC(z - \sigma)^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

} For K-Series
 $\sigma = 1$
 $R : 10^7 \text{ m}^{-1}$
 \hookrightarrow Rydbergs Const.

$$\therefore \sqrt{\nu} = a(z - \sigma)$$

$$\sqrt{RC \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)}$$

\hookrightarrow Square root of frequency is linearly proportional to Atomic number.

